CASE REPORT

Autoimmune haemolytic anaemia in a newborn infant

M Motta, A Cavazza, C Migliori, G Chirico

Arch Dis Child Fetal Neonatal Ed 2003;88:F341-F342

The case is reported of an infant with autoimmune haemolytic anaemia of perinatal onset. Combined treatment with steroids and cyclosporin was necessary to improve haemolysis and reduce the high transfusion requirements. Treatment was discontinued at 13 months of age. The child was healthy at the follow up at 24 and 36 months of age.

Autoimmune haemolytic anaemia (AIHA) is a rare disease; the annual incidence is about one case in 80 000 live births. It is more common than acquired aplastic anaemia and less common than immune thrombocytopenic purpura.¹ Infants of any age can be affected, although neonatal occurrence is considered exceptional. We describe a patient with severe haemolytic anaemia of perinatal onset.

CASE REPORT

The male infant was born by normal vaginal delivery at term with a birth weight of 3120 g. Apgar scores were 9 at one and five minutes. The direct antiglobulin test (DAT) on cord blood was slightly positive (titre 1:64). Twelve hours after birth, he was transfused with packed red cells because of severe anaemia. Two weeks later, he needed another blood transfusion, and at 23 days of life he was transferred to our hospital because of the persistence of anaemia. His family history showed no significant disease; his parents were non-consanguineous. The mother's blood type was A Rh positive, and the infant's was O positive.

Physical examination showed pallor, hepatosplenomegaly, and no lymphadenomegaly.

Haemoglobin concentration was 59 g/l (17% packed cell volume; red blood cells $1870 \times 10^9/l$), reticulocytes were absent, platelets were $314 \times 10^9/l$. Leucocyte count was $8.6 \times 10^9/l$ (neutrophils 54%, lymphocytes 36%, and monocytes 6%). A blood smear showed anisopoikilocytosis and spherocytosis.

DAT was positive (titre 1:250). Serum lactate dehydrogenase concentration was 1314 U/l.

Clotting tests (prothrombin time, partial thromboplastin time, fibrinogen) were normal, and the autoantibody assay (anti-nuclear, anti-extractable nuclear antigens, anti-DNA, anti-cardiolipin) was negative. Investigations for HIV, Epstein-Barr virus, cytomegalovirus, herpes simplex virus 1-2, and hepatitis A, B, and C viruses did not indicate a congenital infection. Serology for parvovirus showed positivity of IgG and negativity of IgM in both mother and infant.

A tibial bone marrow aspiration showed normal cellularity, the presence of megakaryocytes, and normal granulocytic and lymphocytic lineages. A paucity of erythroid precursors, blocked at the stage of basophil erythroblast, was observed. No abnormal blast cells or malignant cytological abnormalities were found.

Cultures of peripheral blood cells, obtained by separation of mononuclear cells and stimulated with several cytokines (granulocyte colony stimulating factor, erythropoietin, interleukin 3), showed normal growth of haematopoietic progenitor cells.²

Figure 1 Number of transfusions (indicated by arrows) and schedule of treatment related to packed cell volume and reticulocyte counts, during the first 15 months of life in an infant with autoimmune haemolytic anaemia.

Lymphocyte phenotype was characterised by a reduction in total T cells, with inversion of T4/T8 ratio and relative increase in T activated lymphocytes and B cells: CD3, 33%; CD3⁺/DR⁺, 18%; CD4, 16%; CD8, 18%; CD5, 50%; HLA-DR, 58%; CD19, 24%; CD16, 30%; TCRγδ⁺/CD4⁺/CD8⁺, 1%; CD95, 78%.

Serial measurements of polyspecific DAT showed a stable positivity with titre 1:256 up to first 8 weeks of age. These findings, along with a negative indirect Coombs test and irregular antibody test in the mother, suggested AIHA.

A monospecific anti-human globulin test was positive for anti-IgG, and negative for anti-IgA, anti-IgM, anti-C3c, and anti-C3d.

Immunosuppressive treatment with methylprednisolone at a dose of 2 mg/kg/day was started. After four weeks, because of the persistence of significant haemolysis (high transfusion

Abbreviations: AIHA, autoimmune haemolytic anaemia; DAT, direct antiglobulin test

Table 1 Reported cases of young infants with autoimmune haemolytic anaemia (AIHA)

Reference	Age at onset (weeks)	Diagnosis	Treatment	Outcome
Win <i>et al</i> ¹	12	AIHA	Not reported	Favourable
Satake <i>et al</i> ²	8	AIHA AI-enteropathy	Steroids immunoglobulin cyclosporin A	Resolution of anaemia
Satake <i>et al</i> ²	4	AIHA AI-enteropathy	Steroids	Improvement of anaemia; died at 6 months from severe diarrhoea
Ritz & Haber ³	6	AIHA	Steroids, ACTH	Resolution of anaemia
Hadnagy ⁴	At birth	AIHA	Steroids, splenectomy	No improvement
Laskj <i>et al</i> ⁵	10	AIHA	—	Favourable
Gasser & Hollander ⁶	7	AIHA	Steroids, splenectomy	Died at 4 months

ACTH, Adrenocorticotrophic hormone; AI, autoimmune.

requirements, increased lactate dehydrogenase, positive DAT, positive cross matching test) and inhibition of bone marrow erythropoiesis (reticulocytopenia), two courses of high dose methylprednisolone (20 mg/kg/day for three consecutive days) followed by intravenous immunoglobulin (2 g/kg in two doses) were given; erythropoiesis was recovered. Because of persisting haemolysis, a maintenance treatment with steroid and 10 mg/kg/day cyclosporin was given, with a significant reduction in the transfusion requirement after two weeks (fig 1).

No appreciable side effects of cyclosporin were recorded. The last transfusion was at the age of 6 months; cortisone tapering was started two months later. Cyclosporin treatment was suspended at the age of 11 months, and treatment was finally discontinued at 13 months of age, when the DAT gave a negative result for the first time.

As a consequence of steroid treatment, the child initially showed hyposomia (length < 3rd centile). However, clinical evaluation at 24 and 36 months of age showed that length and growth had been regained (> 25th centile), and there was no recurrence of haemolysis.

DISCUSSION

To our knowledge, only seven young infants with AIHA have so far been reported (table 1). Six responded to treatment with improvement of anaemia, although for only two could treatment be discontinued without relapse. One of these, who was also affected by autoimmune enteropathy, died at 6 months of age with severe watery diarrhoea. One child remained dependent on steroids, and underwent splenectomy. One baby died as a consequence of refractory AIHA.

During the first 6 weeks of life, an infant's ability to form antibodies is limited by an immature immune and reticulo-endothelial system. Early immunological stimulation usually results in the production of low titre IgM specific antibodies, with a delayed switch from the IgM to the IgG class. However, the impaired responsiveness of the neonate is variable and the possibility of synthesis of autoimmune antibodies cannot be excluded.

Neonatal occurrence of AIHA is very rare. In only one of the reported patients⁴ was the diagnosis made at birth as in our case. As AIHA is so rare, few data on clinical management and treatment strategies are available. The use of steroids is widely accepted, although long term treatment during infancy is associated with appreciable side effects, such as neurological and somatic growth retardation, hypertension, and hypertrophic cardiomyopathy.

Tapering of steroid treatment should be considered when DAT becomes negative and the patient's haemoglobin concentration and reticulocyte count remain satisfactory. Other immunosuppressive drugs such as cyclosporin, azathioprine, and cyclophosphamide are not often used in children, although they can be useful for refractory forms because the long term use of high dose steroids is undesirable.⁹

Splenectomy is sometimes considered for children with chronic AIHA, although it should be avoided in young infants because of the high risk of sepsis and mortality.

Transfusion may be complicated by the high requirement and the difficulty of obtaining compatible blood.¹⁰ When the antigenic specificity is panreactive, the cross match may well not identify any compatible blood units. In such cases, the blood bank designates certain units that are considered the "least incompatible" with the patient's serum.

In our case the use of high dose steroid and intravenous immunoglobulin successfully treated the hypoplastic phase, and the use of cyclosporin controlled the disease (fig 1).

Authors' affiliations

M Motta, A Cavazza, C Migliori, G Chirico, Division of Neonatology and Neonatal Intensive Care, Spedali Civili, Brescia, Italy

Correspondence to: Dr Chirico, Divisione di Neonatologia, Spedali-Civili, 25123 Brescia, Italy; gaechir@tin.it

Accepted 21 September 2002

REFERENCES

- Ware RE, Rosse WF. Autoimmune hemolytic anemia. In: Nathan DG, Orkin SH, eds. *Nathan and Oski's hematology of infancy and childhood*. 5th ed. Philadelphia: WB Saunders, 1997:499-522.
- Gasparoni A, Ciardelli L, Avanzini MA, *et al*. Immunophenotypic changes of fetal cord blood hematopoietic progenitor cells during gestation. *Pediatr Res* 2000;**47**:825-9.
- Win N, Koye T, Mir N, *et al*. Autoimmune haemolytic anemia in infancy with anti-Kpb specificity. *Vox Sang* 1996;**71**:187-8.
- Satake N, Nakanishi M, Okano M, *et al*. A Japanese family of X-linked auto-immune enteropathy with haemolytic anemia and polyendocrinopathy. *Eur J Pediatr* 1993;**152**:313-15.
- Ritz ND, Haber A. Autoimmune hemolytic anemia in a 6-week-old child. *J Pediatr* 1962;**61**:904-10.
- Hadnagy CS. Severe chronic autoimmune hemolytic anemia presenting hemolytic disease of the newborn. *Lancet* 1989;**2**:749.
- Lasky B, Wake EJ, Bain HW, *et al*. Autohemolytic anemia in young infants. *J Pediatr* 1961;**59**:42.
- Gasser C, Hollander L. Acquired acute haemolytic anemia. *Revue d'Hematologie* 1951;**6**:316.
- Rackoff WR, Manno CS. Treatment of refractory Evans syndrome with alternate-day Cyclosporin and Prednisone. *Am J Pediatr Hematol Oncol* 1994;**16**:156-9.
- Jefferies LC. Transfusion therapy in autoimmune hemolytic anemia. *Hematol Oncol Clin North Am* 1994;**8**:1087-104.