

Introduction There is some controversy in the literature regarding the outcomes of pregnancies complicated by antepartum bleeding of unknown origin (ABUO).

Objective To explore the risk of adverse maternal and perinatal outcomes in women with ABUO occurring after the first trimester of pregnancy.

Methods Cohort study based on data extracted from the Aberdeen Maternity and Neonatal Databank. The study population was all primigravidae delivering in Aberdeen Maternity Hospital between 1976 and 2010. Exposure was antepartum haemorrhage occurring after the first trimester not attributable to placenta praevia or abruption. Data were analysed using univariate and multivariate statistical methods.

Results Between 1976 and 2010, there were 7,517 women with ABUO and 68,423 women without. Women with ABUO were more likely to be smokers, belong to lower social class and have slightly higher body mass index. Multivariate analysis revealed that non-specific APH was a significant risk factor for induced labour (OR = 1.23, 95% CI = 1.16, 1.31), preterm delivery (OR = 2.30, 95% CI = 2.11, 2.50), postpartum haemorrhage (OR = 1.15, 95% CI = 1.06, 1.25), Apgar score less than 7 at 1 minute (OR = 1.12, 95% CI = 1.05, 1.21), and at 5 minutes (OR = 1.25, 95% CI = 1.04, 1.50). There was no significant association detected with preeclampsia, mode of delivery or perinatal death.

Conclusion Pregnancies complicated by ABUO are at greater risk of delivery related and neonatal adverse outcomes attributable to preterm birth, some of which is iatrogenic.

PP23 MEASUREMENTS OF AMNIOTIC FLUID: ASSOCIATION AND PREDICTION OF SMALL FOR GESTATIONAL AGE AND COMPROMISE OF FETAL WELLBEING: A SYSTEMATIC REVIEW AND META-ANALYSIS

doi:10.1136/archdischild-2013-303966.303

^{1,2}RK Morris, ^{2,3}CH Meller, ^{1,4}JA Tamblin, ⁵GM Malin, ⁸RD Riley, ^{1,2,8}MD Kilby, ⁶SC Robson, ⁷KS Khan. ¹Division of Reproduction, Genes and Development, University of Birmingham, Birmingham, UK; ²Fetal Medicine Centre, Birmingham Women's Hospital, Birmingham, UK; ³Obstetrics Division, Hospital Italiano de Buenos Aires, Buenos Aires, Argentina; ⁴University Hospital North Staffordshire, Stoke-On-Trent, UK; ⁵School of Clinical Sciences, University of Nottingham, Nottingham, UK; ⁶Institute of Cellular Medicine, Newcastle University, Newcastle, UK; ⁷Women's Health Research Unit, The Blizard Institute, Barts and The London School of Medicine, Queen Mary, University of London, UK; ⁸School of Health and Population Sciences, University of Birmingham, Birmingham, UK

Objective Evaluate the strength of association and predictive value of measurements of amniotic fluid volume (AFV) for small for gestational age and compromise of fetal wellbeing.

Design Systematic literature review with random effects meta-analysis to compute summary odds ratios (OR) to assess prognostic association and assess predictive ability with sensitivity, specificity and likelihood ratios. Study characteristics, design, methodological and reporting quality were objectively assessed.

Data Sources and Eligibility Systematic search (inception to October 2011) for studies comparing AFV measures and outcomes of fetal size or wellbeing.

Results 43 studies reporting 244,493 fetuses included. Strong associations between oligohydramnios (heterogenic definition) and birth weight < 10th centile [OR 6.31 (4.15–9.58)] in a high risk population (6 studies, 28510 fetuses), and mortality [Neonatal death OR 8.72 (2.43–31.26) 6 studies, 55735 fetuses, and perinatal mortality in a high risk population OR 11.54 (4.05–32.9) 2 studies, 27891 fetuses] were identified. There was no significant association between oligohydramnios and abnormal cord pH or adverse perinatal outcome.

There was no significant association between polyhydramnios (heterogenic definitions) and poor fetal growth [Birth weight < 10th centile OR 0.37 (0.07–1.95)]. A strong association between polyhydramnios (maximum pool depth > 8 cm or amniotic fluid index ≥ 25 cm) and birth weight > 90th centile [OR 11.41 (7.09–18.36) 1 study, 3960 fetuses] was found. Despite strong associations predictive ability was poor with significant heterogeneity despite subgroup analysis.

Conclusion Oligohydramnios is associated with poor fetal growth and mortality. Polyhydramnios is associated with BW > 90th centile. Despite strong associations, overall predictive ability was poor. MPD accuracy was slightly improved over AFI and is the recommended technique.

PP24 PLACENTAL VOLUME CAN BE ACCURATELY MEASURED USING TWO- AND THREE-DIMENSIONAL ULTRASOUND NEAR TERM

doi:10.1136/archdischild-2013-303966.304

^{1,2}LE Higgins, ¹CP Sibley, ^{1,2}AEP Heazell, ^{1,2}ED Johnstone. ¹Maternal and Fetal Health Research Centre, University of Manchester, Manchester, UK; ²St. Mary's Hospital, Central Manchester University Hospitals NHS Trust, Manchester, UK

Introduction Stillbirth and fetal growth restriction are associated with lower placental weights compared to live, appropriately grown infants. Sonographic measures of placental size have been used to predict later pregnancy complications but have not been related back to true placental size. We aimed to validate measures of placental size in the third trimester of pregnancy.

Methods Ultrasonographic placental examination was performed in 30 singleton pregnancies of 35–42 weeks' gestation within seven days of delivery. Placental length, width and depth were measured using two-dimensional (2D) ultrasound, placental volume was estimated by a novel elliptical model, Kliman placental gas gauge and two three-dimensional (3D) ultrasound techniques (rotational (VOCAL) and slicing methods). Following delivery, the placenta was measured, weighed and volume determined. Sonographic and true placental measures were compared using linear regression.

Results Elliptical (but not Kliman placental gas gauge) 2D and all 3D ultrasound estimates of placental volume each related significantly with true placental volume with similar accuracy (Table 1) but not to placental weight ($p > 0.05$). VOCAL demonstrated higher accuracy than the elliptical model at the smallest placental volumes.

Abstract PP24 Table 1

	r ²	p
Ellipse	0.27	0.0033
Kliman placental gas gauge	0.054	0.21
VOCAL		
30°	0.31	0.0018
15°	0.67	0.0042
Slicing		
10 mm	0.24	0.0064
5 mm	0.29	0.0025

Conclusion True placental volume can be accurately predicted using 2D and 3D ultrasound in the third trimester. 3D ultrasound may improve detection of the abnormally small placenta. Further research is required to establish whether this measurement can predict poor pregnancy outcomes related to placental disease.

PP25 INTERPREGNANCY CHANGES IN MATERNAL WEIGHT AND BODY MASS INDEX

doi:10.1136/archdischild-2013-303966.305

DA Crosby, M Collins, V O'Dwyer, A O'Higgins, N Farah, MJ Turner. Coombe Women and Infant's University Hospital, Dublin, Ireland

Objective This longitudinal study examined changes in maternal weight and Body Mass Index (BMI) in the early pregnancy between a woman's first and second baby.

Methods We studied women more than 18 years old with a singleton pregnancy who delivered their first baby weighing ≥ 500 grammes in 2009 and who re-attended for antenatal care with a subsequent ongoing pregnancy before January 1st 2012. Maternal weight and height were measured accurately before 18 weeks gestation in both pregnancies and BMI was calculated.

Results Of the 3284 primigravidae, the mean weight at the first antenatal visit was 66.4 kg (SD 12.7). The mean BMI was 24.5 kg/m² (SD 4.6), and 11.3% (n = 370) were obese. Of these 3284 women, 1220 (37.1%) re-attended for antenatal care before 2012 after confirmation of an ongoing pregnancy. Of the 1220 women who re-attended, 788 (64.6%) had gained weight (mean 4.6 kg SD 3.9), 402 (33.0%) had lost weight (mean 3.0 kgs SD 2.9) and 30 (2.4%) had maintained their weight. As a result, 20.2% (n = 247) were now in a higher BMI category and 4.8% (n = 58) had become obese; 5.8% (n = 71) were in a lower BMI category and 1.2% (n = 15) were no longer obese. These early pregnancy weight changes were influenced by maternal age, but not by the duration of the interpregnancy interval.

Conclusion As two thirds of women gain weight in the short-term after delivery of their first baby, we recommend that the advice women get before and during pregnancy needs to be reinforced postpartum.

PP26 SUSTAINABLE VOLUNTEERING AND ITS EFFECT ON UGANDAN STILLBIRTH RATES


doi:10.1136/archdischild-2013-303966.306

¹K MacLeod, ²L Ackers. ¹Mersey Deanery, Liverpool, UK; ²University of Liverpool, Liverpool, UK

Background The Ugandan Maternal and Newborn Hub is a THET funded initiative focused on implementing Sustainable Volunteering Projects (SVPs) to improve maternal and newborn health. Annually, there are 2.65 million stillbirths worldwide and almost 90% occur in low-resource settings. Up to 70% of stillbirths occur in the intrapartum period; frequently these are associated with suboptimal care.

Methods Through a SVP, from January to June 2012, we undertook a quality improvement project aiming to reduce intrapartum stillbirths at Mbarara Hospital in Uganda. Our quality improvements included regular audit presentations, implementation of a labour ward board and skills and drills training. We classified suboptimal care into types of delay using Thaddeus & Maine Three Delay Model.

Results In this 6 months period, there were 102 intrapartum stillbirths. Most stillbirths were singleton pregnancies (99%) with a mean maternal age of 24 years and a mean parity of 2. 34% of women were unbooked. Suboptimal factors contributing to



Abstract PP26 Table Stillbirths Figures Mbarara Hospital 2012

intrapartum stillbirths were identified in 86.4% of cases, 22% were related to care whilst admitted. Intrapartum stillbirths occurring after hospital admission reduced after introduction of quality improvement measures: see table.

Discussion Basic and comprehensive emergency obstetric care is an important intervention to reduce intrapartum stillbirths in the developing world. SVPs are an effective way of individualising quality improvement measures and improving outcomes.

PP27 MICRO-VASCULAR DISEASE AT BOOKING IN T1DM AND ASSOCIATED RISK OF DEVELOPING PRE-ECLAMPSIA

doi:10.1136/archdischild-2013-303966.307

¹N Patel, ²A Brackenridge, ³A Kanji, ⁴D Pasupathy, ⁴D Rajasingam. ¹Division of Women's Health, Women's Health Academic Centre, King's College London, King's Health Partners, London, UK; ²Department of Diabetes, Guy's and St Thomas' NHS Foundation Trust, London, UK; ³School of Medicine, King's College London, London, UK; ⁴Division of Women's Services, Guys and St Thomas' NHS Foundation Trust, London, UK

Background and Aims Micro-vascular disease affects the majority of T1DM patients and is directly related to both duration and glycemic control of diabetes. The aim of this study is to identify the association of micro-vascular disease at booking and subsequent risk of developing pre-eclampsia during pregnancy.

Methods Retrospective cohort study of women with T1DM (n = 37) on continuous subcutaneous insulin infusions booked at Guy's & St Thomas' Hospital between November 2010 and April 2012. Micro-vascular disease at booking was, defined as nephropathy or/and retinopathy at booking.

Results The prevalence of micro-vascular disease at booking was 54.1% (n = 20). No significant difference were seen in age (33.5 years vs 35.0 years; p = 0.28), HbA1c (7.5 vs 6.9; p = 0.11), length of diabetes (16.0 years versus 23.0 years; p = 0.14) compared to women without micro-vascular disease.

Presence of micro-vascular disease was significantly associated with the pre-eclampsia (55.0% vs 17.6%; p = 0.02; unadjusted OR 5.70; CI 1.24–26.26). This association was not explained by maternal age, duration or control of diabetes, parity, BMI and maternal hypertension (adjusted OR 4.92; CI 0.7–32.3).

The rate of LGA and SGA was 51.3% and 5.4% respectively. Women with micro-vascular disease had higher rates of SGA (10.0% vs 0.0%; p < 0.05) and LGA (65.0% vs 35.0%; p < 0.05).

PP28 CERVICAL CERCLAGE: ARE ALL TECHNIQUES SUCCESSFUL?

doi:10.1136/archdischild-2013-303966.308

¹EA Bonney, ²A Atkinson, ²R Lancaster, ¹JJ Walker, ¹NAB Simpson. ¹Division of Clinical Sciences, Leeds Institute of Molecular Medicine, University of Leeds, Leeds, UK; ²School of Medicine, University of Leeds, Leeds, UK

Objective Preterm birth is a global public health issue. In women considered high risk, insertion of a cervical cerclage has been shown to reduce this risk. We present findings from a retrospective cohort evaluating the success of different cerclage procedures (Shirodkar, McDonald and Transabdominal) in a tertiary level obstetric unit.

Study design Retrospective data was collected for 200 women who underwent a cerclage procedure at Leeds Teaching Hospitals NHS Trust between August 2000 and October 2010. Exclusion criteria for the study included multiple pregnancy, insertion of more than one cerclage in a single pregnancy, or an incomplete data record. Success was measured by delivery of a live baby ≥ 34 weeks. Mean gestational age (MGA) for each group was also calculated. Statistical analysis was performed using Fisher's exact test.