CURRENT TOPICS

The pharmacology of inhaled nitric oxide

A D Edwards

In 1980 Furchgott and Zawadski reported that endothelial cells stimulated by acetylcholine produced a vasodilator substance which relaxed vascular smooth muscle.¹ The identity of this endothelium derived relaxing factor (EDRF) remained elusive until 1987 when Palmer and colleagues showed that the effects of nitric oxide (NO) accounted for the known biological actions of EDRF,² and that endothelial cells synthesised NO from L-arginine and oxygen.³

The demonstration that analogues of L-arginine, such as N^G-monomethyl-L-arginine (L-NMMA) and N^G-nitro-L-arginine methyl ester (L-NAME), prevented the production of NO by cells provided an experimental tool for examining the widespread biological roles of NO.⁴ Administration of L-NMMA or L-NAME to animals blocked the vasodilator action of acetylcholine, caused a persistent rise in blood pressure, and reduced the blood volume of organs such as the brain, demonstrating both that NO release contributes to a constant vasodilator tone, and that further release can be initiated by appropriate stimuli.^{5,6} Conversely, administration of drugs which produced NO, such as the nitrate vasodilators, produced a fall in blood pressure.⁵

Several isoforms of the enzyme nitric oxide synthase (NOS) have been characterised – both constitutive and inducible. Constitutive enzymes were found in many tissues, including endothelium and brain.^{5,7} An inducible form of the enzyme, which was expressed after cytokine stimulation and during sepsis, was found in macrophages.^{5,8} NO is now known to be a multifunctional biological mediator with diverse roles in the cardiovascular, neurological, immunological and many other systems.⁹

Agonists and antagonists of NO have many potential medicinal roles, but among the first to be examined in detail is the use of inhaled NO as a specific pulmonary vasodilator. Information about this novel drug is accumulating, although there remain extensive gaps in our knowledge of its pharmacology and toxicology.

Chemistry

Nitric oxide (nitrogen monoxide) is a volatile gas which is the second (nitrogen oxidation state +2) in the sequence of oxy-compounds of nitrogen that includes NO⁻ (oxidation state

+1) and NO⁺ (oxidation state +3). The characteristic chemistry of NO is largely due to the arrangement of the electrons in the valence shell. Nitrogen and oxygen combine to form four bonding and four antibonding orbitals which are filled by 11 electrons, giving the molecule eight bonding and three antibonding electrons, a bond order of 2.5, and a bond length intermediate between double and triple bonding (1.150 Å).¹⁰

The single unpaired electron in the 2* π molecular orbital defines the molecule as a free radical (the presence of this unpaired electron is sometimes emphasised by writing the formula as $\cdot$ NO¹⁰). However, NO does not exhibit the high reactivity characteristic of most free radicals and shows little tendency to dimerise. Nevertheless, in the presence of superoxide radical reactions occur which lead to highly reactive species being formed¹¹; these are described further below.

NO is thermodynamically unstable ($\Delta G^{\circ}_{\text{formation}} = 86.32$ kJ), but decomposition is kinetically hindered and the gas can be stored indefinitely at room temperature and 1 atmosphere pressure. At increased pressure, disproportionation of NO to N₂O and NO₂ can occur. This reaction obeys third order kinetics and at 30°C and 200 atmospheres pressure, some NO initially present may be converted into N₂O and NO₂. This needs to be considered when NO is stored in pressurised cylinders for clinical use.¹⁰

In aqueous solutions NO exhibits solubility and diffusibility similar to other diatomic gases. It does not interact with water, and this property is probably relevant to its pharmacological and biological effects. The diffusion coefficient of NO in tissue and the lipid membrane permeability have not yet been formally quantified.¹¹ NO is not affected by light.

The unpaired electron is antibonding and easily detached to allow NO⁺ to be formed. Although the lifetime of NO⁺ in aqueous media is very short, it has been suggested that the presence in tissue of alternative redox forms of NO might lead to potent changes in biological effects.¹²

Metabolism

Both the effects and metabolism of inhaled NO take place in the lung, where several of the chemical reactions of NO are relevant:

Department of
Paediatrics and
Neonatal Medicine,
Royal Postgraduate
Medical School,
Hammersmith and
Queen Charlotte's
Hospitals, London
W12 0NN
A D Edwards

Correspondence to:
Professor A D Edwards.

REACTIONS WITH HAEM CONTAINING METALLOPROTEINS

Reactions with haem containing proteins are rapid. The association rate constant with Fe(II) in haemoglobin is about 300 times greater than for oxygen. This reaction leads to the formation of Fe(III) haemoglobin (methaemoglobin) and NO_2^- . NO also binds to Fe(III) complexes and Fe(III)NO undergoes a transfer reaction to form Fe(II)NO⁺. NO⁺ can be oxidised to NO_2^- and NO_3^- so that NO is cleared rapidly from the blood.¹¹ The reaction of NO with haem groups is thought to be the predominant interaction in biological systems, and to account both for its activity in smooth muscle cells and its rapid inactivation in blood.¹¹

FREE RADICAL REACTIONS

NO reacts with the superoxide anion (O_2^-), leading to the production of peroxynitrite (OONO^-). At physiological pH, OONO^- has a half life of 1.9 seconds, being protonated to form ONOOH which undergoes spontaneous homolysis to form the highly reactive hydroxyl radical ($\cdot\text{OH}$).¹³ The hydroxyl radical is highly damaging to lipid membranes, and consequently the potential for NO administration to induce tissue injury has led to a great deal of interest in this reaction, and to the role of NO as a cell toxin.¹⁴

AUTOXIDATION

The reaction between NO and O_2 to form NO_2 is rapid at high concentrations of NO, but slow at the concentrations used in trials of inhaled NO treatment. In air, for low concentrations of NO the half life of NO and the rate of NO_2 formation depend on the initial concentration; the half life may vary between one and 500 seconds.¹⁰ It is reported that at NO concentrations of 40 parts per million (ppm) or less, combination of NO with pure oxygen requires 2.56 minutes to yield 5 ppm NO_2 , although at 120 ppm NO this time falls to 0.26 minutes.¹⁵ NO_2 is an environmental toxin which can cause pulmonary oedema, haemorrhage, and bronchiolitis obliterans,¹⁶ and inhalation of NO_2 at concentrations as low as 2 ppm caused lung epithelial damage. A considerable body of published data exists on the effects of inhaled NO_2 , suggesting that 5 ppm is a maximum safe dose.¹⁷ Clinical administration systems should minimise the period of NO and O_2 mixing to prevent extracorporeal NO_2 formation. Within tissue NO has a higher affinity for haemoglobin than oxygen, minimising autoxidation in blood.

FORMATION OF NITROSOTHIOL COMPOUNDS

NO undergoes nitrosation reactions, and in biological systems the formation of nitrosothiols seems to be favoured. The reaction of NO with RSH to form RSNO is rapid but the resulting nitrosothiol is unstable and the reaction is reversible.¹⁰ The presence of nitrosothiols in human plasma supports the

suggestion that nitrosothiols may represent a 'stabilised' form of NO in biological tissue.¹⁸

Pharmacokinetics

Inhaled NO is taken up into the precapillary airspaces and alveoli at a rate many times faster than inhaled oxygen, and the diffusing capacity of the lung is 4.5 times higher for NO than for carbon monoxide.¹⁹ The anatomical proximity of the airspaces to muscular arterioles allows NO to diffuse into contact with the abluminal surface of these vessels.¹⁹

Because of rapid removal of NO by reaction with haemoglobin, the effects of NO inhalation are generally considered to be localised to lung tissue with an effective half life of 2–6 seconds.²⁰ The half life of potential 'stabilised' forms of NO (S-nitrosothiols) is longer, and nitrosothiols may prolong the actions of NO in tissue.^{9, 21} The metabolites of NO are cleared from the body by the kidneys within five to eight hours⁹ and concentrations of methaemoglobin rarely rise above 1–2% during NO treatment.^{22, 23}

Pharmacology

MECHANISMS OF ACTION

Inhaled NO diffuses from alveoli into cells¹⁶ where it binds to the prosthetic haem group of the soluble guanylate cyclase enzyme (GTP pyrophosphatase (cyclising); EC 4.6.1.2).²⁴ NO binding leads to a conformational change similar the effect of oxygen,²⁵ and causes a rise in intracellular cyclic guanylate monophosphate (cGMP), which in turn leads to muscular relaxation and vasodilation (figure). An independent effect of NO on calcium dependent potassium channels has also been described,²⁶ and preliminary evidence suggests a direct activation of G-proteins.²⁷ NO may also stimulate prostanoic synthesis by reacting with the haem in prostaglandin H synthase.²⁸

It remains controversial whether these effects are entirely due to the action of NO itself, or also to nitrosothiol metabolites such

Physiological and pharmacological actions of NO on vascular smooth muscle. Physiological stimuli, such as acetylcholine, activate constitutive nitric oxide synthase (NOS) in endothelial cells, which converts L-arginine and oxygen to L-citrulline and nitric oxide (NO). NO diffuses to smooth muscle cells and activates soluble guanylate cyclase (sGC) which converts GTP to cyclic-GMP, leading to muscular relaxation. Pharmacological administration of NO into the airways acts directly on guanylate cyclase.

as S-nitrosocysteine.^{29 30} Inhaled NO may also have some distant effects mediated by nitrosothiols, as prolongation of the bleeding time has been reported in healthy adult volunteers who inhaled NO,³¹ and rats treated with inhaled NO developed prolonged brainstem evoked responses and impaired learning.³² However, such actions are probably minor.

PHARMACOLOGICAL EFFECTS

NO mediated relaxation of vascular smooth muscle leads to vasodilation⁵ and bronchodilation.^{16 33} In newborn lambs and piglets it has no effect on the normal pulmonary vasculature, but reverses the vasoconstriction induced by hypoxia or vasoconstrictor prostanoids, and this effect is not abolished by concomitant acidemia.^{34–36} Vasodilation can occur in the presence of endothelial damage which abolishes the vasodilatory effect of acetylcholine,³⁷ and it is more effective than inhalation of prostaglandin I₂.³⁸ In piglets inhaled NO reduces the vasoconstriction associated with bacterial sepsis.³⁹

Because inhaled NO is delivered to ventilated lung segments, its vasodilatory effect improves ventilation:perfusion matching and oxygen transport so that improved oxygenation has been observed even without a fall in pulmonary artery pressure.⁴⁰ This effect may be potentiated by the effects of NO on bronchial smooth muscle: inhaled NO reduces the bronchoconstrictor effect of inhaled methacholine in guinea pigs and rabbits, although at inhaled concentrations of 80–300 ppm, higher than those required for vasodilation (5–80 ppm).¹⁶

Other actions of NO may be relevant during inhalation treatment: leakage of albumin into the alveolar space after pulmonary injury is reduced⁴¹; adhesion and aggregation of platelets and leucocytes are inhibited⁴²; the effects of endothelin-1 and other vasoactive agents on pulmonary vessels may be antagonised.^{43 44} NO has a negative feedback effect on NO synthesis, raising concern that inhaled NO may depress endogenous pulmonary vasodilation.^{45–47}

NO has many further biological effects, including inhibition of platelet aggregation⁵ and a negative inotropic effect.⁴⁸ Cytokine activated macrophages use NO to kill tumour cells, fungi, bacteria, viruses and helminths.⁸ NO can disrupt enzymes in both the Krebs cycle (aconitase), mitochondrial complex I (reduced nicotinamide-adenine dinucleotide phosphate-NADH dehydrogenase), and mitochondrial complex II (succinate-NADH dehydrogenase), as well as the rate limiting enzyme in DNA synthesis (ribonucleoside-diphosphate reductase).⁴⁹ NO modulates gene transcription and post-transcription processing⁵⁰ and may play an important part in inflammation: it enhances the effects of cyclooxygenase I and II and stimulates the production of some proinflammatory eicosanoids.²⁸ Immune complex induced injury in rat lung⁵¹ and inflammatory damage to other tissues are reduced by NO inhibitors.^{52 53} NO acts as a

neurotransmitter in the central and peripheral nervous systems, notably in non-adrenergic, non-cholinergic fibres, which are prominent in lung.¹⁶ NO has been implicated as a mediator of cell death in the nervous system.⁵⁴

Toxicology

Inhalational NO is not a licensed drug, and full toxicology data are unavailable. Although the many functions of NO in the body mean that modulation of NO metabolism or effects could have far reaching results, it is generally assumed that systemic toxicity is low because of rapid inactivation of NO on contact with blood.

Studies in rats have shown that inhalation of up to 1500 ppm NO for 15 minutes caused no demonstrable lung injury.⁵⁵ Overdose of 20 000 ppm caused acute pulmonary oedema and methaemoglobinaemia.¹⁶ Unfortunately, such studies are of limited relevance to clinical uses of the drug.

Reports of adverse effects in clinical studies have been rare: paradoxical hypoxia during NO inhalation in one newborn infant has been reported, and it was suggested (without supporting evidence) that this was related to maternal indomethacin treatment.⁵⁶ The increase in bleeding time demonstrated in adult volunteers does not seem to be a significant problem in clinical practice.³¹

Nevertheless, there are theoretical reasons for caution in the use of inhaled NO treatment. The formation of hydroxyl radicals might cause oxidative damage to tissue.¹⁴ NO might impair energy metabolism⁵⁷ or promote inflammation,²⁸ and it has been suggested that NO is involved in mediating the effects of pertussis toxin in airways.⁵⁸ DNA damage in *Salmonella* species due to NO has been recorded, and as cigarette smoke contains NO this has been proposed as one mechanism for the carcinogenic effect of smoking.^{59 60} Some evidence of deteriorations in lung function during inhalation of NO has been produced in studies of adult volunteers: a mean fall in arterial oxygen tension of 7 mm Hg was recorded in 191 normal subjects inhaling NO of 15–20 ppm, and at higher concentrations airways resistance increased.¹⁶

Conclusion

NO is a multifunctional biological mediator with clear pharmacological effects when inhaled into the lung. Considerable gaps remain in our understanding of the pharmacology and toxicology of this novel inhalational treatment, and enthusiasm for NO treatment should be tempered with caution.

I thank Professor M Hughe and Miss S Khan for their invaluable advice.

- 1 Furchgott RF, Zawadzki JV. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature* 1980; 288: 373–6.
- 2 Palmer RM, Ferrige AG, Moncada S. Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* 1987; 327: 524–6.
- 3 Palmer RM, Ashton DS, Moncada S. Vascular endothelial cells synthesize nitric oxide from L-arginine. *Nature* 1988; 333: 664–6.

- 4 Rees DD, Palmer RMJ, Schulz R, Hodson HF, Moncada S. Characterisation of three inhibitors of endothelial nitric oxide synthase in vitro and in vivo. *Br J Pharmacol* 1990; **101**: 746-52.
- 5 Moncada S, Palmer RMJ, Higgs EA. Nitric oxide: Physiology, pathophysiology and pharmacology. *Pharmacol Rev* 1991; **43**: 109-42.
- 6 Takei Y, Edwards AD, Lorek A, Peebles DM, Belai A, Cope M, et al. Effects of Nitro-L-Arginine methyl ester on the cerebral circulation of newborn piglets quantified in vivo by near infrared spectroscopy. *Pediatr Res* 1993; **34**: 354-9.
- 7 Dawson TM, Dawson VL, Snyder SH. A novel neuronal messenger molecule in the brain: the free radical, nitric oxide. *Ann Neurol* 1992; **32**: 297-311.
- 8 Green SJ, Nacy CA, Meltzer MS. Cytokine-induced synthesis of nitrogen oxides in macrophages: a protective host response to Leishmania and other intracellular pathogens. *J Leuk Biol* 1991; **50**: 93-103.
- 9 Anggard E. Nitric oxide: mediator, murderer, and medicine. *Lancet* 1994; **343**: 1199-206.
- 10 Bonner FT, Steadman G. The chemistry of nitric oxide and redox-related species. In: Stamler J, Feelish M, eds. *Methods in nitric oxide research*. Chichester: John Wiley, (in press).
- 11 Stamler JS, Singel DJ, Loscalzo J. Biochemistry of nitric oxide and its redox-activated forms. *Science* 1992; **258**: 1898-902.
- 12 Lipton SA, Choi YB, Pan ZH, Lei SZ, Chen HS, Sucher NJ, et al. A redox-based mechanism for the neuroprotective and neurodestructive effects of nitric oxide and related nitroso-compounds. *Nature* 1993; **364**: 626-32.
- 13 Beckman JS, Beckman TW, Chen J, Marshall PA, Freeman BA. Apparent hydroxyl radical production by peroxynitrite: implications for endothelial injury from nitric oxide and superoxide. *Proc Natl Acad Sci USA* 1990; **87**: 1620-4.
- 14 Radi R, Beckman JS, Buch KM, Freeman BA. Peroxynitrite oxidation of sulfhydryls. The cytotoxic potential of superoxide and nitric oxide. *J Biol Chem* 1991; **266**: 4244-50.
- 15 Foubert L, Fleming B, Latimer R, Jonas M, Odoro A, Borland C, et al. Safety guidelines for use of nitric oxide. *Lancet* 1992; **339**: 1615-6.
- 16 Gaston B, Drazen JM, Loscalzo J, Stamler JS. The biology of nitrogen oxides in the airways. *Am Rev Respir Dis* 1994; **149**: 538-51.
- 17 Centers for Disease Control. Recommendations for occupational safety and health standards. *MMWR* 1988; **37** (suppl): 21.
- 18 Keaney JFJ, Simon DI, Stamler JS, Jaraki O, Scharfstein J, Vita JA, et al. NO forms an adduct with serum albumin that has endothelium-derived relaxing factor-like properties. *J Clin Invest* 1993; **91**: 1582-9.
- 19 Higenbottam T. Inhaled nitric oxide: a magic bullet? *Q J Med* 1993; **86**: 555-8.
- 20 Moncada S. The 1991 Ulf von Euler Lecture. The L-arginine: nitric oxide pathway. *Acta Physiol Scand* 1992; **145**: 201-7.
- 21 Stamler JS, Jaraki O, Osborne J, Simon DI, Keaney J, Vita J, et al. Nitric oxide circulates in mammalian plasma primarily as an S-nitroso adduct of serum albumin. *Proc Natl Acad Sci USA* 1992; **89**: 7674-7.
- 22 Roberts JDJ, Lang P, Bigatello LM, Vlahakes GJ, Zapol WM. Inhaled nitric oxide in congenital heart disease. *Circulation* 1993; **87**: 447-53.
- 23 Kinsella JP, Abman SH. Methaemoglobin during nitric oxide therapy with high-frequency ventilation. *Lancet* 1993; **342**: 615.
- 24 Gruetter CA, Gruetter DY, Lyon JE, Kadowitz PJ, Ignarro LJ. Relationship between cyclic guanosine 3':5'-monophosphate formation and relaxation of coronary arterial smooth muscle by glycyl trinitrate, nitroprusside, nitrite and nitric oxide: effects of methylene blue and methemoglobin. *J Pharmacol Exp Ther* 1981; **219**: 181-6.
- 25 Wolin MS, Wook KS, Ignarro LJ. Guanylate cyclase from bovine lung. A kinetic analysis of the regulation of unpurified soluble enzyme by protoporphyrin IX, haem and nitrosyl-haem. *J Biol Chem* 1982; **257**: 11312-20.
- 26 Bolotina VM, Najibi S, Palacino JJ, Pagano PJ, Cohen RA. Nitric oxide directly activates calcium-dependent potassium channels in vascular smooth muscle. *Nature* 1994; **368**: 850-3.
- 27 Lander HM, Sehajpal PK, Novogrodsky A. Nitric oxide signalling: a possible role for G proteins. *J Immunol* 1993; **151**: 7182-7.
- 28 Salvemini D, Misko TP, Masferrer JL, Seibert K, Currie MG, Needleman P. Nitric oxide activates cyclooxygenase enzymes. *Proc Natl Acad Sci USA* 1993; **90**: 7240-4.
- 29 Feelish M, te Poel M, Zamora R, Deussen A, Moncada S. Understanding the controversy over the identity of EDRF. *Nature* 1994; **368**: 62-5.
- 30 Mathews WR, Kerr SW. Biological activity of S-nitrosothiols: the role of nitric oxide. *J Pharmacol Exp Ther* 1993; **267**: 1529-37.
- 31 Hogman M, Frostell C, Amberg H, Hedenstierna G. Bleeding time prolongation and NO inhalation. *Lancet* 1993; **341**: 1664-5.
- 32 Groll-Knapp E, Haider M, Kienzl K, Handler A, Trimmel M. Changes in discrimination learning and brain activity due to combined exposure to NO and CO in rats. *Toxicol* 1988; **49**: 441-7.
- 33 Barnes PJ. Nitric oxide and airways. *Eur Respir J* 1993; **6**: 163-5.
- 34 Zapol WM, Falke KJ, Hurford WE, Roberts JDJ. Inhaling nitric oxide: a selective pulmonary vasodilator and bronchodilator. *Chest* 1994; **105**: 87-91S.
- 35 Nelin LD, Moshin J, Thomas CJ, Sasidharan P, Dawson CA. The effect of inhaled nitric oxide on the pulmonary circulation of the neonatal pig. *Pediatr Res* 1994; **35**: 20-4.
- 36 Etches PC, Finer NN, Barrington KJ, Graham AJ, Chan WK. Nitric oxide reverses acute hypoxic pulmonary hypertension in the newborn piglet. *Pediatr Res* 1994; **35**: 15-9.
- 37 Wessel DL, Adatia I, Giglia TM, Thompson JE, Kulik TJ. Use of inhaled nitric oxide and acetylcholine in the evaluation of pulmonary hypertension and endothelial function after cardiopulmonary bypass. *Circulation* 1993; **88**: 2128-38.
- 38 Welte M, Zwissler B, Habazettl H, Messmer K. PGI₂ aerosol versus nitric oxide for selective pulmonary vasodilation in hypoxic pulmonary vasoconstriction. *Eur Surg Res* 1993; **25**: 329-40.
- 39 Berger JJ, Gibson RL, Redding GJ, Standaert TA, Clarke WR, Truog WE. Effect of inhaled nitric oxide during group B streptococcal sepsis in piglets. *Am Rev Respir Dis* 1993; **147**: 1080-6.
- 40 Gerlach H, Pappert D, Lewandowski K, Rossaint R, Falke KJ. Long-term inhalation with evaluated low doses of nitric oxide for selective improvement of oxygenation in patients with adult respiratory distress syndrome. *Intensive Care Med* 1993; **19**: 443-9.
- 41 Kurose I, Wolf R, Grisham MB, Granger DN. Modulation of ischemia/reperfusion-induced microvascular dysfunction by nitric oxide. *Circ Res* 1994; **74**: 376-82.
- 42 Dembinska Kiec A, Zmuda A, Wenhryniewicz O, Stachura J, Peskar BA, Gryglewski RJ. Selectin-P-mediated adherence of platelets to neutrophils is regulated by prostanooids and nitric oxide. *Int J Tissue React* 1993; **15**: 55-64.
- 43 Tod ML, Cassin S. Endothelin-1-induced pulmonary arterial dilation is reduced by N^{omega}-nitro-L-arginine in fetal lambs. *J Appl Physiol* 1992; **72**: 1730-4.
- 44 Kourembanas S, McQuillan LP, Leung GK, Faller DV. Nitric oxide regulates the expression of vasoconstrictors and growth factor by vascular endothelium under both normoxia and hypoxia. *J Clin Invest* 1993; **92**: 99-104.
- 45 Buga GM, Griscavage JM, Rogers NE, Ignarro LJ. Negative feedback regulation of endothelial cell function by nitric oxide. *Circ Res* 1993; **73**: 808-12.
- 46 Rengasamy A, Johns RA. Regulation of nitric oxide synthase by nitric oxide. *Mol Pharmacol* 1993; **44**: 124-8.
- 47 Assreuy J, Cunha FQ, Liew FY, Moncada S. Feedback inhibition of nitric oxide synthase activity by nitric oxide. *Br J Pharmacol* 1993; **108**: 833-7.
- 48 Finkel MS, Oddis CV, Jacob TD, Watkins SC, Hattler BG, Simmons RL. Negative inotropic effects of cytokines on the heart mediated by nitric oxide. *Science* 1992; **257**: 387-9.
- 49 Lowenstein CJ, Dinerman JL, Snyder SH. Nitric oxide: a physiologic messenger. *Ann Intern Med* 1994; **120**: 227-37.
- 50 Pantopoulos K, Weiss G, Hentze MW. Nitric oxide and post-transcriptional control of cellular traffic. *Trends Cell Biol* 1994; **4**: 82.
- 51 Mulligan MS, Hevel JM, Marletta MA, Ward PA. Tissue injury caused by deposition of immune complexes is L-arginine dependant. *Proc Natl Acad Sci USA* 1991; **88**: 6338-42.
- 52 Ialenti A, Moncada S, Di Rosa M. Modulation of adjuvant arthritis by endogenous nitric oxide. *Br J Pharmacol* 1993; **110**: 701-6.
- 53 Moncada S, Higgs A. The L-arginine-nitric oxide pathway. *N Engl J Med* 1993; **329**: 2002-12.
- 54 Dalkara T, Moskowitz MA. The complex role of nitric oxide in the pathophysiology of focal cerebral ischaemia. *Brain Pathol* 1994; **4**: 49-57.
- 55 Stavert DM, Lehnert BE. Nitric oxide and nitrogen dioxide as inducers of acute pulmonary injury when inhaled at relatively high concentrations for brief periods. *Inhalational Toxicol* 1990; **2**: 53-67.
- 56 Oriot D, Boussemart T, Berthier M, Bonneau D, Coisne D. Paradoxical effect of inhaled nitric oxide in a newborn with pulmonary hypertension. *Lancet* 1993; **342**: 364-5.
- 57 Cleeter MW, Cooper JM, Darley-Usmar VM, Moncada S, Schapira AVH. Reversible inhibition of cytochrome c oxidase, the terminal enzyme of the mitochondrial respiratory chain, by nitric oxide. *FEBS Lett* 1994; **345**: 50-4.
- 58 Heiss LN, Lancaster JRJ, Corbett JA, Goldman WE. Epithelial autotoxicity of nitric oxide: role in the respiratory cytopathology of pertussis. *Proc Natl Acad Sci USA* 1994; **91**: 267-70.
- 59 Nguyen T, Brunson D, Crespi CL, Penman BW, Wishnok JS, Tannenbaum SR. DNA damage and mutation in human cells exposed to nitric oxide in vitro. *Proc Natl Acad Sci USA* 1992; **89**: 3030-4.
- 60 Routledge MN, Wink DA, Keefer LK, Dippe A. Mutations induced by saturated aqueous nitric oxide in the pSP189 supF gene in human Ad293 and E coli MBM7070 cells. *Carcinogenesis* 1993; **14**: 1251-4.