Nasal high frequency ventilation in neonates with moderate respiratory insufficiency

Mark van der Hoeven, Erik Brouwer, Carlos E Blanco

Abstract

Aim—To investigate the efficacy of nasal high frequency ventilation (nHFV) in newborn infants with moderate respiratory insufficiency.

Method—Twenty one preterm and term neonates were treated with nHFV for respiratory insufficiency. Criteria for starting nHFV were: deterioration on nasal CPAP expressed by a median pH of 7.24 and pCO₂ of 8.3 kPa, or increasing FIO₂. nHFV was delivered using the Infant Star ventilator. Ventilator setting amplitude was 35 cm H₂O; mean airway pressure 7 cm H₂O; and frequency 10 Hz.

Results—pCO₂ decreased significantly from 8.3 kPa to 7.2 kPa after nHFV was started. In five patients nHFV was discontinued after a median period of 6½ hours due to CO₂ retention and high oxygen need, and endotracheal mechanical ventilation was started.

Conclusions—nHFV can reduce pCO₂ in neonates with moderate respiratory insufficiency and, therefore, could be used to decrease the need for endotracheal mechanical ventilation.

(Arch Dis Child Fetal Neonatal Ed 1998;79:F61–F63)

Keywords: nasal high frequency ventilation; idiopathic respiratory distress syndrome

Department of Neonatology, Academic Hospital Maastricht, P Debyelaan 25 PO Box 5800 6202 AZ Maastricht, The Netherlands.
Mark van der Hoeven
Erik Brouwer
Carlos E Blanco

Correspondence to:
Dr Mark van der Hoeven.
Email:
MVH@SKIN.AZM.NL

Accepted 14 January 1998

Over the past two decades assisted mechanical ventilation has been used to manage respiratory insufficiency in newborn infants. However, complications such as airleaks occur in 15 to 48% of the infants who need ventilator support,^{1,2} and 5 to 25% develop bronchopulmonary dysplasia (BPD).³ Both these complications contribute substantially to morbidity in preterm infants. Therefore, a less invasive approach such as nasal continuous positive airway pressure (nCPAP) is frequently used in the initial management of respiratory insufficiency.^{4–6} But 43 to 80% of the neonates with moderate to severe respiratory insufficiency, initially treated with nasal CPAP, will

Table 1 Summary of clinical data of 21 infants studied

	Median (20th–80th percentile)
Birthweight (g)	1010 (750–2170)
Gestation (weeks)	29 (27–32)
Age at study (h)	24 (4–480)
Duration nHFV (h)	36 (12–108)
IRDS	10
TTN	3
Air leak	2
Apnoea	6
Required mechanical ventilation	5

IRDS: idiopathic respiratory distress syndrome; TTN: transient tachypnoea of the newborn

need subsequent mechanical ventilation.⁷ Nasal CPAP has also been used after extubation to facilitate weaning from mechanical ventilation.^{8–11} However, extubation failed in 16 to 40% of cases. In both events subsequent endotracheal mechanical ventilation was needed because of CO₂ retention or increased oxygen need and apnoea of prematurity.

High frequency ventilation (HFV) is a new method of mechanical ventilation based on the delivery of small tidal volumes at a supraphysiological respiratory frequency of 10 Hz.^{12,13} This technique is very effective in eliminating carbon dioxide and is independent of dead space.

We hypothesised that HFV applied on a single nasopharyngeal tube (nHFV) might reduce pCO₂ by augmenting CO₂ diffusion in the pharynx and larynx area and, therefore, decreasing the need for subsequent mechanical ventilation. To test this hypothesis we performed a non-controlled observational study and tested the efficacy of nHFV in infants with moderate respiratory insufficiency.

Methods

Twenty one newborn infants were offered nHFV. All patients were initially treated with nasal CPAP applied via a single nasopharyngeal tube positioned at a depth of 3–4 cm. nHFV was started due to CO₂ retention (median value 8.3 kPa), acidosis (median pH value 7.24), and/or increasing oxygen need

Table 2 Values (median with 20th and 80th percentiles) of blood gases and ventilator settings before and after initiation of nHFV

	Before		After			
	n=19	n=21	n=21	n=14	n=15	n=15
Time (h)	–4.15' (–6.25'– –3.28')	–0.57' (–1.57'– –0.34')	1.30' (0.50'–2.00')	4.24' (3.20'–5.20')	7.38' (6.52'–8.00')	11' (10.00'–13.00')
pH	7.24 (7.20–7.28)	7.24 (7.19–7.29)	7.26 (7.24–7.32)	7.28 (7.24–7.32)	7.31 (7.24–7.42)	7.31 (7.27–7.36)
pCO ₂ (kPa)	8.0 (7.3–8.9)	8.3 (7.4–9.1)	7.2 (6.6–8.3)*	7.1 (6.5–7.9)*	6.5 (5.7–7.9)¶	6.7 (5.9–7.4)¶
Paw (cm H ₂ O)	5 (5–6)	6 (5–7)	7 (6–8)§	7 (6–8)§	7 (6–8)§	7 (6–8)§
Amplitude (cm H ₂ O)			37 (29–46)	35 (28–47)	35 (29–47)	35 (28–48)
FIO ₂	0.35 (0.21–0.5)	0.42 (0.21–0.60)	0.40 (0.21–0.60)	0.40 (0.21–0.50)	0.37 (0.21–0.51)	0.32 (0.21–0.51)

*pCO₂ was significantly different (Wilcoxon p<0.005) from pCO₂ in the two blood gases registered before nHFV was initiated. ¶pCO₂ was significantly different (Wilcoxon p<0.01) from pCO₂ in the two blood gases registered before nHFV. §Paw settings after the start of nHFV were significantly different (Wilcoxon p<0.01) from each of the two Paw pressure values registered before nHFV was initiated.

Figure 1 Course of $p\text{CO}_2$ in blood gas before and two blood gases after start of nHFV. $p\text{CO}_2$ decreased significantly in both samples taken at a median time of 1½ hours (0.50–2 hours) and at 4.24 (3.2–5.2) hours after the start of nHFV compared with that in the sample taken at 0.57 (1.57–0.34) hours before nHFV was started. $p = 0.0012$ and 0.0033 , respectively.

(median FIO_2 0.42) during nasal CPAP. We applied nHFV via the same nasopharyngeal tube by switching the ventilator (Infant Star HFV, Infrasonics, Inc. San Diego, CA, USA) to the HFV mode. This ventilator delivers HFV by flow interruption. NHFV was initiated with the following parameters: mean airway pressure (Paw) was maintained or increased to a higher level; frequency of 10 Hz and amplitude was increased until the infant's chest showed oscillations; FIO_2 was regulated to achieve a peripheral oxygen saturation monitored by pulse oximetry of 86–93%.^{14–15} Arterial or capillary blood gases and ventilator settings were registered at a median time of 4¼ hours (3.28–6.25 hours) and of 0.57 hours (0.34–1.57

Figure 2 Course of $p\text{CO}_2$ in two blood gases measured before (4.15 and 0.57 hours) and four blood gases measured after (1.3, 4.24, 7.38 and 11 hours) the start of nHFV. Median and 20th and 80th percentiles are shown together.

hours) before and 1½ hours (0.50–2 hours), 4.24 hours (3.20–5.20 hours), 7.38 hours (6.52–8 hours), and 11 hours (10–13 hours) after nHFV initiation. Endotracheal mechanical ventilation was considered when blood gases further deteriorated: $\text{pH} < 7.20$; $p\text{CO}_2 > 8.5$ kPa; and $\text{FIO}_2 > 0.80$ or apnoea.

All values are reported as median with 20th and 80th percentiles. A Wilcoxon signed rank sum test for paired data was used to estimate statistical difference for blood gases and ventilator settings. Differences were considered statistical significant when $p < 0.05$.

Results

Twenty one patients were treated with nHFV. Table 1 summarises the clinical data. In five (23%) patients, nHFV was discontinued after a median time of 6½ hours (2–8½ hours), and endotracheal mechanical ventilation was started. Table 2 shows blood gas values and ventilator settings before and after initiating nHFV. NHFV was started in 16 infants because of CO_2 retention ($p\text{CO}_2$ value > 7.5 kPa) and in five infants because of increasing oxygen need ($\text{FIO}_2 > 0.55$). A small but significant reduction in $p\text{CO}_2$ was found ($p = 0.0012$; $p = 0.0033$) after initiation of nHFV compared with the two last blood gases taken before nHFV (figs 1 and 2). Paw pressure was significantly higher after the initiation of nHFV.

In 15/21 (72%) patients, nHFV was initiated early in the course of their respiratory disease (median age of 17 hours; 2–36). Respiratory insufficiency was due to idiopathic respiratory distress syndrome ($n = 10$), transient tachypnoea of the newborn (TTN) ($n = 3$), and hypoventilation ($n = 2$). NHFV was continued for a median (range) period of 35 (2–144) hours. In three patients with severe idiopathic RDS, nHFV was discontinued and endotracheal mechanical ventilation with surfactant replacement was started within 2 to 6 hours because of further deterioration in blood gases or increased oxygen need.

In 6/21 (28%) patients, nHFV was initiated at a median age 9 days (6–21), after a period of endotracheal mechanical ventilation. The diagnoses included air leak syndrome ($n = 2$) and apnoea associated with sepsis and necrotising enterocolitis ($n = 4$). NHFV in this group was continued for a median period of 40 hours (17–126). In two patients in this group with apnoea and sepsis, nHFV was discontinued and endotracheal mechanical ventilation was started because of further deterioration in blood gases and apnoea.

Discussion

This study has shown that nHFV can reduce $p\text{CO}_2$ in patients with nasal CPAP with different respiratory problems. Whether nHFV can be recommended as a valuable mode of respiratory support must be determined in a prospective randomised study. We showed that nHFV was effective in reducing $p\text{CO}_2$ from 8.3 kPa to 7.2 kPa within 2 hours.

The mechanism of CO_2 reduction during HFV is complex and is not completely understood.^{13–16} As oscillations of the chest are

visible during nHFV, direct ventilation of the lung is less important; the larynxes are a significant factor in resistance.¹⁷ Gas transport during nHFV is probably largely improved by the interaction of convective and diffusive gas flux (also called augmented diffusion) in the pharyngeal and laryngeal area, as has been reported during HFV in the larger and medium airways.^{16, 18}

Nasal CPAP has been proposed as a less invasive approach for the management of respiratory distress in very low birthweight infants.⁴⁻⁶ After the introduction of nasal CPAP, a significant decrease in the incidence of intubation was reported.¹⁹ Another report showed that the incidence of BPD was decreased in association with the use of early nasal CPAP and permissive hypercapnia.²⁰ However, the use of nasal CPAP seemed to be limited to a small group of infants, as 43 to 80% of the infants with moderate to severe idiopathic RDS went on to require mechanical ventilation due to CO₂ retention, increased oxygen need, and apnoea of prematurity.⁷ The level of CPAP pressure is associated with an increase in paO₂, indicating an increased functional residual capacity and/or alveolar recruitment.²¹⁻²² But high CPAP pressure was also associated with an increase in paCO₂ due to an increase in dead space, alveolar overdistension, and a decrease in alveolar ventilation.²¹⁻²³ nHFV could be used to control CO₂ retention and to permit the use of a higher P_{aw} pressure so as to reduce oxygen need.

Gaseous distension of the bowel has been mentioned as a disadvantage of the use of nasal CPAP in premature infants. This so-called CPAP belly syndrome has not been associated with necrotising enterocolitis or bowel obstruction and had a benign development.²⁴

In our study some infants who were breathing room air still had clinically significant CO₂ retention. In the alveolar air equation alveolar pO₂ is determined by the inspiratory oxygen concentration minus the alveolar pCO₂, divided by the respiratory exchange ratio. When an alveolar pCO₂ is increased to, for example, 10 kPa and the infant is breathing room air and has a respiratory exchange ratio of 0.8, the alveolar pO₂ will be 8.5 kPa. Assuming that the gradient between alveolar and arterial pO₂ is not severely increased by atelectasis, scatter of ventilation perfusion ratios, or by the presence of a right to left shunt, a clinically significant hypercapnia can exist in association with a low but still normal and acceptable arterial pO₂. In addition, as the P90 (the paO₂ required for 90% oxygen saturation of haemoglobin) in preterm infants was reported to be 5.4 (0.5) kPa,²⁵ there is no clinical or physiological basis to increase the PO₂ value when almost all haemoglobin is bound with oxygen.

In a heterogeneous group of infants with moderate respiratory diseases, nHFV was effective in reducing pCO₂. nHFV might,

therefore, be a valuable addition to the ventilatory management of the newborn infant, and might reduce the need for subsequent mechanical ventilation.

- Hart SM, McNair M, Gasmu MR, *et al*. Pulmonary interstitial emphysema in very low birth weight infants. *Arch Dis Child* 1983;58:612-16.
- Primhak RA. Factors associated with pulmonary air leak in premature infants receiving mechanical ventilation. *J Pediatr* 1983;102:764-8.
- Bancalari E. Neonatal chronic lung disease. In: Fanaroff AA, Martin RJ. *Neonatal-perinatal medicine. Diseases of the fetus and infant*. St Louis: Mosby, 1997:1074-89.
- Jacobsen T, Gronvall J, Petersen S, Andersen GE. Minitouch treatment of very low birth weight infants. *Acta Paediatrica* 1993;83:934-8.
- Kamper J, Wulff K, Larsen C, Lindequist S. Early treatment with nasal continuous positive airway in very low birth weight infants. *Acta Paediatrica* 1993;82:193-7.
- Lundstrom KE. Initial treatment of premature infants: continuous positive airway pressure or ventilation? *Eur J Pediatr* 1996;155:S25-S9.
- Verder H, Robertson B, Greisen G, *et al*. Surfactant therapy and nasal continuous positive airway pressure for newborns with respiratory distress syndrome. Danish-Swedish multicentre study group. *N Engl J Med* 1994;331:1051-5.
- So BH, Tamura M, Mishina J, Watanabe T, Kamoshita S. Application of nasal continuous positive airway pressure to early extubation in very low birthweight infants. *Arch Dis Child* 1995;72:F191-F3.
- Higgins RD, Richter SE, Davis JM. Nasal continuous positive airway pressure facilitates extubation of very low birth weight neonates. *Pediatrics* 1991;88:999-1003.
- Tapia JL, Bancalari A, Gonzalez A, Mercado ME. Does continuous positive airway pressure (CPAP) during weaning from intermittent mandatory ventilation in very low birthweight infants have risks or benefits? A controlled trial. *Pediatr Pulmonol* 1995;5:269-74.
- Chan V, Greenough A. Randomized trial of methods of extubation in acute and chronic respiratory distress. *Arch Dis Child* 1993;68:570-2.
- Clark RH, Null DM. High-Frequency oscillatory ventilation. In: Pomerance JJ, Richardson CJ, eds. *Neonatology for the clinician*. Norwalk, CT: Appleton & Lange, 1993:289-309.
- Clark RH. High-frequency ventilation. *J Pediatr* 1994;124:661-70.
- Wasunna A, Whitelaw AGL. Pulseoximetry in preterm infants. *Arch Dis Child* 1987;62:882-8.
- Joint working group of the British Association of Perinatal Medicine and the Research Unit of the Royal College of Physicians. Development of audit measures and guidelines for good practice in the management of neonatal respiratory distress syndrome. *Arch Dis Child* 1992;67:1221-7.
- Boynton BR, Carlo WA. Pulmonary gas exchange during high-frequency ventilation. In: Boynton BR, Carlo WA, Jobe AH, eds. *New therapies for neonatal respiratory failure. A physiological approach*. Cambridge: Cambridge University Press; 1994:218-44.
- Fredberg JJ. Summary: Pulmonary mechanics during high frequency ventilation. *Acta Anaesthesiol Scand* 1989;33:170-1.
- Chang HK. Mechanisms of gas transport during ventilation by high-frequency oscillation. *J Appl Physiol* 1984;56:553-64.
- Gittermann MK, Fusch C, Gittermann AR, Regazzoni BM, Moessinger AC. Early nasal continuous positive airway pressure treatment reduces the need for intubation in very low birth weight infants. *Eur J Pediatr* 1997;156:384-8.
- Avery MA, Tooley WH, Keller JB, *et al*. Is chronic lung disease in low birth weight infants preventable? A survey of eight centers. *Pediatrics* 1987;79:26-30.
- Osher AB, Gothe B, Simmons DH, Waymost B, Green H. Effects of continuous positive airway pressure after oleic acid-induced lung injury in dogs. *Pediatr Res* 1978;9:923-6.
- Richardson CP, Jung AL. Effects of continuous positive airway pressure on pulmonary function and blood gases of infants with respiratory distress syndrome. *Pediatr Res* 1978;12:771-4.
- Bonta BW, Uauy R, Warshaw JB, Motoyama EK. Determination of optimal continuous positive pressure for the treatment of IRDS by measurement of esophageal pressure. *J Pediatr* 1977;91:449-54.
- Jaille JC, Levin T, Wung JT, Abramson SJ, Ruzal-Shapiro C, Berdon WE. Benign gaseous distension of the bowel in premature infants treated with nasal continuous airway pressure. *Am J Rheumatol* 1992;158:125-7.
- Émond D, Lachance C, Gagnon J, Bard H. Arterial partial pressure of oxygen required to achieve 90 % saturation of hemoglobin in very low birth weight newborns. *Pediatrics* 1993;91:602-4.