CURRENT TOPIC

Feeding issues in preterm infants

A major concern for those taking care of preterm infants is to ensure that nutritional intake meets requirements, thereby ensuring that poor nutrition is not rate limiting on outcome. However, establishment of an adequate intake is difficult during early life in the sick infant. Dietary needs also vary, depending on maturity and nutritional and clinical status. Furthermore, measures of outcome are not widely agreed on. This paper will briefly review some of the principles involved and address some of the practical questions that arise on a day to day basis during nutritional care of these high risk infants.

Determination of nutritional requirements

Recommended dietary intakes are based on needs for maintenance and growth and the assumption that post-natal growth approximates that in utero at the same post conceptional age.¹ However, recommended intakes take time to establish, and, having been established, are commonly interrupted for clinical reasons during the first three to four weeks of life in preterm infants.

Figure 1 illustrates this situation more clearly. Embleton *et al*² compared actual and recommended dietary intake (energy 102 kcal/kg/day; protein 3.0 g/kg/day) in a group of preterm infants (≤ 34 weeks gestation) during the initial hospital stay. By 7 days of age, infants had developed major deficits in energy (~ 400 kcal/kg) and protein (13 g/kg),

which were not recovered by the time of discharge from hospital; the more immature the infant the greater the deficit at discharge.²

It has therefore been suggested that dietary intake must also meet needs for “catch up” growth.³ Whether or not this can be accomplished before hospital discharge remains to be determined, but closer attention must be paid to quantifying and minimising these deficits.

Measures of outcome

The relation between diet and growth appears to be clear cut in that inadequate dietary intake can be directly related to poor growth. However, non-dietary factors also affect growth. In the study of Embleton *et al*,² 50% of the variation in growth was related to diet, and 50% was unexplained. This is not surprising given the differences in size, maturity, and nutritional status at birth and the variable clinical course in these infants. Nonetheless, attention must be paid to the confounding effects of non-nutritional factors when interpreting the effects of diet on growth in these infants.

Weight gain and weight for corrected age are valid measures of adequacy of intake. It is recommended that weight gain approximates that of the normal fetus at the same post conceptional age (10–15 g/kg/day).¹ This may be questioned. After birth, preterm infants lose weight and take variable periods to regain birth weight. Achieved weight therefore will always lag behind expected weight at the same post conceptional age; the more immature the infant the greater the deficit.^{4,5} If infants are to catch up, then weight gain must exceed that found in utero, a rate seldom achieved during the initial hospital stay.

Crown-heel length and length gain are the most sensitive indices of adequacy of nutrient intake,⁶ but accurate and reproducible measures of crown-heel length are difficult to obtain. Knee-heel length, as determined by knemometry, has therefore been used to assess linear growth in preterm infants.^{7,8} Still, recent data suggest that the relation between knee-heel and crown-heel length is not consistent, and the former is less reproducible than the latter when linear growth is being assessed.⁹ Further studies are needed to examine this critical issue more closely.

The relation between diet, brain growth, and behaviour is complex.^{10,11} Studies in term infants show that diet alone may not be adequate in improving behaviour¹² and that attention must be paid to social and environmental factors that deprive infants of normal, parental, emotional, and educative care. Studies in preterm infants have shown that even short periods of dietary manipulation affect behaviour.¹³ However, the same studies show that non-nutritional factors also affect behaviour,¹⁴ and it is not easy to disentangle the effects of one from the other.^{15–17} Nonetheless, development is considered by some to be an important end point when long term outcome is assessed in these high risk infants.

Figure 1 Actual and recommended dietary intake (energy (A) and protein (B)) in a group of preterm infants during the initial hospital stay. * $p < 0.001$. Data taken from Embleton *et al*.²

Figure 2 Normal intestinal function.

To feed or not to feed is not the question; when, what, and how are

Fasting is associated with reduced intestinal motility, intestinal mucosal atrophy, and a longer time to establish enteral nutrition. This is consistent with the idea that enteral nutrition is critical for normal intestinal function (fig 2).¹⁸ However, the introduction of feeds in preterm infants is tempered by concerns about feeding intolerance, gastroesophageal reflux, and/or necrotising enterocolitis.

WHEN TO FEED

Withholding enteral feeds does not appear to prevent necrotising enterocolitis,¹⁹ but small amounts of enteral feeds have been shown to stimulate surges in secretion of intestinal polypeptide hormone thought to be important in postnatal intestinal adaptation.²⁰ Dunn *et al*²¹ prospectively examined the effects of early (day 3 of life) hypocaloric enteral nutrition (10–20 ml/kg/day) and reported less jaundice, osteopenia of prematurity, and earlier establishment of full enteral feeds with the hypocaloric regimen.

Slagle *et al*²² also examined the effects of early low volume enteral feed substrate (12 ml/kg/day) on intestinal function and noted improved feeding tolerance and earlier establishment of full enteral nutrition with the hypocaloric regimen. McLure and Newell²³ have also reported increased intestinal motility in infants receiving minimal enteral nutrition. If withholding feeds does not prevent necrotising enterocolitis, there appears little reason not to begin early hypocaloric enteral feeds in these nutritionally vulnerable infants.

WHAT TO FEED

The initial goal of enteral feeds is to support postnatal intestinal adaptation. Data suggest that tolerance is better²⁴ and the incidence of necrotising enterocolitis is lower²⁵ with human milk than low birthweight infant formula. Hypocaloric feeds with human milk therefore appear desirable during the first two weeks of life. If human milk is not available, term infant formula may be fed until enteral tolerance has been achieved, at which point a preterm infant formula should be introduced.

As tolerance is established, intake must meet growth needs. Preterm infants fed on human milk gain less weight than those fed on fortified human milk.^{26–29} Preterm infants fed on human milk alone also show poorer growth than those fed on preterm infant formula.^{30–34} These findings are not surprising because of the lower concentrations of protein and minerals in human milk.³⁵ The data therefore suggest that preterm infants be fed on fortified human milk, a preterm infant formula, or a combination thereof.

However, fortification of human milk such that intake meets requirements is not clear cut because the protein and mineral content of human milk is not only low but also highly variable.³⁵ At the same time, the nutrient content of human milk fortifiers vary. In effect, the degree of fortifica-

tion needed to sustain adequate growth may vary from day to day. It is worrying that preterm infants fed on current fortification regimens grow less well than those fed on a preterm infant formula.^{36, 37} Further studies must examine this critical issue closely.

Controversy exists about the role of long chain polyunsaturated fatty acids in feeding the preterm infant. The long chain polyunsaturated fatty acids arachidonic acid and docosahexanoic acid are critical for eicosanoid synthesis and cell membrane, retinal, and brain function. However, the synthesis of arachidonic acid and docosahexanoic acid may be limited in the preterm infant, and formula supplementation has been suggested.³⁸ Some studies have suggested a benefit,³⁹ others a transient effect,⁴⁰ others no effect,⁴¹ and some an adverse effect on growth.^{42–43} A recent summary report suggests that further data are needed before final recommendations can be made.⁴⁴

HOW TO FEED

Nasogastric versus orogastric enteral feeds

The narrowest point of the airway is the anterior nasal valve, located just behind the nose. Nasal resistance accounts for ~40% of total airway resistance in neonates.⁴⁵ Passage of a size 5 French nasogastric tube increases airway resistance in preterm infants by 30–50%,⁴⁶ and, presumably, the work of breathing.

Greenspan *et al*⁴⁷ examined pulmonary function in normal neonates with nasogastric and orogastric tubes. Infants with nasogastric tubes had diminished minute ventilation and respiratory rate and increased pulmonary resistance, resistive work of breathing, and peak transpulmonary pressure. The authors concluded that “neonates demonstrate significant pulmonary compromise with nasogastric placement”.

Van Someren *et al*⁴⁸ also examined the effects of nasogastric and orogastric tube placement on respiratory function in preterm infants. An increased incidence of periodic breathing and central apnoea were noted with nasogastric tube placement. These data therefore do not support the routine use of nasogastric tubes in preterm infants who are at risk for apnoea and/or those with respiratory problems such as chronic lung disease of prematurity.

Continuous versus bolus enteral feeds

Enteral feeding stimulates an enterohormonal surge. Data from Aynsley-Green *et al*⁴⁹ suggested that surges in insulin, gastrin, and gastric inhibitory peptide are less with continuous than with bolus feeds. However, preprandial hormone levels were higher with continuous feeds,⁴⁹ but the significance of these findings are not entirely clear.

Toce *et al*⁵⁰ compared growth with feeding related complications in preterm infants randomised to continuous or bolus enteral feeds. Weight gain was greater (+ 3–6 g/kg/day) and the incidence of apnoea tended to be less with continuous enteral feeds. Silvestre *et al*⁵¹ was unable to show any differences in growth or incidence of apnoea between continuously fed and bolus fed infants. However, infants withdrawn from the study because of feeding related problems (recurrent apnoea and bradycardias, abdominal distension, and increased gastric residues) were all bolus fed.⁵¹

Studies in term^{52–53} and preterm^{54–57} infants consistently document falls in oxygenation after feeding. The reasons for this are not entirely clear. Patel *et al*⁵⁸ were unable to show any changes in respiratory function. Others have shown decreased functional residual capacity⁵⁹ and/or decreased lung volumes.⁶⁰ Blondheim *et al*⁶² have also reported increased pulmonary resistance and reduced tidal

volume, minute ventilation, and dynamic compliance in bolus fed compared with continuously fed preterm infants.

Nelle *et al*⁶¹ continuously measured middle cerebral blood flow velocity, oxygenation, and blood pressure before, during, and after bolus tube feeding. Mean blood flow velocity decreased from 5 to 11 minutes after feeding and returned to normal after 17 minutes. There were no changes in oxygenation or blood pressure. The authors concluded that "bolus tube feeding in preterm infants may decrease cerebral perfusion, despite unchanged blood pressure and unchanged oxygen saturation".⁶¹

Given the data presented above and the fact that infants are continuously fed in utero, there appears to be little reason for the routine use of bolus feeds in the small preterm infant. Introduction of bolus feeds seems logical when suck/swallowing coordination has developed and infants are ready to progress from gavage to oral feeds.

Nutrition after discharge from hospital

Postnatal growth retardation appears inevitable during the initial hospital stay, with preterm infants weighing significantly less than expected at hospital discharge.⁴⁻⁶² After discharge, little catch up growth occurs, and infants remain small throughout infancy and childhood.⁶³⁻⁶⁷ Despite these observations, limited attention has been paid to nutrition in these infants after discharge.

In one study, preterm infants (< 1850 g; n = 32) were fed on either a standard term infant or nutrient enriched formula after hospital discharge; weight gain and linear growth were greater with the nutrient enriched formula.⁶⁸ In another study, infants (n < 1750 g; n = 129) were fed on either a standard term or preterm infant formula after discharge from hospital; weight, length, and head growth were greater with the preterm infant formula.⁶² Increased weight gain primarily reflected increased lean body mass, suggesting that the preterm formula more closely met protein requirements.² However, questions remain about these studies.

In the study of Lucas *et al*,⁶⁸ no differences were detected in nutrient intake between the groups, making differences in growth difficult to explain. In that of Cooke *et al*,⁶²⁻⁶⁹ better growth was related to increased protein intake, but, at identical nutrient intakes, boys fed the preterm infant formula grew faster and accreted more lean mass than girls fed the preterm infant formula. This is surprising but suggests that factors other than nutrition may be important determinants of growth in preterm boys and girls.

Nonetheless, both studies suggest that "normal" preterm infants benefit from nutritional intervention after hospital discharge. Clear cut recommendations are difficult because nutritional status, and therefore needs, vary. In both trials, formula with a protein/energy ratio of 2.8 g/100 kcal was fed until term, 6,⁶² or 9 months corrected age.⁶⁸ Feeding until term had no long term effects on growth.⁶² It therefore appears to be logical to feed formula with a similar protein/energy ratio until between 3 and 9 months corrected age, but nutritional status must be closely monitored during and after nutritional intervention.

Conclusion

Feeding practices in preterm infants vary widely between and within special care baby units. In the past, this was understandable because feeding was more art than science and practices largely reflected anecdotal experience. This is no longer the case, and the results of randomised controlled trials now provide a "template" on which nutrition support can be based. The results of any randomised trial may be questioned and the effects of intervention may differ depending on patient population. Nonetheless, which-

ever approach is taken within a given unit it must be consistent enough that outcome can be audited and practice adapted to meet the particular needs of infants in that unit.

R J COOKE

N D EMBLETON

Ward 35, Leazes Wing, Royal Victoria Infirmary, Newcastle upon Tyne
NE1 4LP UK
R.J.Cooke@ncl.ac.uk

- 1 Tsang R, Lucas A, Uauy R, Zlotkin S, eds. *Nutritional needs of the pre-term infant: scientific basis and practical guidelines*. Pawling, NY: Caduceus Medical Publishers, Inc for Williams & Wilkins, 1993.
- 2 Embleton N, Pang N, Perring J, Cooke R. Systematic underfeeding of preterm infants on neonatal intensive care units [abstract]. *Pediatr Res* 1999;45:1653A.
- 3 Kashyap S, Heird W. Protein requirements of low birthweight, very low birthweight, and small for gestational age infants. In: Raiha NCR, ed. *Protein metabolism during infancy*. New York: Raven Press, 1994:133-51.
- 4 Cooke RJ, Ford A, Werkman S, Conner C, Watson D. Postnatal growth in infants born between 700 and 1,500 g. *J Pediatr Gastroenterol Nutr* 1993;16:130-5.
- 5 Ehrenkranz RA, Younes N, Lemons JA, *et al*. Longitudinal growth of hospitalized very low birth weight infants. *Pediatrics* 1999;104:280-9.
- 6 Fomon S. Size and growth. In: Fomon S, ed. *Nutrition of normal infants*. 2nd ed. St Louis: Mosby, 1993:36-84.
- 7 Gibson AT, Pearse RG, Wales JK. Knemometry and the assessment of growth in premature babies. *Arch Dis Child* 1993;69:498-504.
- 8 Kaempf DE, Pfluger MS, Thiele AM, Hermanussen M, Linderkamp O. Influence of nutrition on growth in premature infants: assessment by knemometry. *Ann Hum Biol* 1998;25:127-36.
- 9 Griffin JJ, Pang NM, Perring J, Cooke RJ. Knee-heel length measurement in healthy preterm infants. *Arch Dis Child Fetal Neonatal Ed* 1999;81:F50-5.
- 10 Dobbing J. *Early nutrition and later achievement*. London: Academic Press, 1987.
- 11 Dobbing J. *Developing brain and behaviour*. San Diego: Academic Press, 1997.
- 12 Sinisterra L, McKay H, McKay A, Gomez H, Korgi J. Response of malnourished preschool children to multidisciplinary intervention. In: Brozek J, ed. *Behavioural effect of energy and protein deficits*. DHEW publication no (NIH) 79-1906, 1979:229-38.
- 13 Lucas A, Morley R, Cole TJ, *et al*. Early diet in preterm babies and developmental status in infancy [see comments]. *Arch Dis Child* 1989;64:1570-8.
- 14 Morley R, Cole TJ, Powell R, Lucas A. Mother's choice to provide breast milk and developmental outcome. *Arch Dis Child* 1988;63:1382-5.
- 15 Wainwright PE, Ward GR. Early nutrition and behaviour: a conceptual framework for critical analysis of research. In: Dobbing J, ed. *Developing brain and behaviour: the role of lipids in infant formula*. San Diego: Academic Press, 1997:387-426.
- 16 Singer LT. Methodological considerations in longitudinal studies of infant risk. In: Dobbing J, ed. *Developing brain and behaviour: the role of lipids in infant formula*. San Diego: Academic Press, 1997:209-52.
- 17 Bornstein MH. Nutrition and development. In: Dobbing J, ed. *Developing brain and behaviour: the role of lipids in infant formula*. San Diego: Academic Press, 1997:475-509.
- 18 Johnson LR. Regulation of intestinal growth. In: Green M, Greene HL, eds. *The role of the intestinal tract in nutrient delivery*. Orlando: Academic Press, 1984:1-15.
- 19 Ostertag SG, LaGamma EF, Reisen CE, Ferrentino FL. Early enteral feeding does not affect the incidence of necrotizing enterocolitis. *Pediatrics* 1986;77:275-80.
- 20 Lucas A, Bloom SR, Aynsley-Green A. Postnatal surges in plasma gut hormones in term and preterm infants. *Biol Neonate* 1982;41:63-7.
- 21 Dunn L, Hulman S, Weiner J, Kliegman R. Beneficial effects of early hypocaloric enteral feeding on neonatal gastrointestinal function: preliminary report of a randomized trial. *J Pediatr* 1988;112:622-9.
- 22 Slagle TA, Gross SJ. Effect of early low-volume enteral substrate on subsequent feeding tolerance in very low birth weight infants. *J Pediatr* 1988;113:526-31.
- 23 McClure RJ, Newell SJ. Randomised controlled trial of trophic feeding and gut motility. *Arch Dis Child Fetal Neonatal Ed* 1999;80:F54-8.
- 24 Lucas A. Enteral nutrition. In: Tsang R, Lucas A, Uauy R, Zlotkin S, eds. *Nutritional needs of the preterm infant: scientific basis and practical guidelines*. Pawling, NY: Caduceus Medical Publishers, Inc for Williams & Wilkins, 1993:209-23.
- 25 Lucas A, Cole TJ. Breast milk and neonatal necrotising enterocolitis [see comments]. *Lancet* 1990;336:1519-23.
- 26 Ronnholm KA, Perheentupa J, Siimes MA. Supplementation with human milk protein improves growth of small premature infants fed human milk. *Pediatrics* 1986;77:649-53.
- 27 Carey DE, Rowe JC, Goetz CA, Horak E, Clark RM, Goldberg B. Growth and phosphorus metabolism in premature infants fed human milk, fortified human milk, or special premature formula. Use of serum procollagen as a marker of growth. *American Journal of Diseases in Children* 1987;141:511-15.
- 28 Greer FR, McCormick A. Improved bone mineralization and growth in premature infants fed fortified own mother's milk [published correction appears in *J Pediatr* 1988 Dec;113:1118]. *J Pediatr* 1988;112:961-9.
- 29 Polberger SK, Axelsson IA, Raiha NC. Growth of very low birth weight infants on varying amounts of human milk protein. *Pediatr Res* 1989;25:414-19.
- 30 Tyson JE, Lasky RE, Mize CE, *et al*. Growth, metabolic response, and development in very-low-birth-weight infants fed banked human milk or enriched formula. I. Neonatal findings. *J Pediatr* 1983;103:95-104.
- 31 Davies DP. Adequacy of expressed breast milk for early growth of preterm infants. *Arch Dis Child* 1977;52:296-301.
- 32 Gross SJ. Growth and biochemical response of preterm infants fed human milk or modified infant formula. *N Engl J Med* 1983;308:237-41.
- 33 Modanlou HD, Lim MO, Hansen JW, Sickles V. Growth, biochemical status, and mineral metabolism in very-low-birth-weight infants receiving fortified preterm human milk. *J Pediatr Gastroenterol Nutr* 1986;5:762-7.

- 34 Lucas A, Gore SM, Cole TJ, *et al.* Multicentre trial on feeding low birthweight infants: effects of diet on early growth. *Arch Dis Child* 1984;**59**:722–30.
- 35 Anderson GH, Atkinson SA, Bryan MH. Energy and macronutrient content of human milk during early lactation from mothers giving birth prematurely and at term. *American Journal of Clinical Nutrition* 1981;**34**:258–65.
- 36 Carlson SJ, Ziegler EE. Nutrient intakes and growth of very low birth weight infants. *J Perinatol* 1998;**18**:252–8.
- 37 Schanler RJ, Shulman RJ, Lau C. Feeding strategies for premature infants: beneficial outcomes of feeding fortified human milk versus preterm formula [comment]. *Pediatrics* 1999;**103**:1150–7.
- 38 Aggett PJ, Haschke F, Heine W, *et al.* Comment on the content and composition of lipids in infant formulas. ESPGAN Committee on Nutrition. *Acta Paediatr Scand* 1991;**80**:887–96.
- 39 Lucas A, Morley R, Cole TJ. Randomised trial of early diet in preterm babies and later intelligence quotient. *BMJ* 1998;**317**:1481–7.
- 40 Carlson SE, Cooke RJ, Rhodes PG, Peebles JM, Werkman SH. Effect of vegetable and marine oils in preterm infant formulas on blood arachidonic and docosahexaenoic acids. *J Pediatr* 1992;**120**:S159–67.
- 41 Auestad N, Montalto MB, Hall RT, *et al.* Visual acuity, erythrocyte fatty acid composition, and growth in term infants fed formulas with long chain polyunsaturated fatty acids for one year. Ross Pediatric Lipid Study. *Pediatr Res* 1997;**41**:1–10.
- 42 Carlson SE, Cooke RJ, Werkman SH, Tolley EA. First year growth of preterm infants fed standard compared to marine oil n-3 supplemented formula. *Lipids* 1992;**27**:901–7.
- 43 Jensen CL, Prager TC, Fraley JK, Chen H, Anderson RE, Heird WC. Effect of dietary linoleic/alpha-linolenic acid ratio on growth and visual function of term infants [see comments]. *J Pediatr* 1997;**131**:200–9.
- 44 Hay WW, Jr, Lucas A, Heird WC, *et al.* Workshop summary: nutrition of the extremely low birth weight infant. *Pediatrics* 1999;**104**:1360–8.
- 45 Polgar G, Kong GP. The nasal resistance of newborn infants. *J Pediatr* 1965;**67**:557–67.
- 46 Stocks J. Effect of nasogastric tubes on nasal resistance during infancy. *Arch Dis Child* 1980;**55**:17–21.
- 47 Greenspan JS, Abbasi S, Bhutani VK. Sequential changes in pulmonary mechanics in the very low birth weight (less than or equal to 1000 grams) infant. *J Pediatr* 1988;**113**:732–7.
- 48 van Someren V, Linnert SJ, Stothers JK, Sullivan PG. An investigation into the benefits of resiting nasogastric feeding tubes. *Pediatrics* 1984;**74**:379–83.
- 49 Aynsley-Green A, Adrian TE, Bloom SR. Feeding and the development of enteroincretory hormone secretion in the preterm infant: effects of continuous gastric infusions of human milk compared with intermittent boluses. *Acta Paediatr Scand* 1982;**71**:379–83.
- 50 Toce SS, Keenan WJ, Homan SM. Enteral feeding in very-low-birth-weight infants. A comparison of two nasogastric methods. *American Journal of Diseases of Children* 1987;**141**:439–44.
- 51 Silvestre MA, Morbach CA, Brans YW, Shankaran S. A prospective randomized trial comparing continuous versus intermittent feeding methods in very low birth weight neonates [see comments]. *J Pediatr* 1996;**128**:748–52.
- 52 Blondheim O, Abbasi S, Fox WW, Bhutani VK. Effect of enteral gavage feeding rate on pulmonary functions of very low birth weight infants. *J Pediatr* 1993;**122**:751–5.
- 53 Hammerman C, Kaplan M. Oxygen saturation during and after feeding in healthy term infants. *Biol Neonate* 1995;**67**:94–9.
- 54 Wilkinson A, Yu VY. Immediate effects of feeding on blood-gases and some cardiorespiratory functions in ill newborn infants. *Lancet* 1974;**1**:1083–5.
- 55 Wilkinson AR. Adverse effects of nasogastric feeding tubes and the management of recurrent apnoea [letter]. *Arch Dis Child* 1982;**57**:161.
- 56 Yu VY, Rolfe P. Effect of feeding on ventilation and respiratory mechanics in newborn infants. *Arch Dis Child* 1976;**51**:310–13.
- 57 Krauss AN, Brown J, Waldman S, Gottlieb G, Auld PA. Pulmonary function following feeding in low-birth-weight infants. *American Journal of Diseases of Children* 1978;**132**:139–42.
- 58 Patel BD, Dinwiddie R, Kumar SP, Fox WW. The effects of feeding on arterial blood gases and lung mechanics in newborn infants recovering from respiratory disease. *J Pediatr* 1977;**90**:435–8.
- 59 Heldt GP. The effect of gavage feeding on the mechanics of the lung, chest wall, and diaphragm of preterm infants. *Pediatr Res* 1988;**24**:55–8.
- 60 Pitcher-Wilmott R, Shutack JG, Fox WW. Decreased lung volume after nasogastric feeding of neonates recovering from respiratory disease. *J Pediatr* 1979;**95**:119–21.
- 61 Nelle M, Hoecker C, Linderkamp O. Effects of bolus tube feeding on cerebral blood flow velocity in neonates. *Arch Dis Child Fetal Neonatal Ed* 1997;**76**:F54–6.
- 62 Cooke RJ, Griffin IJ, McCormick K, *et al.* Feeding preterm infants after hospital discharge: effect of dietary manipulation on nutrient intake and growth. *Pediatr Res* 1998;**43**:355–60.
- 63 Georgieff MK, Mills MM, Lindeke L, Iverson S, Johnson DE, Thompson TR. Changes in nutritional management and outcome of very-low-birth-weight infants. *Am J Dis Child* 1989;**143**:82–5.
- 64 Casey PH, Kraemer HC, Bernbaum J, *et al.* Growth patterns of low birth weight preterm infants: a longitudinal analysis of a large, varied sample. *J Pediatr* 1990;**117**:298–307.
- 65 Casey PH, Kraemer HC, Bernbaum J, Yogman MW, Sells JC. Growth status and growth rates of a varied sample of low birth weight, preterm infants: a longitudinal cohort from birth to three years of age. *J Pediatr* 1991;**119**:599–605.
- 66 Hack M, Merkatz IR, McGrath SK, Jones PK, Fanaroff AA. Catch-up growth in very-low-birth-weight infants. Clinical correlates. *American Journal of Diseases in Children* 1984;**138**:370–5.
- 67 Wright K, Dawson JP, Fallis D, Vogt E, Lorch V. New postnatal growth grids for very low birth weight infants. *Pediatrics* 1993;**91**:922–6.
- 68 Lucas A, Bishop NJ, King FJ, Cole TJ. Randomised trial of nutrition for preterm infants after discharge [see comments]. *Arch Dis Child* 1992;**67**:324–7.
- 69 Cooke RJ, McCormick K, Griffin IJ, *et al.* Feeding preterm infants after hospital discharge: effect of diet on body composition. *Pediatr Res* 1999;**46**:461–4.