Non-invasive assessment of shunt and ventilation/perfusion ratio in neonates with pulmonary failure

H L Smith, J G Jones

Abstract

Aims—To make non-invasive measurements of right to left (R-L) shunt and reduced ventilation/perfusion ratio (V_A/Q) in neonates with pulmonary failure and to examine sequential changes in these variables after treatment.

Methods—Twelve neonates with pulmonary failure were studied. They ranged in gestational age from 24 to 37 (median 27) weeks and were 1–39 (median 4) days old. Shunt and reduced V_A/Q were derived from their effects on the relation between inspired oxygen pressure (P_{IO_2}) and arterial oxygen saturation measured with a pulse oximeter (SpO_2). Pairs of P_{IO_2} v SpO_2 data points were obtained by varying P_{IO_2} in a stepwise fashion. A computer algorithm based on a model of pulmonary gas exchange fitted a curve to these data. With P_{IO_2} on the abscissa, an increase in shunt produced a downward movement of the curve, whereas reducing V_A/Q to < 0.8 shifted the curve to the right. The right shift gives a variable that is inversely related to V_A/Q , the $P_{IO_2} - P_{cO_2}$ difference, where P_{cO_2} is mixed capillary oxygen pressure.

Results—Ten of the 12 infants on the first study day had large shunts (range 5.9–31.0%, median 19.9%, normal $< 8\%$) and large $P_{IO_2} - P_{cO_2}$ differences (range 9.7–64.4 kPa, median 19.8 kPa, normal < 7 kPa) equivalent to a median V_A/Q of 0.2 (normal median $V_A/Q = 0.8$). Sequential improvement in shunt and V_A/Q were shown in most infants after treatment.

Sudden large changes in these variables were shown in two infants.

Conclusion—This simple non-invasive method distinguishes between shunt and reduced V_A/Q in neonates with pulmonary failure.

(Arch Dis Child Fetal Neonatal Ed 2001;85:F127–F132)

Keywords: shunt; V_A/Q ; non-invasive; oxygen saturation

A reduced ventilation/perfusion ratio (V_A/Q) and an increased shunt cause quite different, but characteristic, effects on the relation between inspired oxygen pressure (P_{IO_2}) and arterial oxygen saturation (SpO_2).^{1,2} A reduced V_A/Q in the absence of shunt reduces both alveolar oxygen (PAO_2) and oxygen saturation but increasing the P_{IO_2} washes out alveolar nitrogen, restores PAO_2 , and corrects desaturation. In contrast, an isolated shunt reduces oxygen saturation, but increasing P_{IO_2} does not correct this because the shunted blood is never exposed to the increased P_{IO_2} . The non-shunted blood is almost fully saturated, and even administering 100% oxygen adds only a small amount of dissolved oxygen, which cannot compensate for the effect of the shunt.

It may be important in neonates to differentiate between a reduced V_A/Q and a shunt. For example, in an animal model of surfactant deficiency when surfactant was administered into the airways either as a bolus or as an aerosol, V_A/Q worsened with the former but improved with the latter.³ There may also be a prognostic significance of different degrees of

Figure 1 (A) Increasing shunt (from 0 to 40%) lowers the curve. (B) Reducing V_A/Q from 0.8 to 0.1 shifts the curve to the right. The right shift of each P_{IO_2} v SpO_2 curve from the position of the dissociation curve (dashed line) is the $P_{IO_2} - P_{cO_2}$ difference in kPa which includes P_{aCO_2}/R . The 0.8 curve represents the normal lung which intercepts a P_{IO_2} of 21 kPa (vertical line) at 97% SpO_2 .

Cambridge University
Department of
Anaesthesia,
Addenbrooke's
Hospital, Cambridge,
UK
H L Smith
J G Jones

Correspondence to:
Professor Jones, Department
of Anaesthesia, Cambridge
University, Level 4,
Addenbrooke's Hospital,
Cambridge CB2 2QQ, UK
gareth@garjones.demon.co.uk

Accepted 14 May 2001

Table 1 Summary of the clinical information on the first day of the study

Patient code	Days after delivery	Gestation (weeks)	Birth weight (kg)	Admission diagnosis
A	4	37	3.94	Parvovirus infection Anaemia Hydramnios
B	33	28	0.86	RDS
C	8	27	0.69	RDS Surfactant × 2
D	3	31	1.0	RDS Surfactant × 3
E	2	27	0.86	RDS Surfactant × 4
F	39	29	0.84	RDS
G	7	31	1.56	RDS Surfactant
H	1	26	0.82	RDS Surfactant
I	4	26	0.68	RDS Surfactant
J	26	27	0.82	RDS Surfactant
K	1	24	0.55	RDS Surfactant
L	3	24	0.71	RDS, PDA Surfactant

RDS, Respiratory distress syndrome; PDA, patent ductus arteriosus.

V_A/Q and shunt. However, existing methods of quantifying the relative contributions to oxygen desaturation of V_A/Q and shunt are not suitable for routine clinical application in neonates.⁴⁻⁸

We have described^{1,2,9} a simple non-invasive method for the calculation of shunt and reduced V_A/Q in adults which entails the simultaneous measurement of two variables, the inspired oxygen partial pressure (P_{IO_2}) and oxyhaemoglobin saturation using pulse oximetry (SpO_2). By changing the inspired oxygen partial pressure, pairs of P_{IO_2} and SpO_2 values are obtained. The curve relating P_{IO_2} to SpO_2 reflects the shape of the dissociation curve but always lies to the right, and its shape and position vary considerably when gas exchange is impaired.^{1,2,9-11} Thus, with P_{IO_2} on the abscissa:

- Increasing the shunt from 0 to 40% displaces the P_{IO_2} v SpO_2 curve downwards (fig 1A). This reflects the inability of an increase in P_{IO_2} to compensate for an increase in shunt.
- Reducing V_A/Q from 0.8 to 0.1 shifts the curve to the right (fig 1B) because the oxygen gradient increases between P_{IO_2} and P_{AO_2} (or P_{cO_2} , the mixed capillary oxygen). Increasing alveolar CO_2 also shifts the curve to the right by P_{CO_2}/R , where R is the gas exchange ratio. The right shift due to P_{CO_2}/R is small, typically 6 kPa, compared with the effect of reducing V_A/Q from 0.8 to 0.1, which causes a shift of 28 kPa.
- Note that, in fig 1B, the P_{IO_2} v SpO_2 curve representing a V_A/Q of 0.8 corresponds to the curve in normal subjects and gives an SpO_2 of 97% breathing air.

We studied a group of neonates with pulmonary failure to see if we could derive shunt and V_A/Q from the P_{IO_2} v SpO_2 data pairs despite the expected circulatory and ventilatory instabilities in such infants. We then examined the day to day changes to see if we could show

Figure 2 Inverse relation between V_A/Q and the right shift of the P_{IO_2} v SpO_2 curve.

Figure 3 Example of data analysis from infant I. Curve D, the dissociation curve, corresponds to a 0% shunt (see fig 1A) and curve SH to a 21.6% shunt. When curve SH is moved to the right by a $P_{IO_2} - P_{cO_2}$ difference of 26.6 kPa (dashed arrow), it superimposes the data points (closed circles) to give curve Mod.

sequential improvements in shunt and V_A/Q with continuing treatment.

Methods

PATIENTS

Twelve neonates, each with pulmonary failure requiring mechanical ventilation, were studied after local ethics committee approval had been obtained and informed consent obtained from the parents. In seven of these babies, up to four repeat studies were performed at intervals of up to 23 days.

The babies were being treated in the Neonatal Intensive Care Unit, Rosie Maternity Hospital, Addenbrooke's Hospital, Cambridge. All but one of the infants were preterm with intrauterine growth retardation and were admitted for ventilation for respiratory distress syndrome (table 1). All infants were being ventilated through a tracheal tube. A consultant neonatologist not involved in this study managed ventilatory support. One infant, delivered at 37 weeks gestation (infant A), had developed hydrops foetalis after parvovirus infection in utero. In most infants, surfactant had been administered when appropriate through a tracheal tube. Arterial blood gas and

Table 2 Details of shunt and right shift of the P_{iO_2} v SpO_2 curve

Patient*	Shunt (%)	Shunt 95% CI (%)	Shift (kPa), (V_A/Q)	Shift 95% CI (kPa)	r^2
A 4	30.7	29.3 to 32.2	64.4 (<0.05)	61.8 to 67.1	0.73
A 5	27.3	26.4 to 28.3	16.8 (0.27)	14.5 to 19.1	0.78
A 10	20.1	18.6 to 21.8	11.4 (0.49)	8.3 to 14.5	0.89
B	10.8	7.8 to 13.7	13.5 (0.4)	11.5 to 15.6	0.83
C	20.1	18.2 to 21.9	17.5 (0.25)	16.1 to 18.8	0.8
D	17.1	16.0 to 18.2	19.8 (0.2)	18.7 to 20.8	0.93
E 2	19.7	17.0 to 22.5	58.5 (<0.05)	55.1 to 61.9	0.92
E 3	23.1	22.3 to 23.8	26.9 (0.14)	25.5 to 28.4	0.86
E 9	23.5	22.7 to 24.3	56.2 (<0.05)	55.2 to 57.1	0.93
E 25	16.1	13.2 to 18.8	29.0 (0.13)	26.8 to 31.3	0.72
F 39	26.5	25.6 to 27.5	34.6 (0.1)	31.5 to 37.7	0.97
F 43	11.2	9.3 to 13.2	10.3 (0.5)	7.1 to 13.5	0.94
G	21.5	20.2 to 22.8	19.1 (0.2)	16.2 to 21.95	0.92
H 1	31.0	30.0 to 32.0	53.9 (<0.05)	50.5 to 57.3	0.99
H 4	5.9	5.0 to 6.7	9.7 (0.55)	9.0 to 10.4	0.99
I 4	21.6	20.8 to 22.4	26.6 (0.15)	24.9 to 28.2	0.96
I 8	7.3	5.3 to 9.2	13.7 (0.4)	13.4 to 14.1	0.93
J 26	13.3	11.8 to 14.7	22.9 (0.18)	22.2 to 23.5	0.93
J 27	19.0	18.0 to 20.0	22.0 (0.18)	22.5 to 23.5	0.95
K	28.0	27.0 to 28.8	14.2 (0.35)	12.2 to 16.2	0.97
L 3	17.1	16.1 to 18.0	18.6 (0.22)	17.6 to 19.5	0.95
L 7	18.4	17.6 to 19.2	19.7 (0.2)	17.9 to 21.5	0.96

V_A/Q is the ventilation to perfusion ratio in the ventilated but poorly perfused compartment and was derived from the mean shift—that is, $P_{iO_2} - P_{eO_2}$ in kPa—using fig 2. The number after the patient code is the number of days after birth for repeat studies. r^2 indicates the goodness of fit of the gas exchange model to the P_{iO_2} v SpO_2 data points.⁹

haemoglobin values were recorded independently as part of routine clinical management. Monitoring included measurement of inspired oxygen concentration and arterial saturation using the integral gas analyser in the Draeger Babylog 8000 ventilator (to give F_{iO_2} or P_{iO_2}) and an Ohmeda Biox 3700e pulse oximeter (to give SpO_2). The Draeger automatically calibrates itself every 24 hours—for example, at

1100, using piped air and 100% oxygen. Each calibration takes 90 seconds and the instrument is accurate to 5% full scale deflection—that is, 21(1)kPa. The instrument sounds an alarm if the calibration exceeds this range. The routine arterial blood gas results were used periodically to check the pulse oximeter reading. Pulse oximeters are accurate to $\pm 1\%$ in the normal range but much less at low SpO_2 (see Discussion).

PROCEDURE

P_{iO_2} was varied in steps to cause the SpO_2 to vary between 85% and 96%. This level of oxygen has been used in previous research in the measurement of cerebral blood flow using near infrared spectroscopy.¹² The P_{iO_2} was never reduced below 21 kPa.

DATA ANALYSIS AND INTERPRETATION

The pairs of P_{iO_2} and SpO_2 data points for each study were recorded and analysed using a computer algorithm which gave a curve representing a single solution for each subject's data set.⁹ This curve is not a statistical line of best fit but is based on a model of gas exchange used by the computer algorithm to describe the P_{iO_2} v SpO_2 data points by two variables of gas exchange:

- virtual shunt using the shunt equation;
- the difference between inspired oxygen and mixed pulmonary capillary oxygen pressures ($P_{iO_2} - P_{eO_2}$), a function of reduced V_A/Q and P_{CO_2}/R .

Figure 4 The P_{iO_2} v SpO_2 curves from all 12 infants on the first day that they were studied. In every case the curve is shifted to the right of the line marking a P_{iO_2} of 21 kPa, indicating a large reduction in V_A/Q ratio. In infant L, there are two data sets, before (closed circles) and after tracheal aspiration (open circles).

Figure 5 Sequence of changes in P_{IO_2} v S_{PO_2} data points and model fit in seven infants who were studied sequentially.

Figure 2 gives the inverse relation between right shift—that is, $P_{IO_2} - P_{cO_2}$ —and V_A/Q .

The term “virtual” is used if shunt and V_A/Q are calculated with assumed values of haemoglobin, carbon dioxide (P_{CO_2}), respiratory quotient (R), and arteriovenous oxygen difference when the actual values are not available. In this study, R and arteriovenous oxygen difference were assumed.

Results

Table 1 shows the admission diagnosis, gestational age, days after delivery, and weight of the 12 infants.

Figure 3 shows a P_{IO_2} v S_{PO_2} data set from infant I on the first study day (closed circles). A model curve (Mod) was fitted to these data by the computer algorithm which calculated a shunt value of 21.6%. A 21.6% shunt is shown by curve SH which, in the absence of V_A/Q abnormality, asymptotes the dissociation curve D as shown in fig 3. The algorithm fits the SH curve to the data points by displacing it to the right of D by 26.6 kPa. This displacement is the $P_{IO_2} - P_{cO_2}$ difference, which represents the combined effects of a reduced V_A/Q plus P_{CO_2}/R .

Table 2 gives the calculated values of shunt and right shift ($P_{IO_2} - P_{cO_2}$) for all these infants on the first study day. This also shows the V_A/Q derived from $P_{IO_2} - P_{cO_2}$ and the r^2 value, which is the goodness of fit of the model to the data points.⁹

The values for shunt and right shift—that is, $P_{IO_2} - P_{cO_2}$ difference and thus V_A/Q —for each baby shown in table 2 all showed a good fit to the P_{IO_2} v S_{PO_2} data points indicated by the r^2 value.⁹ Figure 4 shows these data points and the computed curves for all 12 infants on the first study day. All these data are plotted on a grid where vertical and horizontal lines mark a P_{IO_2} of 21 kPa (air at sea level) and an S_{PO_2} of 90% respectively. Note that the normal curve, with a median V_A/Q of 0.8, intersects the 21 kPa line at about 97% S_{PO_2} , and this is equivalent to moving the dissociation curve (D in fig

3) to the right by about 6–7 kPa—for example, a normal P_{CO_2}/R . The 95% confidence intervals of the fit of the curve are shown for shunt (upper part of the curve) and right lateral shift (lower part of the curve).

Note that in every case the curve was shifted to the right of the vertical line marking a P_{IO_2} of 21 kPa (air at sea level). There was a good fit of the curve to the data points despite the considerable shunt and right shift, the most severe examples of which were in infants A, E, F, and H. Two curves are shown for infant L. The one on the right was recorded first and that on the left 20 minutes after tracheal suction (see discussion).

Seven babies were studied on more than one occasion (fig 5). Infant A, a neonate with hydrops foetalis secondary to anaemia following an intrauterine parvovirus infection, showed a considerable improvement from day 4 to day 5, and this improvement continued to day 10. The overall reduction in shunt was from 30.7% to 20.1%. More dramatically there was a decrease in $P_{IO_2} - P_{cO_2}$ from 64.4 kPa to 11.4 kPa, equivalent to an improvement in V_A/Q from less than 0.05 to 0.49.

Infant E showed a very unstable clinical pattern with improvement (day 2 to day 3) followed by deterioration (day 3 to day 9). Eventually (day 25) there was the lowest value of shunt, 16.1%, but there was still a very large right shift—that is, a large $P_{IO_2} - P_{cO_2}$ difference—which our gas exchange model equated to the perfused alveoli having a V_A/Q of 0.13.

Infants F and H showed a considerable improvement over the days they were studied, with a reduction in both the shunt and right shift to near normal values.

Infant I, a 26 week old neonate with infant respiratory distress syndrome, was the only baby studied while dependent on a high frequency oscillator. From day 4 to day 8 there was an improvement in both shunt and right shift (figs 4 and 5 and table 2). The shunt

decreased from 21.6% to 7.3% (normal), with the right shift being reduced from 26.6 kPa to 13.7 kPa—that is, a V_A/Q of 0.14 improving to 0.4.

Infant J showed deterioration in shunt from day 26 to 27. A normal echo ruled out a patent ductus arteriosus. However, this baby was hypotensive and had suspected necrotising enterocolitis. Infant L showed a more stable picture, with little change between days 3 and 7.

Discussion

Our gas exchange model derives two variables of impaired gas exchange: shunt and the $P_{IO_2} - P_{cO_2}$ difference representing the effect of a reduced V_A/Q ($0.05 < V_A/Q < 1$). In all 12 infants, we found a good fit of the model to the P_{IO_2} v SpO_2 data points. With a few exceptions the curves were stable—that is, a single shunt equation fitted all the P_{IO_2} v SpO_2 data points obtained from one subject during one data collection period. The results were similar to those seen in a variety of adult patients with pulmonary failure during anaesthesia,⁹ thoracotomy,¹⁰ and after fat embolism.¹¹

Reducing median V_A/Q below unity causes a considerable increase in oxygen gradient.¹ However, some workers, as recently as 1995, have assumed that the entire oxygen gradient is explained only by a shunt.^{13,14} This simplistic approach ignores the effect of an altered V_A/Q distribution on the oxygen gradient and lumps together the effects of a reduced V_A/Q with those of a shunt. In infants with respiratory distress syndrome,^{4,5} the nitrogen gradient method shows a considerable reduction in V_A/Q , which explains about 30% of the oxygen gradient.⁵ Not only were V_A/Q ratios as low as 0.05, but also, in one case, this accounted for almost the whole of an oxygen gradient of 65 kPa.⁵ This large gradient is similar in magnitude to the $P_{IO_2} - P_{cO_2}$ difference seen in our infant A (table 2), which gave a V_A/Q of 0.05.

In some of our infants, the increase in shunt ($> 30\%$) and reduction in V_A/Q (< 0.05) were considerable, the gas exchange abnormalities being as great as those previously seen during thoracic anaesthesia in adults in whom a considerable part of the lung was collapsed.¹⁰

Some infants showed considerable variability in the P_{IO_2} v SpO_2 data. Three examples are as follows. (a) Infant L at the start of the data collection period showed a very large reduction in V_A/Q causing a right shift of 40 kPa but with a very small shunt (fig 4). After tracheal aspiration, there was an improvement in V_A/Q but with an increase in shunt from 6.6 to 17% (open circles fig 4). (b) In infant J, there were two episodes of sudden hypoxaemia when the SpO_2 fell precipitously necessitating the inspired oxygen concentration to be increased to 60%. These episodes may have been caused either by a transient opening of an R-L shunt or, more likely, a fall in cardiac output causing an increased shunt effect.⁹ (c) At times, in an otherwise stable data set, there was more than 6% variability in SpO_2 . This was particularly so

when we were measuring SpO_2 on the steep part of the P_{IO_2} v SpO_2 curve. This was consistent with our previous observations in post-operative adult patients¹⁵, for whom we reported a very unstable pattern of SpO_2 when the P_{IO_2} lay on the steep part of the P_{IO_2} v SpO_2 curve.

Severinghaus and Naifeh¹⁶ examined the accuracy of pulse oximeters. They found that the probe is accurate to $\pm 1\%$ in the normal range, but, with a sudden hypoxic episode, an SpO_2 reading of 55% may be 7% lower than the actual SaO_2 value. Because such inaccuracy occurs only on the very steep part of the oxyhaemoglobin dissociation curve, this has little effect on the derivation of the P_{IO_2} v curve in our studies. The derived P_{IO_2} v SpO_2 curve based on our model of gas exchange obviously does not fit every data point, and the r^2 value and the 95% confidence interval of shunt and right shift of the curve describe this variability (table 2).

The fetal dissociation curve is shifted to the left of the adult curve with a P_{50} of 2.7 kPa compared with the adult value of 3.6 kPa. This difference is reduced after birth and after blood transfusion. This means that the median value of the variable $P_{IO_2} - P_{cO_2}$, which for the whole study was 19.8 kPa, would have been underestimated by less than 1 kPa.

The model is sensitive to a reduced haemoglobin concentration and low cardiac output.⁹ The haemoglobin value for each infant was recorded from routine samples taken on the day of the study, and these values were used in the calculation of shunt and V_A/Q . A cardiac output equivalent to an arteriovenous oxygen difference of 5 ml/dl was assumed.

We conclude that the method shows that the gas exchange model based on the oxyhaemoglobin dissociation curve provides a good fit to the P_{IO_2} v SpO_2 data in the circumstances of our study. The model gave a non-invasive measurement of shunt and reduced V_A/Q in neonates during mechanical ventilation and allowed the prediction of the effect on SpO_2 of changing P_{IO_2} .

We thank Drs James Powell and Jackie Gedney for their assistance with this study. Dr Wilf Kelsall was consultant neonatologist at Addenbrooke's hospital in whose care were the infants described in this paper.

- 1 Roe PG, Jones JG. Analysis of factors which affect the relationship between inspired oxygen partial pressure and arterial oxygen saturation. *Br J Anaesth* 1993;71:488–94.
- 2 Sapsford DJ, Jones JG. Predicting changes in oxygen saturation using pulse oximetry. *Br J Anaesth* 1988;61:497P.
- 3 Schermuly RT, Gunther A, Weissman N, et al. Differential impact of ultrasonically nebulized versus tracheal-instilled surfactant on ventilation-perfusion (V_A/Q) mismatch in a model of acute lung injury. *Am J Respir Crit Care Med* 2000;161:152–9.
- 4 Corbet AJS, Ross JA, Beaudry PH, et al. Ventilation-perfusion relationships as assessed by aADN₂ in hyaline membrane disease. *J Appl Physiol* 1974;36:74–81.
- 5 Hand IL, Shepard EK, Krauss AN, et al. Ventilation-perfusion abnormalities in the preterm infant with hyaline membrane disease: a two-compartment model of the neonatal lung. *Pediatr Pulmonol* 1990;9:206–13.
- 6 Thibeault DW, Poblete E, Auld PAM. Alveolar-arterial oxygen difference in premature infants breathing 100 percent oxygen. *J Pediatr* 1967;71:814–24.
- 7 Roca J, Wagner PD. Principles and information content of the multiple inert gas elimination technique. *Thorax* 1993;49:815–24.
- 8 Melot C. Ventilation-perfusion relationships in acute respiratory failure. *Thorax* 1994;49:1251–8.

- 9 Sapsford DJ, Jones JG. The PiO_2 vs. SpO_2 diagram: a non-invasive measure of pulmonary oxygen exchange. *Eur J Anaesth* 1995;12:375–86.
- 10 de Gray L, Rush EM, Jones JG. A non-invasive method for evaluating the effect of thoracotomy on shunt and ventilation perfusion inequality. *Anaesthesia* 1997;52:630–5.
- 11 Burnstein RM, Newell JP, Jones JG. Sequential changes in gas exchange following traumatic fat embolism. *Anaesthesia* 1998;53:369–81.
- 12 Edwards AD, Wyatt JS, Richardson C, *et al.* Cotside measurement of cerebral blood flow in infants by near infrared spectroscopy. *Lancet* 1988;2:770–1.
- 13 Lawler PGP, Nunn JF. A reassessment of the validity of the iso-shunt graph. *Br J Anaesth* 1984;56:1325–35.
- 14 Hope DA, Jenkins BJ, Willis N, *et al.* Non-invasive estimation of venous admixture: validation of a new formula. *Br J Anaesth* 1995;74:538–43.
- 15 Roe PG, Gadelrab R, Sapsford D, *et al.* Intra operative gas exchange and post operative hypoxaemia. *Eur J Anaesth* 1997;14:203–10.
- 16 Severinghaus JW, Naifeh KH. Accuracy of response of six pulse oximeters to profound hypoxia. *Anesthesiology* 1987;67:551–8.