

# Don't stop now? How long should resuscitation continue at birth in the absence of a detectable heartbeat?

Dominic JC Wilkinson,<sup>1,2,3</sup> Ben Stenson<sup>4</sup>

Shah and colleagues from Western Australia present the latest in a series of papers, suggesting that outcomes for infants who have received prolonged resuscitation in the delivery room are more favourable than before.<sup>1</sup> They report 13 near-term infants who had an Apgar score of 0 at 10 min and were admitted to intensive care. Five of the 13 infants survived and 4 of the 5 appeared to have normal development at 1–2 years of age (one of the infants had hearing impairment). One of the five survivors developed severe spastic quadriplegia.<sup>1</sup>

Similar findings were reported recently in this journal by Kasdorf *et al.*<sup>2</sup> They reported nine infants managed in a New York hospital, and combined their results with data from three of the therapeutic hypothermia trials and one other previously published report.<sup>3</sup> In total, Kasdorf *et al.*<sup>2</sup> analysed a cohort of 90 infants with an Apgar score of 0 at 10 min who were admitted to intensive care. Fifty per cent of the infants survived to discharge from hospital, and 49% of the survivors were developmentally normal at follow-up at 1–2 years.<sup>2</sup> Longer-term developmental outcome data are available for a subgroup of these infants who were enrolled in the US NICHD cooling trial.<sup>4</sup> Twenty four infants in that trial had an Apgar score of 0 at 10 min. Eleven (46%) survived to age 6–7 years, and 5/11 (46%) had mild or no disability at follow-up.<sup>4</sup>

Considered together, these results are in striking contrast to earlier studies and should cause us to question current recommendations in consensus resuscitation guidelines. A systematic review in 2007 identified 94 infants from eight reports.<sup>5</sup> Seventy eight infants (83%) died, while 10/13 (76%) of survivors with available

long-term follow-up were severely impaired. The authors concluded that the outcome for infants with an Apgar score of 0 at 10 min was 'almost universally poor'. Based on this earlier report and other information, the 2010 ILCOR guidelines stated: 'It is appropriate to consider discontinuing resuscitation if there has been no detectable heart rate for 10 minutes'.<sup>6</sup> It is important to recognise that this statement is not a recommendation that resuscitation *should* always be discontinued and was qualified by factors that might be taken into consideration if resuscitation were to be continued. However, the inference is that discontinuing resuscitation at 10 min would be the usual approach.

## ARE CURRENT GUIDELINES APPROPRIATE?

The recent data have some important limitations. The reports only include infants who were admitted to intensive care. Infants in whom prolonged resuscitation did not achieve a sustained return of the circulation are excluded. The cooling trials also excluded infants who were moribund or who died within the first hours after birth. The overall survival rate for infants who respond to resuscitation after having an Apgar score of 0 at 10 min is likely to be somewhat lower than the rate reported here. The newer data were gathered in the context of a practice guided by the historically reported outcomes. If resuscitation and intensive care were continued in more infants with Apgar score of 0 at 10 min, resulting in more survivors, it is possible that the rate of severe disability among survivors might be greater. Death in infants with hypoxic-ischaemic encephalopathy usually follows a decision to limit life-sustaining treatment on the basis of poor neurological prognosis.<sup>7</sup> Longer-term outcomes would be likely to vary in practice settings where decision-making is different.

## WHAT ARE THE ETHICAL JUSTIFICATIONS FOR STOPPING RESUSCITATION?

### Is a normal outcome impossible?

One reason for limiting resuscitation at birth to 10 min may be a belief that

further resuscitation is futile because it is inevitable that the infant will either die or be profoundly impaired.<sup>8</sup> However, the recent papers indicate clearly that in the current era the outcome is not universally poor. Focusing just on the 66 infants in published series who have received therapeutic hypothermia, discontinuing resuscitation at 10 min would have led to the death of 18 infants who do not have severe disability, many of whom are developmentally normal.

### Is a 'poor' outcome highly likely?

An alternative justification for discontinuing resuscitation at 10 min is because the chance of a poor outcome is thought to be so high that resuscitation is not warranted. Here we need to be clear about the reason *why* it is considered too high. This should not be influenced by a high-mortality risk because in the absence of further resuscitation mortality will be universal. The reason here would therefore be a fear of generating additional survivors with severe disability.<sup>9</sup> From the point of view of parents, the outcomes of death or severe disability may have quite different significance. Some parents may fear the possibility of their child surviving in a state of severe impairment, while others may be much more afraid of their child dying even if there is a likelihood of survival with disability. There may be an unwelcome trade-off with increasing duration of resuscitation between avoiding death in a child who could survive with a favourable outcome and avoiding survival in a child who will have profound neuro-disability. Even if the risk of disability can be estimated, identifying a level of disability that is so severe that it is contrary to the child's best interests to provide treatment is extremely challenging.<sup>10</sup>

## WHAT SHOULD RESUSCITATION GUIDELINES RECOMMEND?

### Providing treatment in the best interests of the child

As with all medical treatment for newborn infants, clinicians should be guided by the best interests of the child.<sup>11</sup> It is ethical to withhold life-sustaining treatment where a child's life would be limited in quantity or quality, and treatment would therefore do more harm than good.<sup>11</sup> The evidence summarised above shows that it is far from clear that continued resuscitation beyond 10 min would be harmful. It is, therefore, highly questionable whether medical professionals should make a unilateral decision to cease resuscitation at 10 min in the absence of additional information.

<sup>1</sup>Faculty of Philosophy, Oxford Uehiro Centre for Practical Ethics, University of Oxford, Oxford, UK;

<sup>2</sup>Robinson Institute, Discipline of Obstetrics and Gynaecology, University of Adelaide, Adelaide, Australia;

<sup>3</sup>John Radcliffe Hospital, Oxford, UK;

<sup>4</sup>Neonatal Unit, Royal Infirmary of Edinburgh, Edinburgh, UK

**Correspondence to** Dr Dominic Wilkinson, Oxford Uehiro Centre for Practical Ethics, Suite 8, Littlegate House, St Ebbses St, Oxford OX1 1PT, UK; dominic.wilkinson@philosophy.ox.ac.uk

## Erring on the side of providing treatment

Faced with uncertainty, it is usually preferable to err on the side of providing life-saving treatment.<sup>11</sup> The recently revised Royal College of Paediatrics and Child Health guidelines around limitation of treatment note that 'Neonates should almost always be resuscitated in the labour ward, unless there is a clear decision to do otherwise made with the consent of parents'.<sup>11</sup> Given the uncertainty about outcome and a likely inability to determine the parents' wishes in a meaningfully informed manner, the default approach at 10 min should be to continue resuscitation, take the infant to intensive care if they have a return of circulation and at that stage to involve parents in decision-making about ongoing treatment.

The statistics summarised above, with an estimated chance of survival of 50% and a risk of severe disability in survivors of around 50%, are similar to the outcome figures for infants born alive at 23–24 weeks gestation.<sup>12</sup> In that setting, there is widespread acceptance that parents' views are crucial to decisions.<sup>13</sup> It would not be regarded as acceptable for clinicians to make decisions to limit treatment without knowing the parents' views. Of course, the distinctive feature of decisions about the duration of resuscitation at birth for term infants is that it is extremely difficult to involve parents fully in decision-making. The acuity of the situation, its unpredictability, the intense time-pressure and the involvement of the clinical team in the ongoing resuscitation efforts combine to make it extremely challenging to counsel parents and gauge their views.

A risk of longer resuscitation and of taking infants to intensive care is that this potentially leaves some families facing a more difficult decision that they would have preferred not to have faced about whether or not to withdraw treatment. Some families may be unwilling to acquiesce to a decision to stop life support. This is a particular challenge for religious or cultural groups that traditionally do not accept withdrawal of treatment.<sup>14</sup> However, there is no ethical difference between withholding and withdrawing life-sustaining treatment.<sup>15</sup> Clinicians must be guided primarily by the best interests of the infant. As the paper by Shah indicates, in intensive care units with access to prognostic tests such as MRI and EEG, and for communities and cultures that accept withdrawal of treatment, such a policy is associated with a low rate of survival with severe impairment.<sup>1</sup>

## If not 10 min, when?

On the basis of current evidence, the outcome for infants with an Apgar score of 0 at 10 min is not universally poor. In circumstances where neonatal intensive care (including therapeutic hypothermia) is available, resuscitation at birth should usually continue beyond 10 min. Infants should be supported with intensive care if their circulation is restored. Guidelines on resuscitation should be modified. The question then becomes *how long* resuscitation should continue beyond 10 min in the absence of a heartbeat. This is a much harder question because there are (to our knowledge) little published data on longer durations of resuscitation at birth and outcome. Kasdorf *et al*<sup>2</sup> have called for a registry of all infants with an Apgar score of 0 at 10 min, including infants for whom resuscitation is not successful. However, long-term outcome from any such registry is not likely to be available soon.

Although the series is small, the paper by Shah *et al* provides some pointers. Their five surviving infants all had their first heartbeat documented by 20 min. Of these five, four subsequently had a good outcome. No infant in their report who had their first heartbeat after 20 min of resuscitation survived.<sup>1</sup> It may be possible to obtain information about the timing of the first detectable heartbeat and the final outcome for infants from the other reports. In individual cases, other factors may already be known at the time of birth that should influence decision-making.

## CONCLUSIONS

An Apgar score of 0 at 10 min after birth is not a good enough predictor of outcome to be used as the main basis for decision-making about ongoing resuscitation. There is no clear answer to the question of how long resuscitation should continue after birth. We propose that in most circumstances, resuscitation at birth should continue until 20 min in the absence of a clinically detectable heartbeat. In the face of uncertainty about whether resuscitation should be discontinued, clinicians should opt to provide longer resuscitation, with later consideration of withdrawal of life-sustaining treatment if the clinical course indicates that the prognosis is poor. If practice changes in this way it will be very important for clinicians to audit and report their outcomes in order that the best possible information becomes available to inform future decision-making.

**Twitter** Follow Dominic Wilkinson at @NeonatalEthics

**Contributors** DW and BS conceived and planned this article. DW wrote the first draft and edited the manuscript. BS reviewed and edited the manuscript.

**Funding** DW was supported by a grant from the Wellcome Trust (086041/Z/08/Z) during the preparation of this manuscript.

**Competing interests** BS has been a member of the ILCOR neonatal group and is editor of the Fetal and Neonatal edition of *Archives of Disease in Childhood*. DW has contributed to recent RCPCH guidelines on decisions around life-sustaining treatment and is an associate editor of the *Journal of Medical Ethics*. The views expressed in this article are solely those of the authors and do not represent the views of any other organisation.

**Provenance and peer review** Commissioned; internally peer reviewed.

**Data sharing statement** DW had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**To cite** Wilkinson DJC, Stenson B. *Arch Dis Child Fetal Neonatal Ed* 2015;**100**:F476–F478.

Received 30 May 2015

Accepted 3 July 2015

Published Online First 3 August 2015



Editor's choice  
Scan to access more  
free content



► <http://dx.doi.org/10.1136/fetalneonatal-2014-307825>

*Arch Dis Child Fetal Neonatal Ed* 2015;**100**:F476–F478. doi:10.1136/archdischild-2015-308602

## REFERENCES

- Shah P, Anvekar A, McMichael J, *et al*. Outcomes of infants with Apgar score of zero at 10 minutes: the West Australian experience. *Arch Dis Child Fetal Neonatal Ed* 2015;**100**:F492–4.
- Kasdorf E, Luptook A, Azzopardi D, *et al*. Improving infant outcome with a 10 min Apgar of 0. *Arch Dis Child Fetal Neonatal Ed* 2015;**100**:F102–5.
- Sarkar S, Bhagat I, Dechert RE, *et al*. Predicting death despite therapeutic hypothermia in infants with hypoxic-ischaemic encephalopathy. *Arch Dis Child Fetal Neonatal Ed* 2010;**95**:F423–8.
- Natarajan G, Shankaran S, Luptook AR, *et al*. Apgar scores at 10 min and outcomes at 6–7 years following hypoxic-ischaemic encephalopathy. *Arch Dis Child Fetal Neonatal Ed* 2013;**98**:F473–9.
- Harrington DJ, Redman CW, Moulden M, *et al*. The long-term outcome in surviving infants with Apgar zero at 10 minutes: a systematic review of the literature and hospital-based cohort. *Am J Obstet Gynecol* 2007;**196**:463.e1–5.
- Perman JM, Wyllie J, Kattwinkel J, *et al*. Neonatal resuscitation: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Pediatrics* 2010;**126**:e1319–44.

- 7 Brecht M, Wilkinson DJ. The outcome of treatment limitation discussions in newborns with brain injury. *Arch Dis Child Fetal Neonatal Ed* 2015;100:F155–60.
- 8 Wilkinson DJC, Savulescu J. Knowing when to stop: futility in the ICU. *Curr Opin Anaesthesiol* 2011;24:160–5.
- 9 Janvier A, Mercurio MR. Saving vs creating: perceptions of intensive care at different ages and the potential for injustice. *J Perinatol* 2013;33:333–5.
- 10 Wilkinson D. *Death or disability? The Carmentis Machine and treatment decisions for critically ill children*. Oxford: OUP, 2013.
- 11 Larcher V, Craig F, Bhogal K, *et al*. Making decisions to limit treatment in life-limiting and life-threatening conditions in children: a framework for practice. *Arch Dis Child* 2015;100(Suppl 2):s1–23.
- 12 Moore T, Hennessy EM, Myles J, *et al*. Neurological and developmental outcome in extremely preterm children born in England in 1995 and 2006: the EPICure studies. *BMJ* 2012;345:e7961.
- 13 Wilkinson AR, Ahluwalia J, Cole A, *et al*. Management of babies born extremely preterm at less than 26 weeks of gestation: a framework for clinical practice at the time of birth. *Arch Dis Child Fetal Neonatal Ed* 2009;94:F2–5.
- 14 Sprung CL, Paruk F, Kissoon N, *et al*. The Durban World Congress Ethics Round Table Conference Report: I. Differences between withholding and withdrawing life-sustaining treatments. *J Crit Care* 2014;29:890–5.
- 15 Wilkinson D, Savulescu J. A costly separation between withdrawing and withholding treatment in intensive care. *Bioethics* 2012;26:32–48.