OPEN ACCESS

Estimating Global Burden of Disease due to congenital anomaly: an analysis of European data

Breidge Boyle,^{1,2} Marie-Claude Addor,³ Larraitz Arriola,⁴ Ingeborg Barisic,⁵ Fabrizio Bianchi,⁶ Melinda Csáky-Szunyogh,⁷ Hermien E K de Walle,⁸ Carlos Matias Dias,⁹ Elizabeth Draper,¹⁰ Miriam Gatt,¹¹ Ester Garne,¹² Martin Haeusler,¹³ Karin Källén,¹⁴ Anna Latos-Bielenska,¹⁵ Bob McDonnell,¹⁶ Carmel Mullaney,¹⁷ Vera Nelen,¹⁸ Amanda J Neville,¹⁹ Mary O'Mahony,²⁰ Annette Queisser-Wahrendorf,²¹ Hanitra Randrianaivo,²² Judith Rankin,²³ Anke Rissmann,²⁴ Annukka Ritvanen,²⁵ Catherine Rounding,²⁶ David Tucker,²⁷ Christine Verellen-Dumoulin,²⁸ Diana Wellesley,²⁹ Ben Wreyford,³⁰ Natalia Zymak-Zakutnia,³¹ Helen Dolk¹

For numbered affiliations see end of article.

Correspondence to

Professor Helen Dolk, WHO Collaborating Centre for the Epidemiologic Surveillance of Congenital Anomalies, Centre for Maternal, Fetal and Infant Research, Institute of Nursing and Health Research, Ulster University, Shore Road, Newtownabbey BT37 0QB, UK; h.dolk@ulster.ac.uk

Received 18 August 2016

Revised 15 March 2017

Accepted 18 April 2017

Published Online First

30 June 2017

ABSTRACT

Objective To validate the estimates of Global Burden of Disease (GBD) due to congenital anomaly for Europe by comparing infant mortality data collected by EUROCAT registries with the WHO Mortality Database, and by assessing the significance of stillbirths and terminations of pregnancy for fetal anomaly (TOPFA) in the interpretation of infant mortality statistics.

Design, setting and outcome measures EUROCAT is a network of congenital anomaly registries collecting data on live births, fetal deaths from 20 weeks' gestation and TOPFA. Data from 29 registries in 19 countries were analysed for 2005–2009, and infant mortality (deaths of live births at age <1 year) compared with the WHO Mortality Database. Eight EUROCAT countries were excluded from further analysis on the basis that this comparison showed poor ascertainment of survival status.

Results According to WHO, 17%–42% of infant mortality was attributed to congenital anomaly. In 11 EUROCAT countries, average infant mortality with congenital anomaly was 1.1 per 1000 births, with higher rates where TOPFA is illegal (Malta 3.0, Ireland 2.1). The rate of stillbirths with congenital anomaly was 0.6 per 1000. The average TOPFA prevalence was 4.6 per 1000, nearly three times more prevalent than stillbirths and infant deaths combined. TOPFA also impacted on the prevalence of postneonatal survivors with non-lethal congenital anomaly.

Conclusions By excluding TOPFA and stillbirths from GBD years of life lost (YLL) estimates, GBD underestimates the burden of disease due to congenital anomaly, and thus declining YLL over time may obscure lack of progress in primary, secondary and tertiary prevention.

INTRODUCTION

According to the 2010 Global Burden of Disease (GBD) study,^{1,2} congenital anomalies accounted for 510 400 deaths worldwide in 2010, 1% of all deaths (6% of neonatal and postneonatal infant deaths, 2.5% of deaths 1–4 years), ranking 23rd among all causes of death.¹ Since deaths due to congenital anomaly tend to be very early in life,³ the burden

What is already known on this topic?

- ▶ Global Burden of Disease (GBD) estimates of years of life lost due to congenital anomaly have been declining over time worldwide.
- ▶ GBD estimates include only live births.
- ▶ It is difficult to obtain accurate cause of death data from death certificates for infant and child deaths.

What this study adds?

- ▶ EUROCAT data (2005–2009) show terminations of pregnancy for congenital anomaly were almost three times more frequent than infant deaths and stillbirths with congenital anomaly combined.
- ▶ In countries where termination of pregnancy is illegal, the rate of infant mortality with congenital anomaly is higher than in other European countries.
- ▶ Ignoring termination of pregnancy and stillbirth underestimates the GBD associated with congenital anomaly and obscures lack of progress in primary, secondary and tertiary prevention.

in years of life lost (YLL) is higher — congenital anomaly ranked 14th among causes of death.¹

The GBD estimates were based on the WHO Mortality Database,⁴ for which death certification is the major data source. For low-income/middle-income countries, they are likely to be underestimated due to the absence of accurate cause of death data^{5,6} — Modell *et al* have calculated that mortality due to congenital anomaly for the under-5 age group is likely to be a fourfold underestimation,⁶ and have proposed that better mortality estimates would be produced by combining prevalence estimates with estimated survival in different healthcare systems. Further caveats in interpreting GBD statistics for congenital anomaly are that stillbirths and terminations of

CrossMark

To cite: Boyle B, Addor M-C, Arriola L, *et al*. *Arch Dis Child Fetal Neonatal Ed* 2018;**103**:F22–F28.

pregnancy for fetal anomaly (TOPFA) after prenatal diagnosis are not counted among deaths. The practice of civil registration of perinatal deaths due to severe or lethal congenital anomaly as stillbirths or very early neonatal deaths can vary in time and place due to differences in registration practices⁷ and in resuscitation practices at the edge⁸ of viability.^{8–10} Thus it is illogical to count deaths in the first week of life in GBD, but not stillbirths.

EUROCAT is a network of population-based congenital anomaly registries in Europe covering approximately 30% of the European Union (EU) birth population as well as some non-EU countries. Prevalence data¹¹ are based on a well-validated, harmonised system.¹² The aims of this paper are to compare infant mortality data collected by EUROCAT registries with that provided in the WHO Mortality Database, and to discuss the significance of stillbirths and TOPFA in the interpretation of infant mortality statistics. In Europe, TOPFA is legal in all countries represented in EUROCAT, except Ireland and Malta.

METHODS

The EUROCAT central database contains standard data on congenital anomaly cases from population-based congenital anomaly registries.^{12–14} Cases include live births, fetal deaths from 20 weeks' gestation (including stillbirths) and TOPFA (of any gestational age). Survival (yes/no) to 1 week (days 0–6) is recorded for all liveborn cases by all registries. Age at death (in days) is also available for 29 registries in 19 countries, at least up to age 1 year, but may not be complete as many registries do not have access to death certificates.¹³ These 29 registries include two associate registries (Finland and Sweden), which provided aggregate data regarding survival but do not contribute individual data to the central database. In the 19 countries, 3%–100% of national births are covered by EUROCAT registries (table 1).

For ascertainment of survival, 9 out of 29 registries have systematic access to death certificates (table 1); 9 have systematic access to hospital notes up to at least 1 year; 2 registries have systematic access to hospital notes up to 1 week or discharge; and 8 out of 29 have no systematic data source.

The WHO Mortality Database⁴ includes the total number of deaths under 1 year (infant deaths) by cause of death. Stillbirths are not included. Numbers of all infant deaths and infant deaths due to congenital anomaly were downloaded for the years 2005–2009, by country, for all countries participating in EUROCAT (table 1). The population for which both WHO and EUROCAT data were available (at the time of download) comprised 2715 012 live births. Infant mortality recorded by EUROCAT is slightly differently defined — deaths of registered cases are recorded, but the cause of death (which is not recorded) may occasionally not be related to the congenital anomaly, and we therefore refer to this as 'deaths of infants with congenital anomaly'.

WHO rates per 1000 births of 'infant death/mortality due to congenital anomaly' for the entire country were compared with EUROCAT rates per 1000 births of 'infant death/mortality with congenital anomaly' for the portion of the country covered by EUROCAT. WHO data are based on year of death while EUROCAT data are based on year of birth. The comparison (and all further analyses of EUROCAT data) was based on years (of death/birth) present in both databases. There were 11 EUROCAT countries (19 registries) that recorded rates of infant mortality with congenital anomaly of 80% or more than the rate recorded by WHO (figure 1), thus indicating that coverage of deaths up to 1 year was reasonably complete. Data from 17 registries in

9 of these countries (ie, excluding Finland and Sweden) were included in further analyses of pregnancy outcome and age at death.

EUROCAT records all major congenital anomalies, excluding a specified list of minor anomalies and conditions that are poorly specified or related to immaturity at birth.¹² Major congenital anomalies for this paper were classified into subgroups according to EUROCAT Guide 1.3¹² based on International Classification of Diseases (ICD) codes, further classified by isolated/multiply malformed/syndrome status according to the EUROCAT multiple malformation algorithm.¹⁵ For this paper, cases were also grouped according to chances of survival: lethal anomalies (anencephaly, bilateral renal agenesis, trisomy 13 and 18), anomalies associated with TOPFA (Down syndrome, spina bifida, analysed separately) and anomalies usually liveborn and requiring surgery^{16 17} (severe congenital heart defects, digestive system anomalies, abdominal wall defects, craniosynostosis and orofacial clefts).

Analyses included prevalence per 1000 births/live births of infant deaths with congenital anomaly, proportion of infant deaths among live births by age at death and prevalence of congenital anomaly cases from 'all diagnosed cases' including TOPFA, to all cases surviving to 20 weeks (ie, excluding TOPFA <20 weeks), to birth (live birth or stillbirth or late fetal death from 20 weeks' gestation), to live birth, to age 1 week, 28 days and 1 year.

Perinatal death/mortality is here defined as all fetal deaths from 20 weeks' gestation (including stillbirths) and first-week deaths, expressed per 1000 total births; infant mortality: deaths at age <1 year per 1000 live births; early neonatal mortality: first week deaths per 1000 live births; late neonatal mortality: deaths from 7 days to 27 days per 1000 live births; and post-neonatal mortality: deaths from 1 month to 1 year per 1000 live births.

This research is in compliance with the Helsinki Declaration (<http://www.wma.net/en/30publications/10policies/b3/index.html>). Permission for the study as part of the EUROCAT project was granted by the Institute of Nursing and Health Research, School of Nursing Research Governance Filter Committee at Ulster University on behalf of the Ulster University Ethics Committee.

RESULTS

According to the WHO data, 17%–43% of infant mortality was attributed to congenital anomaly, with higher attributable rates recorded in Malta (43%) and Ireland (42%) (table 1).

After excluding the eight EUROCAT countries with poor survival data (infant mortality with congenital anomaly more than 20% lower than recorded by WHO), there remained seven countries where EUROCAT estimates were between 80% and 100% of the WHO rate, in six of which some of the difference may be explained by geographical non-representativeness. In four countries, EUROCAT estimates were higher than WHO. The greatest differences were in Finland, where WHO mortality was 0.92 per 1000 but the EUROCAT estimate (with 100% national coverage) was 29% higher at 1.19 per 1000 (table 1), and Ireland, where WHO mortality was 1.63 but the EUROCAT estimate (with 62% national coverage) was 26% higher at 2.05 per 1000 (table 1). The average (EUROCAT) infant mortality with congenital anomaly in the 11 analysed countries was 1.1 per 1000 (figure 2).

The average total prevalence of congenital anomaly cases (including live births, fetal deaths more than 20 weeks and

Table 1 Death from congenital anomaly (CA) in the first year of life (infant deaths), comparing EUROCAT data with that reported by WHO per country for selected years 2005–2009*

WHO		EUROCAT							
Country	Years where WHO data are available	All infant deaths (n)	Deaths <1 year due to CA per 1000 live births	Infant deaths due to CA (%)	Country covered by EUROCAT (%)	Data source for infant deathst	Liveborn CA cases registered (n)	Infant deaths with CA (n)	Deaths of infants with CA per 1000 live birthst
Austria, Styria	2005–2009	1464	1.01	27	13	(B)	1323	41	0.81
Belgium, Antwerp and Hainaut	2005–2006	964	0.98	24	27	(D)(E)	1217	55	0.87
Croatia, Zagreb	2005–2009	1121	1.80	34	16	(B)	641	42	1.26
Denmark, Odense	2005–2006	506	1.09	28	8	(B)	290	11	1.06
Finland	2005–2009	825	0.92	33	100	(A)	12 890	353	1.19
France - Ile de la Reunion*	2005–2006	331	1.23	17	100	(C)	570	39	0.84
Germany, Mainz and Saxony-Anhalt	2005–2009	12 679	0.96	26	3	(A)(B)	3014	49	0.47
Hungary*	2008–2009	1048	1.38	25	100	(D)	6209	172	0.89
Ireland, Cork and Kerry, Dublin and South East Ireland	2005–2009	1245	1.63	42	62	(A)(A)(B)	4166	412	2.05
Italy, Tuscany and Emilia Romagna	2006–2008	5987	0.93	26	13	(A) (C)	4051	58	0.27
Malta	2005–2009	118	2.56	43	100	(A)	529	60	3.02
Netherlands, North	2005–2009	3893	1.26	30	10	(B)	2014	106	1.21
Poland, Wielkopolska	2005–2009	11 564	1.93	32	10	(D)	4777	138	0.73
Portugal, South	2007–2009	1065	0.81	23	21	(D)	545	7	0.11
Spain-Basque Country	2005–2009	8523	0.93	26	4	(B)	1463	84	0.80
Sweden	2005–2009	1366	0.77	30	100	(A)	8492	331	0.63
Switzerland, Vaud	2005–2007	926	1.31	31	10	(B)	650	20	0.90
Ukraine	2005–2006 2008–2009	18 542	2.59	31	6	(A)	2383	374	2.75
UK, Wales, N England, EMSYCAR, Wessex, Thames Valley, SW England	2005–2009	18 571	1.15	23	32	(A)(B)(D)(D)(D)	24 539	1141	0.95

*Data are given for both WHO and EUROCAT for the years for which WHO data are available; WHO data are national data, EUROCAT data are for the areas covered by EUROCAT registries. In Hungary WHO data were available for 2005–2009 but EUROCAT data only available 2008–2009. EUROCAT infant mortality data for France were available only for Ile de la Reunion and WHO publishes mortality data for Ile de la Reunion separately from the rest of France.

tData source for EUROCAT registries: A=systematic access to death certificates +/– other sources; B=systematic access to hospital notes up to at least 1 year but not death certificates; C=systematic access to hospital notes up to 1 week or discharge; D=no systematic data source; E=not known. Where there is more than one registry per country, the codes given are in the same order as the registries appear in column 1.

#The EUROCAT rate (deaths of infants with congenital anomaly) would be expected to be higher than the rate of deaths due to congenital anomaly (WHO) as a small number of infants who were diagnosed as having a congenital anomaly will die of other causes.

Figure 1 Prevalence per 1000 live births of infant deaths due to congenital anomaly (WHO) and infant deaths with congenital anomaly (EUROCAT) for 29 EUROCAT registries in 19 countries, 2005–2009. *Registries and years as stated in Table 1.

TOPFA) in the 11 countries was 26.9 per 1000 births. Of these 77.0% were live births surviving infancy.

A total of 17% of congenital anomaly cases were TOPFA, or a prevalence of 4.6 per 1000 births (figure 2), varying from 0 (Malta) to 7.5 per 1000 (Spain-Basque Country) (figure 2). A total of 2.1% were stillbirths, a prevalence of 0.6 per 1000 births, also varying by country (figure 2). Early neonatal deaths accounted for 49% of all first-year deaths. The prevalence of perinatal deaths and infant deaths was higher in those countries where TOPFA is illegal (figure 2). In Ireland 10.6% of all congenital anomaly cases were perinatal deaths, and in Malta 8.9%. Ukraine differed from the other countries in having relatively high perinatal and infant mortality in spite of having a TOPFA rate at the lower end of the range (figure 2). Spain-Basque Country had the highest overall mortality

(figure 2), most of this being due to TOPFA, and lowest perinatal mortality.

Overall, TOPFA was four times more common than infant deaths and nearly three times more common than all stillbirths and infant deaths combined (figure 2). Infant mortality was highest for multiply malformed live births (15% of cases) and lowest for cases with an isolated anomaly (nearly 4%, figure 3).

The survival pattern and prevalence of survivors at 1 year differs between countries where TOPFA is legal and illegal (figure 4). For lethal anomalies, the prevalence of survivors converges at 1 week of life between countries where TOPFA is legal and illegal (figure 4). For Down syndrome and spina bifida, where TOPFA is frequent but infant mortality is low, the difference between countries where TOPFA is legal and illegal is greatest and persists up to 1 year of life (figure 4). For congenital anomaly typically liveborn and requiring

Figure 2 Prevalence per 1000 births of infant deaths with congenital anomaly, by age at death and country, for 19 EUROCAT registries in 11 countries, 2005–2009. *TOPFA, terminations of pregnancy for fetal anomaly. *Years as stated in Table 1. Eleven countries (19 EUROCAT Registries) which ascertain more than 80% of the infant mortality attributed to congenital anomaly recorded by WHO: Austria = Styria; Belgium = Antwerp and Hainaut; Denmark = Odense; Finland; Ireland = Dublin, SE Ireland and Cork & Kerry; Malta; Northern Netherlands; Spain = Basque Country, Sweden; UK = Wales, SW England, Wessex, Northern England, EMSYCAR and Thames Valley; Ukraine.

Figure 3 Proportion of infant deaths among live births with congenital anomaly, by age at death, and association with multiple malformations or syndromes, for 17 EUROCAT registries in nine countries, 2005–2009. *Years as stated in Table 1. Nine countries (17 full member EUROCAT registries) which ascertain more than 80% of the infant mortality attributed to congenital anomaly recorded by WHO: Austria = Styria; Belgium = Antwerp and Hainaut; Denmark = Odense; Ireland = Dublin, SE Ireland and Cork & Kerry; Spain = Basque Country; UK = Wales, SW England, Wessex, Northern England, EMSYCAR and Thames Valley; Ukraine. *Years as stated in Table 1.

Figure 4 Prevalence of congenital anomaly cases per 1000 births, from before 20 weeks' pregnancy up to 1 year of age, by legality of TOPFA* for lethal anomalies, Down syndrome, spina bifida, and surgically treated anomalies, for 17 EUROCAT registries in nine countries,** 2005–2009.***
 *TOPFA is illegal in Malta and Ireland. **Nine countries (15 full EUROCAT registries) which ascertain more than 80% of the infant mortality attributed to congenital anomaly recorded by WHO: Austria = Styria; Belgium = Antwerp and Hainaut; Denmark = Odense; Ireland = Dublin, SE Ireland and Cork & Kerry; Malta; Northern Netherlands; Spain = Basque Country; UK = Wales, SW England, Wessex, Northern England, EMSYCAR and Thames Valley; Ukraine. ***years as stated in Table 1 [^]Lethal anomalies are anencephaly, bilateral renal agenesis and trisomy 13 and 18 ^{^^}Anomalies typically requiring surgery are: severe congenital heart defects, digestive system anomalies, abdominal wall defects, craniosynostosis and oro-facial clefts. See Public Health Indicators Khoshnood et al (8) all diagnosed cases" including TOPFA, to all cases surviving to 20 weeks (i.e. excluding TOPFA <20 weeks), to birth (livebirth or stillbirth or late fetal death from 20 weeks gestation), to livebirth, to age one week, 28 days and 1 year.

surgery, the survival pattern converges at 1 year of age in the two groups of countries (figure 4).

Among all TOPFAs, 42% were at 20 weeks' gestational age or later, varying according to anomaly category (figure 4).

Thirty-five per cent of babies (live or still) with lethal anomalies were stillborn (figure 4), as against 4%–9% for the other categories (figure 4). Infants with lethal anomalies accounted for 16% of all recorded infant mortality with a congenital anomaly and infants with surgically treatable anomalies 52%.

DISCUSSION

Infant mortality due to congenital anomaly in Europe, according to the WHO mortality data based on death certificates, ranges from one-fifth to two-fifths of infant deaths. Strategies to reduce infant mortality due to congenital anomaly are therefore an important part of meeting the millennium goals to reduce infant mortality¹⁸ and the sustainable development goals for 2030,¹⁹ and we need accurate and comprehensive data to monitor progress. We found that EUROCAT national estimates of infant mortality could be up to 29% higher than WHO estimates based on death certificates alone, assumed to be mainly cases where congenital anomaly had not been given as the primary cause of death on death certificates. On the other hand, we also found evidence of under-recording of mortality by some EUROCAT congenital anomaly registries that was unlikely to be explained by limitations of the extrapolation of regional to national data. Congenital anomaly registries should

be adequately resourced for, and data protection legislation should allow, their role in establishing fundamental population health statistics.

EUROCAT data show that TOPFA is on average four times more frequent than infant deaths, and nearly three times more frequent than stillbirths and infant deaths combined. Infant mortality due to congenital anomaly in a country is higher when the prevalence of TOPFA is lower, and it follows that increases over time in the TOPFA rate would lower the infant mortality rate. There is a danger then that apparent improvements in infant mortality due to congenital anomaly (and associated YLL) in GBD estimates could, via exclusion of TOPFA, mask lack of progress in primary prevention,²⁰ lack of improvement in tertiary prevention via surgery⁵ and/or lack of improvement in secondary prevention via early diagnosis (prenatally or postnatally), leading to more successful treatment.²¹ This may be leading to misinterpretation of GBD statistics, such as the fall in YLL between 1990 and 2010.^{1 2} In addition, the current GBD estimates omit any appropriate weighting of the disability associated with loss of wanted pregnancies.

In Europe in the last 30 years, little progress has been made towards primary prevention of congenital anomalies, and this has occasioned EUROCAT to issue recommendations to encourage integrated primary prevention policies.²² Although periconceptional folic acid is a known protective factor for neural tube defects, evidence to date shows that neural tube defect prevalence has not declined²³ due to failure of public health policy and implementation. It has only been possible to bring this to light by

including data on TOPFA in surveillance, showing that increases in TOPFA have led to decreases in live birth prevalence, but that there has been no overall change in total prevalence.²³

The exclusion of stillbirths from GBD burden of disease estimates has the greatest impact for lethal congenital anomalies. We had too few data to examine differences in civil registration (recording the birth of a baby with lethal anomaly as live or still) between countries, but this is likely¹⁰ for sociocultural and health system reasons.

For conditions such as spina bifida and Down syndrome with low postnatal mortality,²⁴ our data show an impact of TOPFA on the prevalence of survivors and thus on Disability Adjusted Life Year (DALYs)²⁵ rather than on YLLs. Investment in health and social services should reflect the greater needs in countries with lower TOPFA rates.

For surgically treatable anomalies (selected indicator conditions excluding spina bifida¹⁶), we found that the prevalence of survivors converges at 1 year, indicating that TOPFAs concerns the more severe cases with higher risk of mortality. European infant mortality for these conditions was much less than described for low-income and middle-income countries,⁵ and given the importance of improving surgical treatment in these countries, we suggest that separate monitoring of surgically treatable congenital anomaly be undertaken. Prenatal diagnosis may also be improving outcomes for these conditions,²⁶ for postnatal surgery and increasingly also fetal surgery. In Europe on average, these anomalies represent at least 52% of all infant deaths with congenital anomaly.

Higher infant mortality due to congenital anomaly should be expected in countries where TOPFA is illegal,²⁷ and socioeconomic differences within countries may occur where prenatal diagnosis or TOPFA is linked to ability to pay.^{28 29} However, there are some problems in the interpretation of data. There is evidence that, in those countries where abortion is illegal, some women are travelling abroad for TOPFA.³⁰ On the other hand, we slightly overestimated the impact of TOPFA on mortality, since some early TOPFA (of the 58% under 20 weeks' gestation) may without intervention have resulted in a spontaneous abortion by 20 weeks. From 10 weeks' gestation, 89% Down syndrome, 77% trisomy 18 and 73% of trisomy 13 are estimated to survive to 20 weeks' gestation³¹ while survival is higher for non-chromosomal conditions and from later gestational ages.

The majority (77% in our data) of children with congenital anomalies in the developed world are liveborn and survive infancy often with significant disability and need for treatment and rehabilitative services. The full burden of disease requires assessment of DALYs.²⁵ The validity of the disability weighting system for congenital anomalies³² used in GBD requires further examination.⁶ Almost all the DALYs currently attributed to congenital anomalies in GBD estimates are related to early death rather than disability.²⁵

The strengths of this paper are the broad geographical coverage and the high quality of case data recorded, including all pregnancy outcomes (except miscarriages before 20 weeks) and well-validated diagnoses. We analysed mortality as prevalence per 1000 births rather than proportion of all congenital anomaly cases, because proportions can be affected by variable ascertainment of live births diagnosed after the early neonatal period. The main limitation of interpretation is extrapolation from regional registries to a national scale for comparison with the WHO estimates, where there may be economic, social or healthcare differences between regions.²⁶ We classified a limited number of conditions as 'lethal', although this term has been criticised and its application found to be variable in the

literature,³² and we found a small number of deaths later than 1 year among 'lethal' conditions. Our case categories were based on EUROCAT Public Health Indicators,¹⁶ which do not include all congenital anomalies and make certain case classification decisions, such as analysing spina bifida on its own rather than included in surgically treatable anomalies, which could affect comparability with other studies.

CONCLUSIONS

GBD currently only considers the deaths of liveborn congenital anomaly cases and excludes TOPFA and stillbirths. This underestimates the societal health impact. TOPFA distorts both YLL and DALY estimates. Awareness of these issues is crucial in interpreting changes in GBD estimates in relation to primary, secondary or tertiary prevention.

Author affiliations

¹EUROCAT: WHO Collaborating Centre for the Surveillance of Congenital Anomalies, University of Ulster, Coleraine, UK

²School of Nursing and Midwifery, Queens University Belfast, Belfast, UK

³Division of Medical Genetics, CHUV, Lausanne, Switzerland

⁴Registro Anomalias Congénitas CAV Subdirección de Salud Pública Av Navarra, San Sebastian, Spain

⁵Children's Hospital Zagreb, School of Medicine, University of Zagreb, Zagreb, Croatia

⁶CNR Institute of Clinical Physiology, Via Moruzzi, Pisa, Italy

⁷Hungarian Congenital Abnormality Registry, National Public Health and Medical Officer Service, Budapest, Hungary

⁸Department of Genetics, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

⁹Departamento de Epidemiologia, Registo Nacional de Anomalias Congénitas Av Padre Cruz, Lisbon, Portugal

¹⁰Department of Epidemiology Public Health, East Midlands & South Yorkshire (EMSYCAR), University of Leicester, Leicester, UK

¹¹Department of Health Information and Research, Guardamangia, Malta

¹²Department of Paediatric, Hospital Lillebaelt, Kolding, Denmark

¹³Department of Obstetrics and Gynecology, Medical University of Graz, Graz, Austria

¹⁴Swedish National Board of Health and Welfare and Department of Reproduction Epidemiology, Institution of Clinical Sciences, University of Lund, Lund, Sweden

¹⁵Department of Medical Genetics, University of Medical Sciences, Poznan, Poland

¹⁶Health Service Executive, Dublin, Ireland

¹⁷Health Service Executive, Kilkenny, Ireland

¹⁸Department of Environment, PIH, Province of Antwerp, Antwerp, Belgium

¹⁹Azienda Ospedaliero-Universitaria di Ferrara Corso Giovecca, Ferrara, Italy

²⁰Health Service Executive, Cork, Ireland

²¹Birth Registry Mainz Model, Children's Hospital University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany

²²Register of Reunion Island, Centre Hospitalo-Universitaire, St Pierre La Reunion, Reunion, UK

²³Institute of Health and Society, Newcastle University, Newcastle, UK

²⁴Malformation Monitoring Centre, Saxony-Anhalt, Medical Faculty Otto-von-Guericke University, Magdeburg, Germany

²⁵National Institute for Welfare and Health (THL), Helsinki, Finland

²⁶National Perinatal Epidemiology Unit, University of Oxford, Oxford, UK

²⁷Public Health Wales, Congenital Anomaly Register and Information Service for Wales (CARIS), Swansea, UK

²⁸Centre de Génétique Humaine IPG Institut de Pathologie et de Génétique Avenue G Lemaître, Charleroi, Belgium

²⁹Faculty of Medicine, University of Southampton and Wessex Clinical Genetics Service, Southampton, UK

³⁰School of Clinical Sciences, University of Bristol, Bristol, UK

³¹Khmelnytsky Perinatal Center, OMNI-Net Ukraine Birth Defects Program, Khmelnytsky, Ukraine

Contributors HD and BB defined the research question, designed the study and interpreted the analysis. BB and HD co-wrote the paper. BB prepared and analysed the data. M-CA, LA, IB, FB, MC-S, HEKdW, CMD, ED, MG, EG, MH, KK, AL-B, BM, CM, VN, AJN, MO, AQ-W, HR, JR, AnKR, AnnR, CR, DT, CV-D, DW, BW and NZ-Z provided data and commented on drafts.

Funding Executive Agency for Health and Consumers Grant Agreement: 2013 3307 UU (OG).

Competing interests None declared.

Ethics approval Ethical approval has been granted for the collection of these data.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the global burden of disease study 2010. *Lancet* 2012;380:2095–128.
- Murray CJ, Ezzati M, Flaxman AD, et al. GBD 2010: a multi-investigator collaboration for global comparative descriptive epidemiology. *Lancet* 2012;380:2055–8.
- Liu L, Johnson HL, Cousens S, et al. Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. *Lancet* 2012;379:2151–61.
- WHO. Mortality Statistics. 2012 <http://apps.who.int/healthinfo/statistics/mortality/who.htm> (accessed 18 Aug 2013).
- Sitkin NA, Ozgediz D, Donkor P, et al. Congenital anomalies in low- and middle-income countries: the unborn child of global surgery. *World J Surg* 2015;39:36–40.
- Modell B, Berry RJ, Boyle CA, et al. Global regional and national causes of child mortality. *Lancet* 2012;380:1556.
- Pharoah PO. Anomalies occur in registrations of fetal deaths in multiple pregnancies. *BMJ* 1999;319:188.
- Singh J, Fanaroff J, Andrews B, et al. Resuscitation in the "gray zone" of viability: determining physician preferences and predicting infant outcomes. *Pediatrics* 2007;120:519–26.
- Blencowe H, Cousens S, Jassir FB, et al. Lancet Stillbirth Epidemiology Investigator Group. National, regional, and worldwide estimates of stillbirth rates in 2015, with trends from 2000: a systematic analysis. *Lancet Glob Health* 2016;4:e98–108.
- McDonnell R, Delany V, O'Mahony MT, et al. Neural tube defects in the republic of Ireland in 2009–11. *J Public Health* 2015;37:fd016.
- Boyd PA, Haesusler M, Barisic I, et al. Paper 1: the EUROCAT network--organization and processes. *Birth Defects Res A Clin Mol Teratol* 2011;91(Suppl 1):S2–5.
- EUROCAT. EUROCAT: guide 1.3 and Reference documents. 2013. <http://www.eurocat-network.eu/content/EUROCAT-Guide-1.3.pdf> (accessed 7 Apr 2014).
- Greenlees R, Neville A, Addor MC, et al. Paper 6: eurocat member registries: organization and activities. *Birth Defects Res A Clin Mol Teratol* 2011;91(Suppl 1):S51–100.
- Calzolari E, Barisic I, Loane M, et al. Epidemiology of multiple congenital anomalies in Europe: a EUROCAT population-based registry study. *Birth Defects Res A Clin Mol Teratol* 2014;100:270–6.
- Khoshnood B, Greenlees R, Loane M, et al. Paper 2: EUROCAT public health indicators for congenital anomalies in Europe. *Birth Defects Res A Clin Mol Teratol* 2011;91(Suppl 1):S16–22.
- Boyd PA, Rounding C, Chamberlain P, et al. The evolution of prenatal screening and diagnosis and its impact on an unselected population over an 18-year period. *BJOG* 2012;119:1131–40.
- Lozano R, Wang H, Foreman KJ, et al. Progress towards Millennium Development Goals 4 and 5 on maternal and child mortality: an updated systematic analysis. *Lancet* 2011;378:1139–65.
- United Nations. Sustainable Development Goals. 2015 <https://sustainabledevelopment.un.org/?menu=1300> (accessed 26 Jan 2016).
- Dolk H; EUROCAT Project Management Committee. What is the "primary" prevention of congenital anomalies? *Lancet* 2009;374:378.
- Reiss I, Schaible T, van den Hout L, et al. Standardized postnatal management of infants with congenital diaphragmatic hernia in Europe: the CDH EURO Consortium consensus. *Neonatology* 2010;98:354–64.
- Taruscio D, Arriola L, Baldi F, et al. European recommendations for primary prevention of congenital anomalies: a joined effort of EUROCAT and EUROPLAN projects to facilitate inclusion of this topic in the National Rare Disease Plans. *Public Health Genomics* 2014;17:115–23.
- Khoshnood B, Loane M, de Walle H, et al. Long term trends in prevalence of neural tube defects in Europe: population based study. *BMJ* 2015;351:h5949.
- Tennant PW, Pearce MS, Bythell M, et al. 20-year survival of children born with congenital anomalies: a population-based study. *Lancet* 2010;375:649–56.
- Murray CJ, Vos T, Lozano R, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the global burden of disease study 2010. *Lancet* 2012;380:2197–223.
- Kancherla V, Oakley GP, Brent RL. Urgent global opportunities to prevent birth defects. *Semin Fetal Neonatal Med* 2014;19:153–60.
- Gatt M, England K, Grech V, et al. Contribution of congenital anomalies to neonatal Mortality Rates in Malta. *Paediatr Perinat Epidemiol* 2015;29:401–6.
- Budd JL, Draper ES, Lotto RR, et al. Socioeconomic inequalities in pregnancy outcome associated with down syndrome: a population-based study. *Arch Dis Child Fetal Neonatal Ed* 2015;100:F400–4.
- Hutcheon JA, Bodnar LM, Simhan HN. Medicaid pregnancy termination funding and racial disparities in congenital anomaly-related infant deaths. *Obstet Gynecol* 2015;125:163–9.
- Loane M, Morris JK, Addor MC, et al. Twenty-year trends in the prevalence of down syndrome and other trisomies in Europe: impact of maternal age and prenatal screening. *Eur J Hum Genet* 2013;21:27–33.
- EUROCAT. Prevalence data. 2014 <http://www.eurocat-network.eu/accessprevalencedata/prevalencetables> (accessed 7 Apr 2014).
- Salomon JA, Wang H, Freeman MK, et al. Healthy life expectancy for 187 countries, 1990–2010: a systematic analysis for the global burden disease study 2010. *Lancet* 2012;380:2144–62.
- Wilkinson DJC, Thiele P, Watkins A, et al. Fatally flawed? A review and ethical analysis of lethal congenital malformations. *BJOG* 2012;119:1302–8.