



OPEN ACCESS

# Neonatal admission and mortality in babies born in UK alongside midwifery units: a national population-based case-control study using the UK Midwifery Study System (UKMidSS)

Rachel Rowe ,<sup>1</sup> Aung Soe,<sup>2</sup> Marian Knight ,<sup>1</sup> Jennifer J Kurinczuk,<sup>1</sup> UK Midwifery Study System (UKMidSS)

<sup>1</sup>National Perinatal Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, UK

<sup>2</sup>Oliver Fisher Neonatal Intensive Care Unit, Medway Maritime Hospital, Medway NHS Foundation Trust, Gillingham, UK

## Correspondence to

Associate Professor Rachel Rowe, National Perinatal Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford OX3 7LF, UK; rachel.rowe@npeu.ox.ac.uk

Received 3 March 2020

Revised 14 July 2020

Accepted 21 August 2020

Published Online First

30 October 2020

## ABSTRACT

**Objectives** To determine the incidence of and risk factors for neonatal unit admission, intrapartum stillbirth or neonatal death without admission, and describe outcomes, in babies born in an alongside midwifery unit (AMU).

**Design** National population-based case-control study. **Method** We used the UK Midwifery Study System to identify and collect data about 1041 women who gave birth in AMUs, March 2017 to February 2018, whose babies were admitted to a neonatal unit or died (cases) and 1984 controls from the same AMUs. We used multivariable logistic regression, generating adjusted OR (aOR) with 95% CIs, to investigate maternal and intrapartum factors associated with neonatal admission or mortality.

**Results** The incidence of neonatal admission or mortality following birth in an AMU was 1.2%, comprising neonatal admission (1.2%) and mortality (0.01%). White 'other' ethnicity (aOR=1.28; 95% CI=1.01 to 1.63); nulliparity (aOR=2.09; 95% CI=1.78 to 2.45);  $\geq 2$  previous pregnancies  $\geq 24$  weeks' gestation (aOR=1.38; 95% CI=1.10 to 1.74); male sex (aOR=1.46; 95% CI=1.23 to 1.75); maternal pregnancy problem (aOR=1.40; 95% CI=1.03 to 1.90); prolonged (aOR=1.42; 95% CI=1.01 to 2.01) or unrecorded (aOR=1.38; 95% CI=1.05 to 1.81) second stage duration; opiate use (aOR=1.31; 95% CI=1.02 to 1.68); shoulder dystocia (aOR=5.06; 95% CI=3.00 to 8.52); birth weight <2500 g (aOR=4.12; 95% CI=1.97 to 8.60), 4000–4999 g (aOR=1.64; 95% CI=1.25 to 2.14) and  $\geq 4500$  g (aOR=2.10; 95% CI=1.17 to 3.76), were independently associated with neonatal admission or mortality. Among babies admitted (n=1038), 18% received intensive care. Nine babies died, six following neonatal admission. Sepsis (52%) and respiratory distress (42%) were the most common discharge diagnoses.

**Conclusions** The results of this study are in line with other evidence on risk factors for neonatal admission, and reassuring in terms of the quality and safety of care in AMUs.

## INTRODUCTION

Most babies in high-income countries are born in hospital obstetric units (OU),<sup>1 2</sup> but in the UK around 15% of births take place in midwifery-led

## What is already known on this topic?

- ▶ National guidance recommends that women at low risk of complications should have a choice of planned birth setting.
- ▶ For low-risk women planned midwifery unit birth is as safe for babies and associated with reduced intervention, compared with planned hospital obstetric unit birth.
- ▶ Neonatal admission following birth in a midwifery unit, where most women are low risk, not induced and give birth vaginally, is a potential 'near-miss' event.

## What this study adds?

- ▶ Around 1% of babies born in the UK alongside midwifery units are admitted to neonatal care, are stillborn or die soon after birth without admission.
- ▶ Few babies had diagnoses or outcomes indicative of serious intrapartum-related adverse outcome.
- ▶ Risk factors for neonatal admission or mortality identified in our study are known risk factors for adverse neonatal outcome in term infants in other settings.

settings.<sup>3 4</sup> Around 80% of these are in alongside midwifery units (AMU), on the same site as an OU. For women at low risk of complications, planned birth in an AMU is associated with less intervention,<sup>5</sup> including a 60% reduction in the likelihood of caesarean section, with no difference in neonatal outcomes, compared with planned birth in an OU.<sup>6</sup> Most women planning birth in AMUs are at low risk, but around 4% have pre-existing risk factors<sup>6 7</sup>; many AMUs now explicitly admit women with selected risk factors.<sup>8 9</sup>

National guidance recommends transfer to an OU when complications occur during labour in a midwifery-led setting.<sup>10</sup> However, around 40% of adverse perinatal outcomes in births planned in midwifery-led settings occur in births in the planned setting, that is, when no transfer took place.<sup>6</sup> Most babies admitted to neonatal care are born in OUs.



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

**To cite:** Rowe R, Soe A, Knight M, et al. *Arch Dis Child Fetal Neonatal Ed* 2021;**106**:F194–F203.

Neonatal admission, or death without admission, following birth in an AMU, where most women are at low risk of complications, labour spontaneously and give birth vaginally, is a potential indicator of a 'near-miss' or adverse outcome event where different management might have made a difference to outcome. Factors associated with term admission and reasons for admission are well-documented,<sup>11–15</sup> but there is no evidence about admission following birth in a midwifery-led setting. This study aimed to: (i) determine the incidence of and risk factors for admission to neonatal care, intrapartum stillbirth or neonatal death without admission, in babies born in an AMU and (ii) describe reasons for neonatal admission.

## METHODS

### Study design

We carried out a national, population-based, case-control study.

### Cases and controls

We identified and collected data about women who gave birth in an AMU in the UK between 1 March 2017 and 28 February 2018, and whose baby was admitted to a neonatal unit, for at least 4 hours, within 48 hours of birth or before discharge home or who was stillborn or died within 48 hours of birth without admission to neonatal care (cases). We refer to this outcome as 'neonatal admission or mortality'. Controls were the two women not meeting the case definition who gave birth in the same AMU immediately before each case.

### Data collection

We collected data using the UK Midwifery Study System (UKMidSS), a national research infrastructure covering all 123 AMUs in all four countries of the UK. UKMidSS midwife 'reporters' in each AMU reported cases, and the number of AMU births, in response to monthly emails, and entered anonymised data directly from medical records using study-specific forms in a secure web-based environment.<sup>16</sup>

We compared the number of deaths reported for this study with those reported to national perinatal surveillance (MBRRACE-UK)<sup>17</sup> in AMU births over the same period. Where this identified potentially 'missing' deaths, not reported to UKMidSS, we contacted UKMidSS and MBRRACE-UK reporters in the relevant units to cross-check using MBRRACE-UK and UKMidSS ID numbers (no identifiers were disclosed). Where a previously unreported case was identified, data were collected as described above.


### Data

We considered maternal sociodemographic factors; pre-existing clinical characteristics and those arising during pregnancy and intrapartum and birth-related factors, as potential explanatory variables (online supplementary table S1) and collected the neonatal outcomes listed in online supplementary table S2.

### Analysis

We estimated the incidence of neonatal admission or mortality (combined and separately) using the total reported births as the denominator, with 95% CIs.

We described characteristics of cases and controls, and neonatal outcomes for cases. We used unconditional logistic regression to investigate univariable associations between explanatory variables and the primary outcome, and built a multivariable model, calculating unadjusted OR and adjusted OR (aOR) with 95% CI. Conditional logistic regression was not used since cases and



**Figure 1** Reported and confirmed cases and controls with reasons for exclusion.

controls were 'convenience matched' on the basis of time of birth only.<sup>18</sup> We used a prespecified conceptual framework approach to multivariable model building, adding variables to the model in stages from distal to proximal (maternal and pre-existing clinical characteristics first, followed by maternal clinical characteristics arising during pregnancy, maternal intrapartum factors and finally, birth-related factors). Variables were considered for inclusion in the model if  $p < 0.05$  in the univariable analysis, or if univariable analyses indicated that their association with the outcome was confounded by another variable. The contribution of each variable to the fit of the data to the model was tested for significance using the Wald test, and variables for which  $p < 0.05$  were retained in the model. We used robust variance estimation to allow for the clustering of women within units.

In a post hoc analysis, we compared the highest level of care received and reasons for admission in centres where the number of cases reported or the incidence of admission was at or above the 95th centile, with other centres, using the  $\chi^2$  test.

Our approach to handling missing data, and sample size and power calculations are presented in online supplementary box S1.

We used Stata V.15SE for all analyses.<sup>19</sup>

## RESULTS

### Response and incidence

All 123 AMUs in the UK participated (100% of eligible units), with a 99.8% response to monthly report requests.

In total, 1063 cases were reported (figure 1). There were 1041 confirmed cases, and 1984 controls, from a total of 87 102 women giving birth in AMUs. Among the 1041 cases, two women had an intrapartum stillbirth and one woman's baby died after birth without admission to neonatal care. Six of the 1038 surviving babies admitted to neonatal care subsequently died. The incidence of neonatal admission or mortality following birth in an AMU was 1.2% (95% CI 1.1 to 1.3), comprising neonatal admission (1.2%) and mortality (0.01%).

Overall, 110 out of 123 (85.4%) AMUs in the UK reported at least one confirmed case during the study period. The incidence of neonatal admission or mortality in each AMU ranged from 0% to 4.2% (median 1.0%; IQR 0.5%–1.6%).

### Univariable analysis

Among sociodemographic and pre-existing clinical characteristics of women, ethnicity, parity and previous pregnancy problems were statistically significantly associated with neonatal admission or mortality (table 1). Clinical characteristics arising during pregnancy that were statistically significantly associated

**Table 1** Sociodemographic, pre-existing and pregnancy-related clinical characteristics of women

	Cases n=1041		Controls n=1984		Unadjusted analysis		P value
	n	%	n	%	OR	(95% CI)	
<b>Maternal age (years)</b>							0.30
Under 20	43	4.1	55	2.8	1.55	(0.98 to 2.46)	
20–24	177	17.0	325	16.4	1.08	(0.85 to 1.37)	
25–29	312	30.0	619	31.2	1		
30–34	342	32.9	628	31.7	1.08	(0.89 to 1.32)	
35–39	155	14.9	324	16.3	0.95	(0.74 to 1.22)	
≥40	12	1.2	33	1.7	0.72	(0.40 to 1.30)	
Missing	0		0				
<b>Ethnic group</b>							0.004
White (UK and Ireland)	641	61.6	1210	61.0	1		
White (other)	170	16.3	246	12.4	1.30	(1.05 to 1.62)	
Asian	140	13.5	289	14.6	0.91	(0.76 to 1.10)	
Black	51	4.9	134	6.8	0.72	(0.50 to 1.03)	
Other	39	3.8	105	5.3	0.70	(0.46 to 1.06)	
Missing	0		0				
<b>Socioeconomic status</b>							0.06
Higher managerial, admin, prof	301	28.9	540	27.2	1		
Intermediate	202	19.4	362	18.3	1.00	(0.81 to 1.24)	
Routine and manual	237	22.8	499	25.2	0.85	(0.69 to 1.06)	
Unemployed/Student	100	9.6	160	8.1	1.12	(0.88 to 1.43)	
Employed, job unrecorded or uncodable	66	6.3	123	6.2	0.96	(0.70 to 1.33)	
Employment status not recorded	135	13.0	300	15.1	0.81	(0.65 to 1.00)	
<b>Area deprivation quintile*</b>							0.29
First (least deprived)	175	17.0	374	19.0	1		
Second	192	18.6	390	19.9	1.05	(0.82 to 1.35)	
Third	211	20.5	409	20.8	1.10	(0.87 to 1.40)	
Fourth	242	23.5	408	20.8	1.27	(0.98 to 1.64)	
Fifth (most deprived)	210	20.4	383	19.5	1.17	(0.92 to 1.49)	
Missing	11		20				
<b>Smoking status</b>							0.71
Non-smoker during pregnancy	903	86.9	1720	86.7	1		
Smoker during pregnancy	125	12.0	235	11.8	1.01	(0.80 to 1.28)	
Not recorded	11	1.1	29	1.5	0.72	(0.32 to 1.61)	
Missing	2		0				
<b>Previous pregnancies ≥24 weeks</b>							<0.0001
0	523	50.2	704	35.5	1.98	(1.70 to 2.31)	
1	328	31.5	875	44.1	1		
2	137	13.2	280	14.1	1.31	(1.03 to 1.65)	
3 or more	53	5.1	125	6.3	1.13	(0.77 to 1.67)	
Missing	0		0				
<b>BMI at booking (kg/m<sup>2</sup>)</b>							0.44
<18.5	23	2.2	56	2.8	0.73	(0.42 to 1.26)	
18.5–24.9	529	50.8	1063	53.6	0.88	(0.75 to 1.05)	
25–29.9	296	28.4	526	26.5	1		
30–35.0	89	8.6	154	7.8	1.03	(0.76 to 1.38)	
>35.0	26	2.5	34	1.7	1.36	(0.78 to 2.37)	
Not recorded	78	7.5	151	7.6	0.92	(0.73 to 1.16)	
Missing	0		0				
<b>Pre-existing medical risk factor†</b>							0.69
None	1026	98.6	1956	98.7	1		
One or more	15	1.4	25	1.3	1.14	(0.59 to 2.21)	
Missing	0		3				
<b>Problems in a previous pregnancy‡ (multiparous women only)</b>							0.03
None	494	95.4	1247	97.4	1		
One or more	24	4.6	33	2.6	1.83	(1.05 to 3.20)	

Continued

Table 1 Continued

	Cases n=1041		Controls n=1984		Unadjusted analysis		
	n	%	n	%	OR	(95% CI)	P value
Missing	0		0				
Current pregnancy maternal problem§							0.07
None	950	91.6	1847	93.4	1		
One or more	87	8.4	131	6.6	1.29	(0.98 to 1.71)	
Missing	4		6				
Current pregnancy fetal problem¶							0.03
None	1018	98.2	1960	99.1	1		
One or more	19	1.8	18	0.9	2.03	(1.06 to 3.88)	
Missing	4		6				
Sex of baby							<0.0001
Male	630	60.8	1009	50.9	1.50	(1.27 to 1.76)	
Female	406	39.2	973	49.1	1		
Missing	5		2				
Gestation at birth (weeks)							0.20
36–37	37	3.6	50	2.5	1.46	(0.91 to 2.35)	
38	127	12.3	213	10.8	1.18	(0.90 to 1.53)	
39	295	28.5	623	31.5	0.93	(0.78 to 1.12)	
40	362	34.9	714	36.1	1		
41–42	216	20.8	380	19.2	1.12	(0.94 to 1.34)	
Missing	4		4				

\*Area deprivation quintiles created using children in low-income families measure.<sup>37</sup>

†Essential hypertension; confirmed cardiac disease; thromboembolic disorder; atypical antibodies; hyperthyroidism; diabetes; renal disease; epilepsy.

‡Unexplained stillbirth/neonatal death or previous death related to intrapartum difficulty; previous baby with neonatal encephalopathy; primary PPH requiring treatment/transfusion; shoulder dystocia; previous caesarean section; other.

§GBS; BMI >35 kg/m<sup>2</sup>; post-term pregnancy; pre-eclampsia/pregnancy-induced hypertension; preterm prelabour rupture of membranes; substance misuse/alcohol; gestational diabetes; other.

¶Malpresentation; small for gestational age; reduced fetal movements; fetal abnormality. BMI, body mass index; GBS, group B streptococcus; PPH, postpartum haemorrhage.

with neonatal admission or mortality were: current pregnancy fetal problem and sex of the baby (table 1). Details of medical risk factors, and pregnancy problems are shown in online supplemental tables S3 and S4.

Labour-related factors significantly associated with neonatal admission or mortality were: fetal complication at the start of labour care, stage of labour at admission, prolonged second stage of labour, immersion in water, opiates for pain relief, fetal heart rate concerns or other fetal complications identified during labour and consulting an obstetrician for fetal compromise (table 2). Details of complications are shown in online supplemental tables S5–S7. All four birth-related factors: shoulder dystocia, birth weight, birth in water and mode of birth were statistically significantly associated with neonatal admission or mortality (table 2).

### Multivariable analysis: factors associated with neonatal admission or mortality

Multivariable analysis identified white ‘other’ ethnicity; nulliparity; two or more previous pregnancies  $\geq 24$  weeks’ gestation; male sex; maternal current pregnancy problem (most common group B streptococcus (GBS) and body mass index (BMI)) >35 kg/m<sup>2</sup>, see online supplemental table S2); prolonged or unrecorded duration of second stage of labour; opiates during labour; shoulder dystocia and birth weight <2500 g, 4000–4999 g and  $\geq 4500$  g, as independently associated with neonatal admission or mortality (table 3).

### Neonatal outcomes

Around three-quarters (78%) of the cases admitted to neonatal care were admitted from the birth room, rather than from the postnatal ward (table 4). Less than half (43%) received any neonatal resuscitation and in around 80% of those this comprised stimulation/positioning, inflation breaths and oxygen/ventilation breaths only. Among cases who were resuscitated, 17% were intubated and 2% received neonatal resuscitation drugs. Around one in five cases (18%) who were admitted to neonatal care received intensive care. The most common reasons for admission were respiratory problems and suspected infection, and this was reflected in diagnoses on discharge.

### Centres with high numbers of cases

Compared with other centres, in centres with more cases or higher incidence, a higher proportion of babies received special care as the highest level of neonatal care (64% vs 48%,  $p < 0.001$ ), fewer babies were admitted because of respiratory problems (60% vs 73%,  $p < 0.0001$ ) and more babies were admitted with suspected infection (52% vs 40%,  $p = 0.001$ ) (online supplemental table S8).

### DISCUSSION

Around 1% of babies born in UK AMUs during the study period were admitted to neonatal care, stillborn or died within 48 hours of birth without admission. Less than half of the babies admitted to neonatal care required resuscitation and in around 80% of those this comprised stimulation/positioning, inflation breaths

Table 2 Labour and birth-related factors

	Cases n=1041		Controls n=1984		Unadjusted analysis		
	n	%	n	%	OR	(95% CI)	P value
Maternal complications identified at start of labour care*							0.30
None	1012	97.8	1943	98.3	1		
One or more	23	2.2	34	1.7	1.30	(0.79 to 2.13)	
Missing	6		7				
Fetal complications identified at start of labour care†							<0.0001
None	960	92.8	1908	96.5			
One or more	75	7.2	69	3.5	2.16	(1.52 to 3.06)	
Missing	6		7				
Stage of labour at admission							0.03
Latent	243	23.5	430	21.7	1.07	(0.90 to 1.26)	
Active first stage	709	68.5	1340	67.6	1		
Passive second stage	27	2.6	80	4.0	0.64	(0.43 to 0.94)	
Active second stage	56	5.4	131	6.6	0.81	(0.57 to 1.14)	
Missing	6		3				
Duration of first stage of labour‡							0.21
Within guidance	881	84.6	1706	86.0	1		
Possibly prolonged	20	1.9	21	1.1	1.84	(0.92 to 3.68)	
Not recorded	140	13.5	257	12.9	1.05	(0.85 to 1.31)	
Duration of second stage of labour§							0.02
Within guidance	882	84.7	1752	88.3	1		
Possibly prolonged	70	6.7	92	4.6	1.51	(1.06 to 2.15)	
Not recorded	89	8.6	140	7.1	1.26	(0.97 to 1.65)	
Immersion in water during labour							0.002
No	539	52.1	1152	58.2	1		
Yes	496	47.9	829	41.8	1.28	(1.10 to 1.49)	
Missing	3		3				
Pethidine/Diamorphine during labour							0.002
No	887	85.7	1779	89.8	1		
Yes	148	14.3	203	10.2	1.46	(1.16 to 1.85)	
Missing	6		2				
Duration between pethidine/diamorphine and birth (hours)¶							0.29
Mean (SD)	3.8	(2.4)	3.5	(2.6)	1.05	(0.96 to 1.15)	
Median (IQR)	3.3	(2.0 to 5.4)	2.9	(1.5 to 4.7)			
Missing (n)	22		39				
Fetal heart rate concerns identified							<0.0001
No	962	93.0	1942	98.0	1		
Yes	73	7.0	39	2.0	3.78	(2.62 to 5.45)	
Missing	6		3				
Maternal complications identified during labour (before birth)**							0.78
None	1019	98.5	1948	98.3	1		
One or more	16	1.6	33	1.7	0.93	(0.54 to 1.59)	
Missing	6		3				
Fetal complications identified during labour (before birth)††							<0.0001
None	798	77.1	1860	93.9	1		
One or more	237	22.9	121	6.1	4.57	(3.61 to 5.77)	
Missing	6		3				
Obstetrician consulted for maternal compromise during labour							0.58
No	1021	98.7	1959	98.8	1		
Yes	14	1.3	23	1.2	1.17	(0.67 to 2.03)	
Missing	6		2				
Obstetrician consulted for fetal compromise during labour							<0.0001
No	968	93.5	1925	97.1	1		
Yes	67	6.5	57	2.9	2.34	(1.63 to 3.35)	
Missing	6		2				
Shoulder dystocia							<0.0001

Continued

Table 2 Continued

	Cases n=1041		Controls n=1984		Unadjusted analysis		
	n	%	n	%	OR	(95% CI)	P value
No	972	93.9	1958	98.8	1		
Yes	63	6.1	24	1.2	5.29	(3.27 to 8.55)	
Missing	6		2				
Birth weight (g)							<0.0001
<2500	25	2.4	12	0.6	4.65	(2.34 to 9.24)	
2500–2999	130	12.6	233	11.7	1.25	(0.95 to 1.63)	
3000–3499	383	37.0	855	43.2	1		
3500–3999	330	31.9	672	33.9	1.10	(0.90 to 1.33)	
4000–4499	137	13.2	183	9.2	1.67	(1.29 to 2.16)	
≥4500	31	3.0	26	1.3	2.66	(1.57 to 4.52)	
Missing	5		3				
Birth in water							0.01
No	789	76.2	1425	71.9	1		
Yes	246	23.8	557	28.1	0.80	(0.67 to 0.95)	
Missing	6		2				
Mode of birth							0.03
Spontaneous vertex	1014	98.0	1962	99.0	1		
Vaginal breech††	8	0.8	1	0.1	15.48	(1.87 to 128.32)	
Instrumental	13	1.3	18	0.9	1.40	(0.62 to 3.13)	
Missing	6		3				

\*Maternal tachycardia; hypertension; proteinuria; maternal fever; vaginal blood loss; prolonged membrane rupture; pain differing from contractions.

†Significant meconium; non-significant meconium; abnormal presentation; high/free floating head; suspected fetal growth restriction/macrosomia; suspected anhydramnios/polyhydramnios; fetal heart rate abnormality; fetal heart rate decelerations; reduced fetal movements in last 24 hours.

‡From start of active first stage to start of active second stage: within guidance ≤12 hours (nulliparous and multiparous); possibly prolonged >12 hours (nulliparous and multiparous).

§From start of active second stage to birth: within guidance ≤2 hours (nulliparous), ≤1 hour (multiparous); possibly prolonged >2 hours (nulliparous), >1 hour (multiparous).

¶In those who received pethidine/diamorphine.

\*\*Maternal tachycardia; hypertension; maternal fever; vaginal blood loss; prolonged membrane rupture; pain differing from contractions.

††Significant meconium; confirmed/suspected first stage delay; confirmed/suspected second stage delay; obstetric emergency; abnormal presentation; transverse/oblique lie; high/free floating head; fetal heart rate abnormality; fetal heart rate decelerations.

‡‡All vaginal breech births were undiagnosed before admission. One woman had a precipitate birth, shortly after admission. In the remaining eight, abnormal presentation was noted on admission or during labour and five women were seen by an obstetrician in the AMU. AMU, alongside midwifery unit.

and oxygen/ventilation breaths only. The most common reasons for admission were respiratory problems and suspected infection; this was reflected in discharge diagnoses. We found significant variation between units in rates and reasons for admission.

National statistics show that the overall incidence of neonatal admission in term babies is around 6%<sup>1</sup> and the incidence of intrapartum-related stillbirth or neonatal death at term is 0.28 per 1000 total births.<sup>20</sup> The lower incidence of these outcomes found in our study reflects the predominantly 'low risk' characteristics of the study population.

Babies of women with identified maternal pregnancy complications were 1.4 times more likely to be admitted to neonatal care. In line with other evidence,<sup>7</sup> the most common of these complications were GBS colonisation and BMI >35 kg/m<sup>2</sup>. National guidance advises that women with a BMI >35 kg/m<sup>2</sup>, and those with GBS for whom antibiotics in labour would be recommended, should be advised to plan birth in an OU rather than a midwifery-led setting, the latter group so they can receive intrapartum antibiotic prophylaxis.<sup>10 21</sup> Almost half of UK midwifery units now report admitting women who require antibiotics for GBS<sup>9</sup> and admission of women with a BMI >35 kg/m<sup>2</sup> is also widespread.<sup>8</sup> 'Higher risk' women in AMUs are more likely to be transferred before birth than women at 'low risk' of complications, but around 70% of 'higher risk' women admitted to AMUs give birth there without transfer.<sup>7</sup> Planned birth in

an AMU is associated with a significant reduction in the likelihood of having a Caesarean section compared with planned OU birth,<sup>6</sup> and there is evidence of good outcomes for women with higher BMI in AMUs.<sup>8</sup> Individual care plans are recommended for women at 'higher risk' of complications who plan to give birth outside an OU.<sup>10</sup>

Even in term appropriately grown babies, boys are more likely than girls to have lower Apgar scores, need neonatal resuscitation, develop respiratory problems and be admitted for neonatal care.<sup>22</sup> Our finding, that boys were almost 1.5 times more likely than girls to be admitted for neonatal care, in a population in which 99% of women had a spontaneous vaginal birth, suggest that these differences are not explained by an increased risk of instrumental or operative birth,<sup>22</sup> and are independent of birth weight.

Evidence about the duration of labour and neonatal outcomes is mixed,<sup>23</sup> but recent studies point to an increased risk of adverse neonatal outcomes with prolonged second stage of labour.<sup>15 24 25</sup> We found that the likelihood of neonatal admission or mortality was 1.4 times higher when the second stage of labour was 'possibly prolonged'.<sup>10</sup> Given available data it was not possible to determine why these women were not transferred to an OU, as national guidance recommends, nor whether the outcome for the baby would have been different had transfer taken place. For some women, for example, free text comments indicated

**Table 3** Factors associated with neonatal admission or mortality

	Cases n=1040		Controls n=1982		Unadjusted analysis		Adjusted analysis n=3002		P value
	n	%	n	%	OR	(95% CI)	aOR*	(95% CI)	
<b>Ethnic group</b>									0.01
White (UK and Ireland)	641	61.6	1210	61.0	1		1		
White (other)	170	16.3	246	12.4	1.30	(1.05 to 1.62)	1.28	(1.01 to 1.63)	
Asian	140	13.5	289	14.6	0.91	(0.76 to 1.10)	0.94	(0.76 to 1.16)	
Black	51	4.9	134	6.8	0.72	(0.50 to 1.03)	0.73	(0.51 to 1.04)	
Other	39	3.8	105	5.3	0.70	(0.46 to 1.06)	0.68	(0.44 to 1.05)	
<b>Previous pregnancies ≥24 weeks</b>									<0.0001
0	523	50.2	704	35.5	1.98	(1.70 to 2.31)	2.09	(1.78 to 2.45)	
1	328	31.5	875	44.1	1		1		
2 or more	190	18.3	405	20.4	1.25	(1.00 to 1.56)	1.38	(1.10 to 1.74)	
<b>Sex of baby</b>									<0.0001
Female	406	39.2	973	49.1	1		1		
Male	630	60.8	1009	50.9	1.50	(1.27 to 1.76)	1.46	(1.23 to 1.75)	
<b>Current pregnancy maternal problem†</b>									0.03
None	950	91.6	1847	93.4	1		1		
One or more	87	8.4	131	6.6	1.29	(0.98 to 1.71)	1.40	(1.03 to 1.90)	
<b>Duration of second stage of labour‡</b>									0.01
Within guidance	882	84.7	1752	88.3	1		1		
Possibly prolonged	70	6.7	92	4.6	1.51	(1.06 to 2.15)	1.42	(1.01 to 2.01)	
Not recorded	89	8.6	140	7.1	1.26	(0.97 to 1.65)	1.38	(1.05 to 1.81)	
<b>Pethidine/Diamorphine during labour</b>									0.04
No	887	85.7	1779	89.8	1		1		
Yes	148	14.3	203	10.2	1.46	(1.16 to 1.85)	1.31	(1.02 to 1.68)	
<b>Shoulder dystocia</b>									<0.0001
No	972	93.9	1958	98.8	1		1		
Yes	63	6.1	24	1.2	5.29	(3.27 to 8.55)	5.06	(3.00 to 8.52)	
<b>Birth weight (g)</b>									<0.0001
<2500	25	2.4	12	0.6	4.65	(2.34 to 9.24)	4.12	(1.97 to 8.60)	
2500–2999	130	12.6	233	11.7	1.25	(0.95 to 1.63)	1.20	(0.91 to 1.59)	
3000–3499	383	37.0	855	43.2	1		1		
3500–3999	330	31.9	672	33.9	1.10	(0.90 to 1.33)	1.10	(0.90 to 1.34)	
4000–4499	137	13.2	183	9.2	1.67	(1.29 to 2.16)	1.64	(1.25 to 2.14)	
≥4500	31	3.0	26	1.3	2.66	(1.57 to 4.52)	2.10	(1.17 to 3.76)	

\*Adjusted for all other variables in the model.

†GBS; BMI >35 kg/m<sup>2</sup>; post-term; pre-eclampsia/pregnancy-induced hypertension; preterm prelabour rupture of membranes; substance misuse/alcohol; gestational diabetes.

‡From start of active second stage to birth: within guidance ≤2 hours (nulliparous), ≤1 hour (multiparous); possibly prolonged >2 hours (nulliparous), >1 hour (multiparous).

BMI, body mass index; GBS, group B streptococcus.

that when concerns were identified late in labour an obstetrician attended the AMU to expedite birth, in preference to transfer, but available data did not permit further investigation. This might also explain at least some of the 31 instrumental births in an AMU in our study. Free text comments also indicated that, for some women, fetal concerns such as significant meconium and fetal heart decelerations were only identified in advanced labour or close to the time of birth. This might explain why these women were not transferred, as recommended by national guidance,<sup>10</sup> however the extent to which this also explained the lack of consultation with an obstetrician, seen in several cases, was not clear.

Opioids are widely used for pain relief in labour,<sup>26</sup> but are associated with neonatal respiratory depression.<sup>27–28</sup> In line with other studies,<sup>15–29</sup> we found that the likelihood of neonatal admission or mortality was 1.3 times higher in babies of women who used opioids for pain relief in labour.

Our study confirms low (<2500g) and high (>4000g) birth weight as independent risk factors for adverse neonatal outcome.<sup>29–32</sup> Fetal macrosomia is a risk factor for shoulder

dystocia.<sup>33–34</sup> Almost half of the cases with shoulder dystocia in our study had a birth weight >4000g, but shoulder dystocia was associated with a fivefold increase in the likelihood of neonatal admission after adjusting for birth weight. Shoulder dystocia is a risk factor for birth trauma injuries,<sup>35</sup> but in our study only a small proportion of the babies with documented shoulder dystocia had a discharge diagnosis of birth injury.

Finally, there are national initiatives aimed at reducing unnecessary neonatal admissions, avoiding separation of mother and baby, and standardising admission criteria,<sup>13–16</sup> and some evidence that the provision of transitional care is increasing.<sup>4</sup> Our study provides further evidence of variation in local neonatal unit admission policies,<sup>14</sup> with significant variation between units in admission rates and reasons for admission.

This was a national population-based study, which reduces the risk of bias associated with local, hospital-based studies. All eligible units participated, with over 99% response to monthly report requests and complete data returned for over 95% of reported cases, reducing the possibility of selection bias. There are nevertheless some potential limitations. We aimed to identify

Table 4 Neonatal outcomes

	Cases n=1041		Controls n=1984	
	n	%	n	%
<b>Apgar score &lt;7 at 5 min</b>				
Yes	222	21.5	10	0.5
Missing	7		2	
<b>Neonatal resuscitation</b>				
Yes	448	43.2	51	2.6
Missing	5		2	
<b>Type of resuscitation (among those who received resuscitation)</b>				
Stimulation	375	83.7	47	92.2
Positioning/Managing airway	362	80.8	27	52.9
Five inflation breaths	386	86.2	31	60.8
Oxygen	261	58.3	11	21.6
Ventilation breaths	277	61.8	10	19.6
Intubation	77	17.2	0	
Chest compression	32	7.1	1	2.0
Neonatal resuscitation drug	9	2.0	0	
<b>Hierarchy of resuscitation (among those who received resuscitation)*</b>				
Airway (A): stimulation/positioning only	15	3.4	17	33.3
Breathing 1 (B1): A or five inflation breaths	73	16.3	17	33.3
Breathing 2 (B2): A or B1 or oxygen/ventilation breaths	275	61.4	16	31.4
Breathing 3 (B3): A or B1 or B2 or intubation	51	11.4	0	
Chest compression (C): A or B or chest compression	25	5.6	1	2.0
Drugs: A or B or C or neonatal drugs	9	2.0	0	
<b>Neonatal team consulted while baby in midwifery unit</b>				
Yes	893	86.2	136	6.9
Missing	5		4	
<b>Primary reason neonatal team consulted</b>				
Respiratory problems	572	64.6	26	19.7
Suspected infection	31	3.5	16	12.1
Suspected perinatal asphyxia	89	10.1	14	10.6
Meconium aspiration	54	6.1	20	15.2
Congenital anomaly	18	2.0	12	9.1
Feeding problems	18	2.0	4	3.0
Physical trauma/birth injury	11	1.2	6	4.6
Other	93	10.5	34	25.8
Missing	7		4	
<b>Age of baby when neonatal team first consulted (hours)</b>				
Mean (SD)	4.5	(7.2)	6.9	(10.6)
Median (IQR)	0.6	(0.7 to 6.3)	2.2	(0.3 to 8.5)
<b>Skin to skin</b>				
Yes	885	86.1	1921	97.2
Missing	13		7	
<b>Initiation of breast feeding</b>				
Yes	794	77.0	1622	82.0
Missing	10		6	
<b>Age of baby at neonatal admission (hours)</b>				
Mean (SD)	n=1038		n=13†	
Mean (SD)	6.4	(8.4)	10.1	(14.4)
Median (IQR)	2.3	(0.6 to 9.5)	2.1	(2.0 to 11.5)

Continued

Table 4 Continued

	Cases n=1041		Controls n=1984	
	n	%	n	%
<b>Where was baby admitted from</b>				
Birth room	802	77.7		
Postnatal ward	230	22.3		
Missing	6			
<b>Highest level of care baby received</b>				
Intensive care	182	17.7		
High dependency care	319	31.0		
Special care	527	51.3		
Missing	10			
<b>Reasons for admission‡</b>				
Respiratory problems	731	70.4		
Suspected infection	438	42.2		
Suspected perinatal asphyxia	102	9.8		
Meconium aspiration	86	8.3		
Hypoglycaemia	57	5.5		
Congenital anomaly	46	4.4		
Feeding problems	44	4.2		
Jaundice	32	3.1		
Cardiac problems	23	2.2		
Pulse oximetry	15	1.5		
Abnormal movements	14	1.4		
Hypothermia	13	1.3		
Physical trauma/birth injury	11	1.1		
Maternal substance abuse	6	0.6		
Other§	32	3.1		
<b>Diagnoses on discharge‡</b>				
Sepsis	536	51.9		
Respiratory distress syndrome	429	41.6		
Congenital pneumonia	41	4.0		
Transient tachypnoea of the newborn	64	6.2		
Pneumothorax	18	1.7		
Hypoxic ischaemic encephalopathy	74	7.2		
Meconium aspiration syndrome	62	6.0		
PPHN	13	1.3		
Hypoglycaemia	63	6.1		
Jaundice	68	6.6		
Feeding problems	30	2.9		
Birth injury	8	0.8		
Congenital anomaly	56	5.4		
Cardiac problems	20	1.9		
Neonatal abstinence syndrome/social	4	0.4		
Normal	16	1.6		
Insufficient information/not specified	78	7.6		
Other	17	1.6		

\*Mutually exclusive hierarchy in which each category includes those babies who received that type of resuscitation, excluding those who also received any resuscitation type higher in the hierarchy.

†Thirteen babies in the control group were admitted to neonatal care for less than <4 hours. Other data for these 13 babies not shown because of small numbers.

‡More than one reason for admission/discharge diagnosis could be given.

§Other includes: shoulder dystocia, observation, skin rash, intrauterine growth restriction/low birth weight and reason not specified.



all women whose baby was admitted to neonatal care or who died following birth in an AMU. We checked against deaths reported to MBRRACE-UK, so are confident that we are unlikely to have missed deaths, but had no other sources of data against which to validate other reported cases, so it is possible that we may have missed some admissions. We were dependent on anonymised routine data from medical records so did not have data on a number of factors of interest, including, for example, staffing levels, time of day, day of the week or whether there was an agreed plan for care of the woman in an AMU in the presence of risk factors.

## CONCLUSIONS

The results of this study are broadly reassuring and in line with existing evidence about the quality and safety of care in AMUs. Relatively few babies had diagnoses of suspected asphyxia or meconium aspiration, which might be indicative of serious intrapartum-related adverse outcome. Many of the factors we identified as associated with neonatal admission or mortality are known risk factors for adverse neonatal outcome in term infants in other settings. Midwives should continue to practice in line with national guidance in relation to the management of risk factors and emerging complications in women labouring in AMUs.

**Twitter** Rachel Rowe @RachelRowe3, Marian Knight @Marianknight and UK Midwifery Study System (UKMidSS) @NPEU\_UKMidSS

**Acknowledgements** The authors would like to thank all the UKMidSS reporting midwives across the UK who responded to monthly report requests and entered data, and the MBRRACE-UK reporters who helped identify potentially 'missing' cases. The authors would also like to thank UKMidSS Research Midwives Paula Jenkins and Ginny Mounce for their contribution to managing data collection and supporting UKMidSS reporters.

**Collaborators** The UKMidSS Steering Group (in addition to RR, AS, MK and JJK): Mervi Jokinen (Chair), Royal College of Midwives; Catherine Williams (Vice Chair), Lay member; Meena Bhatia (from June 2019), Oxford University Hospitals NHS Foundation Trust; Jan Butler (until December 2018), Cambridge University Hospitals NHS Foundation Trust; Philippa Cox (from February 2019), Homerton University Hospital NHS Foundation Trust; Abi Holmes, Cardiff & Vale University Health Board; Karen Joash (from June 2019), Imperial College Healthcare NHS Trust; Eddie Morris, Royal College of Obstetricians and Gynaecologists; Jackie O' Neill (from February 2019), Northern Health and Social Care Trust; Dharmindra Pasupathy (until February 2020), Guy's and St Thomas' NHS Foundation Trust; Rachel Plachcinski, lay member; Jane Rogers, University Hospitals Southampton NHS Foundation Trust; Julia Sanders, Cardiff and Vale University Health Boards; Anna Temke (February–December 2019), South Tees Hospitals NHS Foundation Trust; Phyllis Winters (until June 2019), NHS Tayside.

**Contributors** RR conceived the study and developed the protocol with input from JJK and MK. RR managed data collection, wrote the analysis plan and carried out analysis with input from JJK, MK, AS and the UKMidSS Steering Group. RR drafted the manuscript with input from all authors. All authors were involved in interpretation of data, review and revision of the draft manuscript and approval of the final version.

**Funding** This paper presents independent research arising from a National Institute for Health Research (NIHR) Post Doctoral Fellowship awarded to Rachel Rowe (PDF-2014-07-006). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

**Competing interests** RR, MK and JJK report grants from the National Institute for Health Research (NIHR) during the conduct of this study. AS has nothing to disclose.

**Patient consent for publication** Not required.

**Ethics approval** UKMidSS and this study received ethics approval from the National Research Ethics Service (NRES) Committee South West—Frenchay (REC ref. 15/SW/0166) in May 2015.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available on reasonable request. Requests for access to the dataset underlying our findings will be considered by the National Perinatal Epidemiology Unit Data Sharing Committee and should be addressed to the Director, Jennifer Kurinczuk, jenny.kurinczuk@npeu.ox.ac.uk in the first instance.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

## ORCID iDs

Rachel Rowe <http://orcid.org/0000-0003-2994-3240>

Marian Knight <http://orcid.org/0000-0002-1984-4575>

## REFERENCES

- 1 NMPA Project Team. National maternity and perinatal audit: clinical report 2019. Based on births in NHS maternity services between 1 April 2016 and 31 March 2017. London RCOG; 2019.
- 2 Shaw D, Guise J-M, Shah N, *et al*. Drivers of maternity care in high-income countries: can health systems support woman-centred care? *Lancet* 2016;388:2282–95.
- 3 Walsh D, Spiby H, Grigg CP, *et al*. Mapping midwifery and obstetric units in England. *Midwifery* 2018;56:9–16.
- 4 Blotkamp A. National maternity and perinatal audit: Organisational report 2019. London NMPA Project Team, RCOG; 2019.
- 5 Scarf VL, Rossiter C, Vedam S, *et al*. Maternal and perinatal outcomes by planned place of birth among women with low-risk pregnancies in high-income countries: a systematic review and meta-analysis. *Midwifery* 2018;62:240–55.
- 6 Hollowell J, Puddicombe D, Rowe R. The birthplace national prospective cohort study: perinatal and maternal outcomes by planned place of birth. Birthplace in England research programme. Final report part 4. London NIHR Service Delivery and Organisation programme; 2011.
- 7 Hollowell J, Rowe R, Townend J, *et al*. The birthplace in England national prospective cohort study: further analyses to enhance policy and service delivery decision-making for planned place of birth. *Health Services and Delivery Research* 2015;3:1–264.
- 8 Rowe R, Knight M, Kurinczuk JJ, *et al*. Outcomes for women with BMI>35kg/m<sup>2</sup> admitted for labour care to alongside midwifery units in the UK: A national prospective cohort study using the UK Midwifery Study System (UKMidSS). *PLoS One* 2018;13:e0208041.
- 9 Glenister C, Burns E, Rowe R. Local guidelines for admission to UK midwifery units compared with national guidance: a national survey using the UK midwifery study system (UKMidSS). *PLoS One* 2020;15:e0239311.
- 10 National Institute for Health and Care Excellence. Intrapartum care: care of healthy women and their babies during childbirth. NICE guideline CG190. London NICE; 2014.
- 11 Tracy SK, Tracy MB, Sullivan E. Admission of term infants to neonatal intensive care: a population-based study. *Birth* 2007;34:301–7.
- 12 Alkhaat A, Hutchinson M, Jacques A, *et al*. Evaluation of the frequency and obstetric risk factors associated with term neonatal admissions to special care units. *Aust N Z J Obstet Gynaecol* 2013;53:277–82.
- 13 NHS Improvement. Reducing harm leading to avoidable admission of full-term babies into neonatal units. findings and resources for improvement, 2017. Available: [https://improvement.nhs.uk/documents/764/Reducing\\_term\\_admissions\\_final.pdf](https://improvement.nhs.uk/documents/764/Reducing_term_admissions_final.pdf)
- 14 Battersby C, Michaelides S, Upton M, *et al*. Term admissions to neonatal units in England: a role for transitional care? A retrospective cohort study. *BMJ Open* 2017;7:e016050.
- 15 Burgess APH, Katz J, Pessolano J, *et al*. Determination of antepartum and intrapartum risk factors associated with neonatal intensive care unit admission. *J Perinat Med* 2016;44:589–96.
- 16 Rowe RE, Kurinczuk JJ, Hollowell J, *et al*. The UK midwifery study system (UKMidSS): a programme of work to establish a research infrastructure to carry out national studies of uncommon conditions and events in midwifery units. *BMC Pregnancy Childbirth* 2016;16:1–6.
- 17 National Perinatal Epidemiology Unit. MBRRACE-UK: mothers and babies: reducing risk through audits and Confidential enquiries across the UK, 2019. Available: <https://www.npeu.ox.ac.uk/mbrance-uk> [Accessed 14 Jan 2020].
- 18 Bland JM, Altman DG. Matching. *BMJ* 1994;309:1128.
- 19 StataCorp. *Stata statistical software: release 15*. College Station, TX: StataCorp LLC, 2017.
- 20 Draper ES, Kurinczuk JJ, Kenyon S. MBRRACE-UK 2017 perinatal Confidential enquiry: term, singleton, intrapartum stillbirth and intrapartum-related neonatal death.

- Leicester: The Infant Mortality and Morbidity Studies, Department of Health Sciences, University of Leicester, 2017.
- 21 Hughes RG, Brocklehurst P, Steer PJ, *et al.* Prevention of early-onset neonatal group B streptococcal disease: Green-top guideline No. 36. *BJOG* 2017;124:e280–305.
  - 22 Dunn L, Prior T, Greer R, *et al.* Gender specific intrapartum and neonatal outcomes for term babies. *Eur J Obstet Gynecol Reprod Biol* 2015;185:19–22.
  - 23 Altman MR, Lydon-Rochelle MT. Prolonged second stage of labor and risk of adverse maternal and perinatal outcomes: a systematic review. *Birth* 2006;33:315–22.
  - 24 Altman M, Sandström A, Petersson G, *et al.* Prolonged second stage of labor is associated with low Apgar score. *Eur J Epidemiol* 2015;30:1209–15.
  - 25 Sandström A, Altman M, Cnattingius S, *et al.* Durations of second stage of labor and pushing, and adverse neonatal outcomes: a population-based cohort study. *J Perinatol* 2017;37:236–42.
  - 26 Bricker L, Lavender T. Parenteral opioids for labor pain relief: a systematic review. *Am J Obstet Gynecol* 2002;186:S94–109.
  - 27 Wong CA. Advances in labor analgesia. *International journal of women's health* 2010;1:139–54.
  - 28 Jones L, Othman M, Dowswell T, *et al.* Pain management for women in labour: an overview of systematic reviews. *Cochrane Database Syst Rev* 2012;3:CD009234.
  - 29 Yi Wen P, Broom E, Flatley C, *et al.* Maternal demographic and intrapartum antecedents of severe neonatal outcomes at term. *J Matern Fetal Neonatal Med* 2020;33:1–6.
  - 30 Malin GL, Morris RK, Riley R, *et al.* When is birthweight at term abnormally low? A systematic review and meta-analysis of the association and predictive ability of current birthweight standards for neonatal outcomes. *BJOG* 2014;121:515–26.
  - 31 Temerinac D, Chen X, Sütterlin M, *et al.* Influence of fetal birth weight on perinatal outcome in planned vaginal births. *Arch Gynecol Obstet* 2014;289:313–8.
  - 32 Rossi AC, Mullin P, Prefumo F. Prevention, management, and outcomes of macrosomia: a systematic review of literature and meta-analysis. *Obstet Gynecol Surv* 2013;68:702–9.
  - 33 Bahar AM. Risk factors and fetal outcome in cases of shoulder dystocia compared with normal deliveries of a similar birthweight. *Br J Obstet Gynaecol* 1996;103:868–72.
  - 34 Ouzounian JG, Gherman RB. Shoulder dystocia: are historic risk factors reliable predictors? *Am J Obstet Gynecol* 2005;192:1933–5.
  - 35 Gherman RB, Chauhan S, Ouzounian JG, *et al.* Shoulder dystocia: the unpreventable obstetric emergency with empiric management guidelines. *Am J Obstet Gynecol* 2006;195:657–72.
  - 36 Scottish Government. The best start: a five-year forward plan for maternity and neonatal care in Scotland, 2017. Available: <https://www.gov.scot/publications/best-start-five-year-forward-plan-maternity-neonatal-care-scotland-9781786527646/>
  - 37 HM Revenue & Customs. Personal tax credits: children in low-income families local measure: 2014 snapshot as at 31 August 2014: 30 September 2016, 2016. Available: <https://www.gov.uk/government/statistics/personal-tax-credits-children-in-low-income-families-local-measure-2014-snapshot-as-at-31-august-2014-30-september-2016> [Accessed 14 Jan 2020].

## SUPPLEMENTARY INFORMATION

Table S1: Potential explanatory variables and categorisation

Type of characteristic Variable (unit or how derived)	Categorisation
<b>Maternal social demographic</b>	
Maternal age (years)	Under 20; 20-24;25-29; 30-34; 35-39; ≥40
Ethnic group	White (UK & Ireland); White (other); Asian; Black; Other
Socioeconomic status (derived from woman's occupation, or partner's if woman not in work or with unrecorded/uncodable occupation, using three class NS-SEC)	Higher managerial, admin, prof; Intermediate; Routine and manual; Unemployed/student; Employed, job unrecorded or uncodable; Employment status not recorded
Area deprivation quintile (derived from women's postcode entered by midwives in to postcode look-up website which returned score from Children in Low-income Families Measure, converted into quintiles)	1st (least deprived); 2 <sup>nd</sup> ; 3 <sup>rd</sup> ; 4 <sup>th</sup> ; 5th (most deprived)
Smoking status	Non-smoker during pregnancy; Smoker during pregnancy; Not recorded
<b>Pre-existing clinical factors</b>	
Previous pregnancies ≥24 weeks' gestation	0; 1; 2; ≥3
Body mass index (BMI) at booking (kg/m <sup>2</sup> )	<18.5; 18.5-24.9; 25-29.9; 30-35.0; >35.0; Not recorded
Pre-existing medical risk factors (see Table S3)	None; One or more
Problems in a previous pregnancy (see Table S3)	None; One or more
<b>Clinical factors arising during pregnancy</b>	
Maternal current pregnancy problem (see Table S4)	None; One or more
Fetal current pregnancy problem (see Table S4)	None; One or more
Sex of baby	Male; Female
Gestation at birth (weeks)	36-37; 38; 39; 40; 41-42
<b>Intrapartum-related</b>	
Maternal complications identified at start of labour care (see Table S5)	None; One or more
Fetal complications identified at start of labour care (see Table S5)	None; One or more
Stage of labour at admission	Latent; Active 1st stage; Passive 2nd stage; Active 2nd stage
Duration of 1 <sup>st</sup> stage of labour <sup>1</sup>	Within guidance; Possibly prolonged; Not recorded <sup>1</sup>
Duration of 2 <sup>nd</sup> stage of labour <sup>2</sup>	Within guidance; Possibly prolonged; Not recorded <sup>2</sup>
Immersion in water during labour	No; Yes
Pethidine/diamorphine during labour	No; Yes
Duration between pethidine/diamorphine and birth (hours)	Mean (SD); Median (IQR)
Fetal heart rate concerns identified	No; Yes

Type of characteristic Variable (unit or how derived)	Categorisation
Maternal complications identified during labour, before birth (see Table S6)	None; One or more
Fetal complications identified during labour, before birth (see Table S6)	None; One or more
Obstetrician consulted for compromise during labour (see Table S6)	No; Yes
<b>Birth-related</b>	
Shoulder dystocia	No; Yes
Birthweight (g)	<2500; 2500-2999; 3000-3499; 3500-3999; 4000-4499; ≥4500
Birth in water	No; Yes
Mode of birth	Spontaneous vertex; Vaginal breech; Instrumental

<sup>1</sup> From start of active 1<sup>st</sup> stage to start of active 2<sup>nd</sup> stage: Categorised using NICE CG190 as: Within guidance ≤12 hours (nulliparous and multiparous); Possibly prolonged >12 hours (nulliparous and multiparous)

<sup>2</sup> From start of active 2<sup>nd</sup> stage to birth: Categorised using NICE CG190 as: Within guidance ≤2 hours (nulliparous), ≤1 hour (multiparous); Possibly prolonged >2 hours (nulliparous), >1 hour (multiparous)

Table S2: Neonatal outcomes and categorisation

Neonatal outcome (unit or how derived)	Categorisation
Apgar score at 5 minutes	≥7; <7
Neonatal resuscitation	No; Yes
Type of resuscitation	Stimulation; Positioning/managing airway; Five inflation breaths; Oxygen; Ventilation breaths; Intubation; Chest compression; Neonatal resuscitation drug
Neonatal team consulted while baby in midwifery unit	No; Yes
Primary reason for neonatal consultation	Respiratory problems; Suspected infection; Suspected perinatal asphyxia; Meconium aspiration; Congenital anomaly; Feeding problems; Physical trauma/birth injury; Other
Age of baby when neonatal team first consulted (hours)	Mean (SD); Median (IQR)
Skin to skin care	No; Yes
Initiation of breastfeeding	No; Yes
Age of baby at neonatal admission (hours)	Mean (SD); Median (IQR)
Where was baby admitted from	Birth room; Postnatal ward
Highest level of care baby received	Intensive care; High dependency care; Special care
Reasons for admission	Respiratory problems; Suspected infection; Suspected perinatal asphyxia; Meconium aspiration; Hypoglycaemia; Congenital anomaly; Feeding problems; Jaundice; Cardiac problems; Pulse oximetry; Abnormal movements; Hypothermia; Physical trauma/birth injury; Maternal substance abuse; Other
Diagnoses on discharge	Sepsis; Respiratory Distress Syndrome; Congenital Pneumonia; Transient Tachypnoea of the Newborn; Pneumothorax; Hypoxic Ischaemic Encephalopathy; Meconium Aspiration Syndrome; PPHN; Hypoglycaemia; Jaundice; Feeding problems; Birth injury; Congenital anomaly; Cardiac problems; Neonatal Abstinence Syndrome/Social; Normal; Insufficient information/not specified; Other

**Approach for handling missing data**

We anticipated that some of the data required to generate body mass index would be unrecorded in women's notes, and that this 'missing' data would not be randomly distributed, so we provided the option at data entry of indicating that height or weight were 'not recorded' and used this category in our analysis.

The only other variable with a substantial proportion of missing data was socioeconomic status. This occurred because women's or their partner's occupation is often not recorded in women's notes, and we know from previous work (e.g. Lindquist A, Knight M, Kurinczuk JJ. Variation in severe maternal morbidity according to socioeconomic position: a UK national case-control study. *BMJ Open* 2013;3:e002742 doi: 10.1136/bmjopen-2013-002742) that these data are not missing at random. Approaches such as multiple imputation would not therefore be appropriate. We therefore included a separate 'missing/not recorded' category in our analyses for these variables.

**Sample size and power**

We anticipated identifying approximately 900 cases over one year, based on an estimated incidence of 1.5% in a population of around 60,000 women (Hollowell J, Puddicombe D, Rowe R, et al. The Birthplace national prospective cohort study: perinatal and maternal outcomes by planned place of birth. Birthplace in England research programme. Final report part 4. London: NIHR Service Delivery and Organisation programme; 2011). The actual number of cases and controls generated 80% power at the 5% level to detect ORs of 1.6 or greater for an exposure variable with a frequency of 5%, and 1.3 or greater for an exposure variable with a frequency of 30%.

*Box S1: Additional methodological detail*

*Table S3: Pre-existing medical and previous pregnancy risk factors*

	Cases (n=1041)		Controls (n=1981)	
	n	%	n	%
<b>Medical risk factors</b>				
Essential hypertension	3	0.3	2	0.1
Confirmed cardiac disease	2	0.2	1	0.1
Thromboembolic disorder	0		5	0.3
Atypical antibodies	5	0.5	4	0.2
Hyperthyroidism	3	0.3	5	0.3
Diabetes	2	0.2	3	0.2
Renal disease	0		3	0.2
Epilepsy	0		2	0.1
<b>Problems in previous pregnancy</b>				
			<b>Multiparous women only</b>	
			<b>n=518</b>	<b>n=1280</b>
Previous unexplained/intrapartum-related stillbirth/death	0		3	0.2
Previous baby with Neonatal Encephalopathy	1	0.2	0	0.0
Previous primary postpartum haemorrhage requiring treatment	11	2.1	19	1.5
Previous shoulder dystocia	5	1.0	3	0.2
Previous Caesarean section	5	1.0	5	0.4
"Other" previous pregnancy problem	2	0.4	3	0.2

Table S4: Current pregnancy problems

	Cases (n=1037)		Controls (n=1978)	
	n	%	n	%
<b>Maternal</b>				
Group B Streptococcus	47	4.5	79	4.0
BMI>35kg/m <sup>2</sup>	26	2.5	34	1.7
Post-term (>42 weeks)	4	0.4	6	0.3
Pre-eclampsia/pregnancy induced hypertension	3	0.3	1	0.1
Preterm prelabour membrane rupture	0		1	0.1
Substance misuse/alcohol dependency	4	0.4	7	0.4
Gestational diabetes	4	0.4	8	0.4
Other	1	0.1	1	0.1
<b>Fetal</b>				
Multiple birth	0		0	
Malpresentation	2	0.2	1	0.1
Small for gestational age	7	0.7	9	0.5
Reduced fetal movements	4	0.4	9	0.5
Fetal abnormality	6	0.6	0	

Table S5: Complications identified at start of labour care

Complications identified at start of labour care	Cases (n=1035 <sup>1</sup> )		Controls (n=1977 <sup>1</sup> )	
	n	%	n	%
<b>Maternal</b>				
Maternal tachycardia	9	0.9	7	0.4
Essential hypertension	2	0.2	4	0.2
Proteinuria	2	0.2	4	0.2
Maternal pyrexia	0		1	0.1
Vaginal blood loss	3	0.3	1	0.1
Prolonged membrane rupture	11	1.1	19	1.0
Pain differing from contractions	1	0.1	1	0.1
<b>Fetal</b>				
Significant meconium	14	1.4	10	0.5
Non-significant meconium	22	2.1	27	1.4
Abnormal presentation	3	0.3	0	
High /free floating head	1	0.1	0	
Suspected fetal growth restriction / macrosomia	5	0.5	4	0.2
Suspected anhydramnios / polyhydramnios	2	0.2	0	
Fetal heart rate abnormality	6	0.6	5	0.3
Fetal heart rate decelerations	9	0.9	6	0.3
Reduced fetal movements in last 24 hours	18	1.7	18	0.9

<sup>1</sup> Those with non-missing data for complications identified at start of labour care

Table S6: Complications identified during labour (before birth)

Complications identified during labour (before birth)	Cases (n=1035)		Controls (n=1981)	
	n	%	n	%
<b>Maternal</b>				
Maternal tachycardia	6	0.6	9	0.5
Hypertension	4	0.4	4	0.2
Maternal pyrexia	5	0.5	3	0.2
Vaginal blood loss	7	0.7	10	0.5
Prolonged membrane rupture	7	0.7	11	0.6
Pain differing from contractions	1	0.1	0	
<b>Fetal</b>				
Significant meconium	120	11.6	49	2.5
Confirmed/suspected 1 <sup>st</sup> stage delay	14	1.4	11	0.6
Confirmed/suspected 2 <sup>nd</sup> stage delay	12	1.2	10	0.5
Obstetric emergency	15	1.5	2	0.1
Abnormal presentation	7	0.7	2	0.1
High /free floating head	2	0.2	0	
Fetal heart rate abnormality	22	2.1	14	0.7
Fetal heart rate decelerations	74	7.2	48	2.4

Note: more than one complication could be identified in each woman

Table S7: Reasons for consultation with obstetrician during labour (before birth)

Primary reason for consulting obstetrician during labour	Cases (n=82 <sup>1</sup> )		Controls (n=80 <sup>1</sup> )	
	n	%	n	%
<b>Maternal</b>				
Maternal tachycardia	1	1.2	1	1.3
Hypertension	0		2	2.5
Maternal pyrexia	0		2	2.5
Vaginal blood loss	1	1.2	0	
Prolonged membrane rupture	1	1.2	4	5.0
Pain differing from contractions	0		1	1.3
Other maternal <sup>2</sup>	11	13.4	15	18.8
<b>Fetal</b>				
Significant meconium	5	6.1	5	6.3
Confirmed/suspected 1 <sup>st</sup> stage delay	6	7.3	9	11.3
Confirmed/suspected 2 <sup>nd</sup> stage delay	4	4.9	11	13.8
Obstetric emergency	14	17.1	1	1.3
Abnormal presentation	7	8.5	1	1.3
Fetal heart rate abnormality	13	15.9	8	10.0
Fetal heart rate decelerations	18	22.0	23	28.8
Reduced fetal movements	4	4.9	2	2.5
Other fetal <sup>2</sup>	4	4.9	3	3.8

<sup>1</sup> Those for whom an obstetrician was consulted during labour and a reason was given

<sup>2</sup> Other, including advising on suitability for AMU, categorised as maternal or fetal on individual basis

Note: More than one reason could be given for each woman



Table S8: Level of care and reasons for admission in centres with high and lower number of cases

	Centres with fewer cases (n=810)		Centres with more cases (n=228)		p value
	n	%	n	%	
<b>Highest level of neonatal care</b>					<0.0001
Intensive care	146	18.3	36	15.8	
High dependency care	272	34.0	47	20.6	
Special care	382	47.8	145	63.6	
Missing	10		0		
<b>Reasons for admission</b>					
Respiratory problems	594	73.3	137	60.1	<0.0001
Suspected perinatal asphyxia	81	10.0	21	9.2	0.72
Hypoglycaemia	40	4.9	17	7.5	0.14
Physical trauma/birth injury	9	1.1	2	0.9	0.76
Feeding problems	30	3.7	14	6.1	0.11
Suspected infection	320	39.5	118	51.8	0.001
Meconium aspiration	61	7.5	25	11.0	0.10
Jaundice	21	2.6	11	4.8	0.09
Congenital anomaly	34	4.2	12	5.3	0.49
Maternal substance abuse	6	0.7	0		0.19
Cardiac problems	18	2.2	5	2.2	1.0
Pulse oximetry	11	1.4	4	1.8	0.66
Abnormal movements	11	1.4	3	1.3	0.96
Hypothermia	11	1.4	2	0.9	0.56
Other	26	3.2	6	2.6	0.65