

Subject index

- 11q23/MLL**, Congenital leukaemia after heavy abuse of permethrin during pregnancy, **F436**
- ABO incompatibility**, Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn, **F6**
- acyclovir**, Relapse of neonatal herpes simplex virus infection, **F483**
- adolescence**, Health and school performance of teenagers born before 29 weeks gestation, **F190**
- advanced neonatal nurse practitioner**, Neonatal transfers by advanced neonatal nurse practitioners and paediatric registrars, **F509**
- amiodarone**, Amiodarone and breast feeding, **F255**
- anaemia**, Late anaemia in Rh haemolytic disease, **F444**
Massive acute haemolysis in neonates with glucose-6-phosphate dehydrogenase deficiency, **F534**
- anaemia of prematurity**, Effects of vitamin E supplementation during erythropoietin treatment of the anaemia of prematurity, **F324**
- anatomy**, Galen (AD 129–200) of Pergamon: anatomist and experimental physiologist, **F441**
- antibiotics**, Neonatal sepsis in Peshawar:, **F160**
Reducing antibiotic use on the neonatal unit by improving communication of blood culture results: a completed audit cycle, **F255**
- antioxidant**, Plasma bilirubin level and oxidative stress in preterm infants, **F119**
- antioxidant cocktail**, Progress in treatment and outcome for children with neonatal haemochromatosis, **F124**
- apoptosis**, Decreased neutrophil apoptosis in tracheal fluids of preterm infants at risk of chronic lung disease, **F245**
- ascites**, Dehydrated hereditary stomatocytosis with transient perinatal ascites, **F438**
- asphyxia**, Effects of gestational age and prenatal and perinatal events on the coagulation status in premature infants, **F319**
Respiratory responses to hypoxia/hypercapnia in small for gestational age infants influenced by maternal smoking, **F217**
S100 protein content of umbilical cord blood in healthy newborns in relation to mode of delivery, **F67**
- assessment**, Neonatal murmurs: are senior house officers good enough?, **F147**
- autoimmune haemolytic anaemia**, Autoimmune haemolytic anaemia in a newborn infant, **F341**
- aviation medicine**, Excessive exposure of sick neonates to sound during transport, **F513**
- avidity**, Effect of a fourth *Haemophilus influenzae* type b immunisation in preterm infants who received dexamethasone for chronic lung disease, **F58**
- bag valve mask**, Changes in resuscitation practice at birth, **F375**
- balloon atrial septostomy**, Bedside balloon atrial septostomy on neonatal units, **F339**
- basal ganglia**, Neonatal lenticulostriate vasculopathy: further characterisation, **F410**
- BiliCheck**, Transcutaneous bilirubin measurement in newborn infants: evaluation of a new spectrophotometric method, **F350**
- bilirubin**, Phototherapy with turquoise versus blue light, **F430**
Plasma bilirubin level and oxidative stress in preterm infants, **F119**
- bioethics**, Consent for clinical research in the neonatal intensive care unit: a retrospective survey and a prospective study, **F280**
- birth**, Erich Bracht (1882–1969) of Berlin and his “breech” manoeuvre, **F76**
- birth weight**, Social trends in singleton births and birth weight in Wirral residents, 1990–2001, **F421**
Updated gestational age specific birth weight, crown-heel length, and head circumference of Chinese newborns, **F229**
- blood transfusion**, Blood transfusion in preterm neonates, **F78**
Decreasing oxygen saturation in very early preterm newborn infants after transfusion, **F163**
- body composition**, Growth and body composition in preterm infants with bronchopulmonary dysplasia, **F46**
- brain**, Cerebral perfusion in infants and neonates: preliminary results obtained using dynamic susceptibility contrast enhanced magnetic resonance imaging, **F525**
Low field strength magnetic resonance imaging of the neonatal brain, **F203**
Neonatal cerebral infarction and visual function at school age, **F487**
Progressive ventricular dilatation (PVD) over the past 22 years, **F257**
Serum S-100 protein does not correlate with cerebral ultrasound scans in preterm infants, **F160**
- brain damage**, S100 protein content of umbilical cord blood in healthy newborns in relation to mode of delivery, **F67**
- brain lesions**, Neonatal shaken baby syndrome—historical inexactitudes, **F161**
Neonatal shaken baby syndrome—lessons to be learned, **F161**
- breast feeding**, Amiodarone and breast feeding, **F255**
Effect of caesarean section on breast milk transfer to the normal term newborn over the first week of life, **F380**
Fever in healthy asymptomatic newborns during the first days of life, **F312**
Neonatal weight loss in breast and formula fed infants, **F472**
- breast milk**, Donor human milk versus formula for preventing necrotising enterocolitis in preterm infants: systematic review, **F11**
Factors associated with weaning in full term and preterm infants, **F296**
- breast milk transfer**, Effect of caesarean section on breast milk transfer to the normal term newborn over the first week of life, **F380**
- breech manoeuvre**, Erich Bracht (1882–1969) of Berlin and his “breech” manoeuvre, **F76**
- bronchopulmonary dysplasia**, Growth and body composition in preterm infants with bronchopulmonary dysplasia, **F46**
- caesarean section**, Effect of caesarean section on breast milk transfer to the normal term newborn over the first week of life, **F380**
Fever in healthy asymptomatic newborns during the first days of life, **F312**
- calcifications**, Subcutaneous fat necrosis of the newborn: hypercalcaemia with hepatic and atrial myocardial calcification, **F343**
- Candida**, Methaemoglobinaemia with concurrent blood isolation of *Saccharomyces* and *Candida*, **F446**
- cardiac**, Neonatal murmurs: are senior house officers good enough?, **F147**
- cardiac abnormalities**, Evaluation of echocardiography on the neonatal unit, **F287**
- cardiac output**, Cardiac output, pulmonary artery pressure, and patent ductus arteriosus during therapeutic cooling after global hypoxia-ischaemia, **F223**
- cardiac tamponade**, Neonatal long lines, **F260**
Pericardial effusion and cardiac tamponade as complications of neonatal long lines: are they really a problem?, **F292**
- cardiopulmonary insufficiency**, Pilot investigation of hypothermia in neonates receiving extracorporeal membrane oxygenation, **F128**
- cardiovascular system**, Andreas Vesalius (1514–1564), Padua, and the fetal “shunts”, **F157**
- carrier detection**, Implications of carrier identification in newborn screening for cystic fibrosis, **F467**
- carriers**, Cystic fibrosis newborn screening and detection of carriers, **F448**
- catheterisation**, Neonatal long lines, **F260**
- central venous**, Neonatal long lines, **F260**
- central venous catheters**, Technique for insertion of percutaneous central venous catheters in the newborn period, **F256**
- central venous line**, Unusual complication of a central venous line in a neonate, **F440**
- cerebral blood flow**, Cerebral perfusion in infants and neonates: preliminary results obtained using dynamic susceptibility contrast enhanced magnetic resonance imaging, **F525**
The light still shines, but not that brightly? The current status of perinatal near infrared spectroscopy, **F263**
- cerebral blood volume**, The light still shines, but not that brightly? The current status of perinatal near infrared spectroscopy, **F263**
- cerebral injury**, Pilot investigation of hypothermia in neonates receiving extracorporeal membrane oxygenation, **F128**

cerebral venous saturation, The light still shines, but not that brightly? The current status of perinatal near infrared spectroscopy, **F263**

chest radiograph, Frequent wheeze at follow up of very preterm infants: which factors are predictive?, **F329**

chest tubes, Congenital idiopathic chylothorax in neonates: chemical pleurodesis with povidone-iodine (Betadine), **F531**

child health, Use of healthcare resources, family function, and socioeconomic support during the first four years after preterm birth, **F415**

childhood leukaemia, Absence of leukaemic fusion gene transcripts in preterm infants exposed to diagnostic x rays, **F237**

Chinese, Updated gestational age specific birth weight, crown-heel length, and head circumference of Chinese newborns, **F229**

cholestasis, Intestinal absorption of mixed micellar phylloquinone (vitamin K₁) is unreliable in infants with conjugated hyperbilirubinaemia: implications for oral prophylaxis of vitamin K deficiency bleeding, **F113**

chronic lung disease, Cognitive development at 5.5 years of children with chronic lung disease of prematurity, **F101**
Decreased neutrophil apoptosis in tracheal fluids of preterm infants at risk of chronic lung disease, **F245**
Effect of a fourth *Haemophilus influenzae* type b immunisation in preterm infants who received dexamethasone for chronic lung disease, **F58**
Frequent wheeze at follow up of very preterm infants: which factors are predictive?, **F329**
Neonatologists are using much less dexamethasone, **F432**

chylothorax, Congenital idiopathic chylothorax in neonates: chemical pleurodesis with povidone-iodine (Betadine), **F531**

coagulation factors, Effects of gestational age and prenatal and perinatal events on the coagulation status in premature infants, **F319**

cocaine, Prenatal drug exposure and maternal and infant feeding behaviour, **F391**

cochlear damage, Noise levels within the ear and post-nasal space in neonates in intensive care, **F315**

cognition, Cognitive development at 5.5 years of children with chronic lung disease of prematurity, **F101**

colour coded duplex ultrasound, Brachial artery pseudoaneurysm in a premature infant, **F152**

communication, Reducing antibiotic use on the neonatal unit by improving communication of blood culture results: a completed audit cycle, **F255**

complications, Neonatal long lines, **F260**

computed tomographic angiography, Brachial artery pseudoaneurysm in a premature infant, **F152**

congenital, Urine polymerase chain reaction as a screening tool for the detection of congenital cytomegalovirus infection, **F371**

congenital brachial palsy, Congenital brachial palsy: incidence, causes, and outcome in the United Kingdom and Republic of Ireland, **F185**

congenital leukaemia, Congenital leukaemia after heavy abuse of permethrin during pregnancy, **F436**

consent, Consent for clinical research in the neonatal intensive care unit: a retrospective survey and a prospective study, **F280**
Use of animal surfactant: should we seek consent?, **F351**

cord blood, S100 protein content of umbilical cord blood in healthy newborns in relation to mode of delivery, **F67**

corticosteroids, Neonatologists are using much less dexamethasone, **F432**

CPAP, Nasal CPAP for neonates: what do we know in 2003?, **F168**

crown-heel length, Updated gestational age specific birth weight, crown-heel length, and head circumference of Chinese newborns, **F229**

culture, Reducing antibiotic use on the neonatal unit by improving communication of blood culture results: a completed audit cycle, **F255**

cystic fibrosis, Cystic fibrosis newborn screening and detection of carriers, **F448**
Implications of carrier identification in newborn screening for cystic fibrosis, **F467**

cytokines, Proinflammatory and anti-inflammatory cytokine responses in preterm infants with systemic infections, **F209**

cytomegalovirus, Urine polymerase chain reaction as a screening tool for the detection of congenital cytomegalovirus infection, **F371**

de novo mutation, Sporadic case of fatal encephalopathy with neonatal onset associated with a T158M missense mutation in MECP2, **F251**

deafness, Noise levels within the ear and post-nasal space in neonates in intensive care, **F315**

dehydration, Hypernatraemic dehydration: excess sodium is not the cause, **F349**

delivery mode, S100 protein content of umbilical cord blood in healthy newborns in relation to mode of delivery, **F67**

development, Maternal docosahexaenoic acid supplementation during pregnancy and visual evoked potential development in term infants: a double blind, prospective, randomised trial, **F383**
The EPICure study: growth and associated problems in children born at 25 weeks of gestational age or less, **F492**

developmental care, Excessive exposure of sick neonates to sound during transport, **F513**

dexamethasone, Effect of a fourth *Haemophilus influenzae* type b immunisation in preterm infants who received dexamethasone for chronic lung disease, **F58**
Neonatologists are using much less dexamethasone, **F432**

diagnostic test, Diagnostic value of radiography in cases of perinatal death: a population based study, **F521**

dialysis, Early dialysis in a neonate with intrauterine lisinopril exposure, **F154**

diet, Maternal and umbilical cord erythrocyte omega-3 and omega-6 fatty acids and haemorheology in singleton and twin pregnancies, **F134**

digestion, Intestinal permeability in relation to birth weight and gestational and postnatal age, **F52**

discharge from hospital, Discharging twins separately from neonatal units, **F445**

diseases, Improved outcomes for very low birthweight infants: evidence from New Zealand national population based data, **F23**
Outcomes for high risk New Zealand newborn infants in 1998–1999: a population based, national study, **F15**

docosahexaenoic acid, Maternal docosahexaenoic acid supplementation during pregnancy and visual evoked potential development in term infants: a double blind, prospective, randomised trial, **F383**

drainage, Congenital idiopathic chylothorax in neonates: chemical pleurodesis with povidone-iodine (Betadine), **F531**

drug abuse, Treatment of neonatal abstinence syndrome, **F2**

drug exposure, Comparison of meconium and neonatal hair analysis for detection of gestational exposure to drugs of abuse, **F98**

drugs, Prenatal drug exposure and maternal and infant feeding behaviour, **F391**

early diagnosis, Limitations of ultrasonography for diagnosing white matter damage in preterm infants, **F275**

eating problems, Parental post-traumatic reactions after premature birth: implications for sleeping and eating problems in the infant, **F400**

echocardiogram, Bedside balloon atrial septostomy on neonatal units, **F339**

echocardiography, Cardiac output, pulmonary artery pressure, and patent ductus arteriosus during therapeutic cooling after global hypoxia-ischaemia, **F223**
Effect of early targeted indomethacin on the ductus arteriosus and blood flow to the upper body and brain in the preterm infant, **F477**
Evaluation of echocardiography on the neonatal unit, **F287**

elastase 1, Faecal elastase 1 levels in premature and full term infants, **F106**

endotracheal tube, Endotracheal tube fixation in neonates, **F352**

enoxaparin, Use of low molecular mass heparin (enoxaparin) in newborn infants: a prospective cohort study of 62 patients, **F365**

Erbs palsy, Congenital brachial palsy: incidence, causes, and outcome in the United Kingdom and Republic of Ireland, **F185**

erythrocyte, Maternal and umbilical cord erythrocyte omega-3 and omega-6 fatty acids and haemorheology in singleton and twin pregnancies, **F134**

erythrocytes, Decreasing oxygen saturation in very early preterm newborn infants after transfusion, **F163**

erythropoiesis, Autoimmune haemolytic anaemia in a newborn infant, **F341**

erythropoietin, Effects of vitamin E supplementation during erythropoietin treatment of the anaemia of prematurity, **F324**

ethics, Consent for clinical research in the neonatal intensive care unit: a retrospective survey and a prospective study, **F280**
Obtaining consent for neonatal research, **F166**

- ethnicity**, Effect of ethnicity on mortality of very low birthweight infants in Israel, **F333**
- evaluation**, Developing a neonatal workforce: role evolution and retention of advanced neonatal nurse practitioners, **F426**
- exchange transfusion**, Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn, **F6**
- exchange transfusions**, Treatment of jaundice in low birthweight infants, **F459**
- extracorporeal membrane oxygenation (ECMO)**, Paediatricians' perception of the use of extracorporeal membrane oxygenation to treat meconium aspiration syndrome, **F70**
Pilot investigation of hypothermia in neonates receiving extracorporeal membrane oxygenation, **F128**
- extreme prematurity**, Non-viable delivery at 20–23 weeks gestation: observations and signs of life after birth, **F199**
- extremely low birthweight infants**, A national two year follow up study of extremely low birthweight infants born in 1996–1997, **F29**
- extremely low gestational age**, Health and school performance of teenagers born before 29 weeks gestation, **F190**
- extubation**, Extubation failure due to phrenic nerve injury, **F72**
- eye-hand coordination**, Cognitive development at 5.5 years of children with chronic lung disease of prematurity, **F101**
- faecal elastase 1**, Faecal elastase 1 levels in premature and full term infants, **F106**
- family function**, Use of healthcare resources, family function, and socioeconomic support during the first four years after preterm birth, **F415**
- fatty acids**, Maternal and umbilical cord erythrocyte omega-3 and omega-6 fatty acids and haemorheology in singleton and twin pregnancies, **F134**
Maternal docosahexaenoic acid supplementation during pregnancy and visual evoked potential development in term infants: a double blind, prospective, randomised trial, **F383**
- feeding**, Multipoint measurement of intragastric pH in healthy preterm infants, **F517**
Prenatal drug exposure and maternal and infant feeding behaviour, **F391**
- feeding solids**, Factors associated with weaning in full term and preterm infants, **F296**
- fetal meconium peritonitis**, Fetal meconium peritonitis: the “vanishing hydrocele” sign, **F74**
- fever, dehydration**, Fever in healthy asymptomatic newborns during the first days of life, **F312**
- follow up**, Health and school performance of teenagers born before 29 weeks gestation, **F190**
Use of healthcare resources, family function, and socioeconomic support during the first four years after preterm birth, **F415**
- formula**, Donor human milk versus formula for preventing necrotising enterocolitis in preterm infants: systematic review, **F11**
- formula feeding**, Neonatal weight loss in breast and formula fed infants, **F472**
- formula milk**, Factors associated with weaning in full term and preterm infants, **F296**
- fractional oxygen extraction**, The light still shines, but not that brightly? The current status of perinatal near infrared spectroscopy, **F263**
- fungaemia**, Methaemoglobinaemia with concurrent blood isolation of *Saccharomyces* and *Candida*, **F446**
- gastric acidity**, Multipoint measurement of intragastric pH in healthy preterm infants, **F517**
- gastrointestinal perforation**, Neonatal gastrointestinal perforation, **F75**
- gastrointestinal tract**, Insulin in human milk: postpartum changes and effect of gestational age, **F214**
Probiotics for preterm infants?, **F354**
- gastro-oesophageal reflux**, Multipoint measurement of intragastric pH in healthy preterm infants, **F517**
- gestation**, Updated gestational age specific birth weight, crown-heel length, and head circumference of Chinese newborns, **F229**
- gestational age**, Insulin in human milk: postpartum changes and effect of gestational age, **F214**
Intestinal permeability in relation to birth weight and gestational and postnatal age, **F52**
- glucagon**, An inadequate glycaemic response to glucagon is linked to insulin resistance in preterm infants?, **F62**
- glucose**, An inadequate glycaemic response to glucagon is linked to insulin resistance in preterm infants?, **F62**
- glucose-6-phosphate dehydrogenase deficiency**, Massive acute haemolysis in neonates with glucose-6-phosphate dehydrogenase deficiency, **F534**
- Gram negative organisms**, Neonatal sepsis in Peshawar., **F160**
- growth**, Growth and body composition in preterm infants with bronchopulmonary dysplasia, **F46**
Neonatal weight loss in breast and formula fed infants, **F472**
The EPICure study: growth and associated problems in children born at 25 weeks of gestational age or less, **F492**
Weaning preterm infants: a randomised controlled trial, **F302**
- growth standards**, Updated gestational age specific birth weight, crown-heel length, and head circumference of Chinese newborns, **F229**
- gut**, Neonatal gastrointestinal perforation, **F75**
- haemoglobin**, Decreasing oxygen saturation in very early preterm newborn infants after transfusion, **F163**
- haemolysis**, Massive acute haemolysis in neonates with glucose-6-phosphate dehydrogenase deficiency, **F534**
- haemolytic disease**, Late anaemia in Rh haemolytic disease, **F444**
- haemolytic disease of newborns**, High dose intravenous immunoglobulin in haemolytic disease of neonates, **F444**
Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn, **F444**
- Haemophilus influenzae type b vaccine**, Effect of a fourth *Haemophilus influenzae* type b immunisation in preterm infants who received dexamethasone for chronic lung disease, **F58**
- haemorheology**, Maternal and umbilical cord erythrocyte omega-3 and omega-6 fatty acids and haemorheology in singleton and twin pregnancies, **F134**
- hair analysis**, Comparison of meconium and neonatal hair analysis for detection of gestational exposure to drugs of abuse, **F98**
- Hashimoto's thyroiditis**, Do we need to assess the thyroid function in the infants of mothers with Hashimoto's thyroiditis?, **F258**
- head circumference**, Updated gestational age specific birth weight, crown-heel length, and head circumference of Chinese newborns, **F229**
- hearing loss**, Noise levels within the ear and post-nasal space in neonates in intensive care, **F315**
- heart**, Evaluation of echocardiography on the neonatal unit, **F287**
Neonatal murmurs: are senior house officers good enough?, **F147**
- heart rate**, Motion resistant pulse oximetry in neonates, **F505**
- heat loss**, Swaddling and heat loss, **F256**
- herpes simplex virus**, Relapse of neonatal herpes simplex virus infection, **F483**
- high dose intravenous immunoglobulin**, High dose intravenous immunoglobulin in haemolytic disease of neonates, **F444**
Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn, **F444**
- history**, Andreas Vesalius (1514–1564), Padua, and the fetal “shunts”, **F157**
Galen (AD 129–200) of Pergamun: anatomist and experimental physiologist, **F441**
Hugh Downman, MD (1740–1809) of Exeter and his poem on infant care, **F253**
- human milk**, Insulin in human milk: postpartum changes and effect of gestational age, **F214**
- hydrocele**, Fetal meconium peritonitis: the “vanishing hydrocele” sign, **F74**
- hyperbilirubinaemia**, Intestinal absorption of mixed micellar phyloquinone (vitamin K₁) is unreliable in infants with conjugated hyperbilirubinaemia: implications for oral prophylaxis of vitamin K deficiency bleeding, **F113**
Massive acute haemolysis in neonates with glucose-6-phosphate dehydrogenase deficiency, **F534**
Phototherapy with turquoise versus blue light, **F430**
Transcutaneous bilirubin measurement in newborn infants: evaluation of a new spectrophotometric method, **F350**
- hypercalcaemia**, Subcutaneous fat necrosis of the newborn: hypercalcaemia with hepatic and atrial myocardial calcification, **F343**
- hypernatraemia**, Hypernatraemic dehydration: excess sodium is not the cause, **F349**
- hyponatraemia**, Hyponatraemia as a consequence of serial liquor punctures in preterm infants with a ventricular access device after posthaemorrhagic hydrocephalus, **F351**

- hypotension**, Hypotension in the very low birthweight infant: the old, the new, and the uncertain, **F450**
- hypothermia**, Cardiac output, pulmonary artery pressure, and patent ductus arteriosus during therapeutic cooling after global hypoxia-ischaemia, **F223**
Pilot investigation of hypothermia in neonates receiving extracorporeal membrane oxygenation, **F128**
Preventing hypothermia at birth in preterm babies: at a cost of overheating some?, **F256**
- hypoxia-ischaemia**, Cardiac output, pulmonary artery pressure, and patent ductus arteriosus during therapeutic cooling after global hypoxia-ischaemia, **F223**
- immunisation**, Effect of a fourth *Haemophilus influenzae* type b immunisation in preterm infants who received dexamethasone for chronic lung disease, **F58**
- incidence**, Congenital brachial palsy: incidence, causes, and outcome in the United Kingdom and Republic of Ireland, **F185**
- indomethacin**, Effect of early targeted indomethacin on the ductus arteriosus and blood flow to the upper body and brain in the preterm infant, **F477**
- infant**, Infant to staff ratios and risk of mortality in very low birthweight infants, **F94**
Nasal CPAP for neonates: what do we know in 2003?, **F168**
Neonatal long lines, **F260**
Neonatal meningitis, **F173**
- infants**, SIDS, smoking, and arousal thresholds: conclusions not supported by data, **F162**
- infarction**, Neonatal cerebral infarction and visual function at school age, **F487**
- infection**, Fever in healthy asymptomatic newborns during the first days of life, **F312**
Neonatal sepsis in Peshawar:, **F160**
Proinflammatory and anti-inflammatory cytokine responses in preterm infants with systemic infections, **F209**
Urine polymerase chain reaction as a screening tool for the detection of congenital cytomegalovirus infection, **F371**
- infection control**, Infection control for SARS in a tertiary neonatal centre, **F405**
- inflammatory response syndrome**, Increased serum levels of interleukin 6 are associated with severe intraventricular haemorrhage in extremely premature infants, **F501**
- informed consent**, Obtaining consent for neonatal research, **F166**
- inotropes**, Hypotension in the very low birthweight infant: the old, the new, and the uncertain, **F450**
- insecticide**, Congenital leukaemia after heavy abuse of permethrin during pregnancy, **F436**
- insulin**, An inadequate glycaemic response to glucagon is linked to insulin resistance in preterm infants?, **F62**
Insulin in human milk: postpartum changes and effect of gestational age, **F214**
- insulin resistance**, An inadequate glycaemic response to glucagon is linked to insulin resistance in preterm infants?, **F62**
- intensive care**, Improved outcomes for very low birthweight infants: evidence from New Zealand national population based data, **F23**
- intensive care units**, Noise levels within the ear and post-nasal space in neonates in intensive care, **F315**
- interleukin 6**, Increased serum levels of interleukin 6 are associated with severe intraventricular haemorrhage in extremely premature infants, **F501**
- interleukin 10**, Decreased neutrophil apoptosis in tracheal fluids of preterm infants at risk of chronic lung disease, **F245**
- interleukin 11**, Neonatal thrombocytopenia: causes and management, **F359**
- intestinal permeability**, Intestinal permeability in relation to birth weight and gestational and postnatal age, **F52**
- intracranial pressure**, Progressive ventricular dilatation (PVD) over the past 22 years, **F257**
- intrauterine exposure**, Early dialysis in a neonate with intrauterine lisinopril exposure, **F154**
- intravenous immunoglobulin**, Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn, **F6**
- intraventricular haemorrhage**, Increased serum levels of interleukin 6 are associated with severe intraventricular haemorrhage in extremely premature infants, **F501**
Progressive ventricular dilatation (PVD) over the past 22 years, **F257**
- Prophylactic indomethacin for preterm infants: a systematic review and meta-analysis, **F464**
- intraventricular haemorrhage: patent ductus arteriosus**, Effect of early targeted indomethacin on the ductus arteriosus and blood flow to the upper body and brain in the preterm infant, **F477**
- intubation**, Changes in resuscitation practice at birth, **F375**
- iodine**, Iodine deficiency associated with parenteral nutrition in extreme preterm infants, **F56**
- iron**, Weaning preterm infants: a randomised controlled trial, **F302**
- jaundice**, Jaundice in low birthweight infants: pathobiology and outcome, **F455**
Phototherapy with turquoise versus blue light, **F430**
Plasma bilirubin level and oxidative stress in preterm infants, **F119**
Treatment of jaundice in low birthweight infants, **F459**
- Jaundice Meter**, Transcutaneous bilirubin measurement in newborn infants: evaluation of a new spectrophotometric method, **F350**
- Jewish**, Effect of ethnicity on mortality of very low birthweight infants in Israel, **F333**
- kernicterus**, Jaundice in low birthweight infants: pathobiology and outcome, **F455**
Treatment of jaundice in low birthweight infants, **F459**
- kidney**, Early dialysis in a neonate with intrauterine lisinopril exposure, **F154**
- lactation**, Effect of caesarean section on breast milk transfer to the normal term newborn over the first week of life, **F380**
- late onset**, A ten year, multicentre study of coagulase negative staphylococcal infections in Australasian neonatal units, **F89**
- lenticulostriate**, Neonatal lenticulostriate vasculopathy: further characterisation, **F410**
- leukaemia**, Congenital leukaemia after heavy abuse of permethrin during pregnancy, **F436**
- limb cooling**, Effect of limb cooling on peripheral and global oxygen consumption in neonates, **F139**
- liquor puncture**, Hyponatraemia as a consequence of serial liquor punctures in preterm infants with a ventricular access device after posthaemorrhagic hydrocephalus, **F351**
- lisinopril**, Early dialysis in a neonate with intrauterine lisinopril exposure, **F154**
- liver**, Intestinal absorption of mixed micellar phyloquinone (vitamin K₁) is unreliable in infants with conjugated hyperbilirubinaemia: implications for oral prophylaxis of vitamin K deficiency bleeding, **F113**
Progress in treatment and outcome for children with neonatal haemochromatosis, **F124**
- low molecular mass heparin**, Use of low molecular mass heparin (enoxaparin) in newborn infants: a prospective cohort study of 62 patients, **F365**
- magnetic resonance imaging**, Cerebral perfusion in infants and neonates: preliminary results obtained using dynamic susceptibility contrast enhanced magnetic resonance imaging, **F525**
Limitations of ultrasonography for diagnosing white matter damage in preterm infants, **F275**
Low field strength magnetic resonance imaging of the neonatal brain, **F203**
Magnetic resonance imaging of preterm brain injury, **F269**
Neonatal cerebral infarction and visual function at school age, **F487**
- Maleco catheter**, Unusual complication of a central venous line in a neonate, **F440**
- maternal**, Management of mothers of neonates with vertically transmitted sexually transmitted infections, **F163**
- maternal-infant interaction**, Prenatal drug exposure and maternal and infant feeding behaviour, **F391**
- meconium analysis**, Comparison of meconium and neonatal hair analysis for detection of gestational exposure to drugs of abuse, **F98**
- meconium aspiration syndrome**, Paediatricians' perception of the use of extracorporeal membrane oxygenation to treat meconium aspiration syndrome, **F70**
- meningitis**, A ten year, multicentre study of coagulase negative staphylococcal infections in Australasian neonatal units, **F89**
Long term outcome of neonatal meningitis, **F179**
Neonatal meningitis, **F173**
- metabolic rate**, Effect of a change in global metabolic rate on peripheral oxygen consumption in neonates, **F143**

- Effect of limb cooling on peripheral and global oxygen consumption in neonates, **F139**
- methaemoglobinaemia**, Methaemoglobinaemia with concurrent blood isolation of *Saccharomyces* and *Candida*, **F446**
- microbial colonisation**, Probiotics for preterm infants?, **F354**
- milk contamination**, Contamination of a milk bank pasteuriser causing a *Pseudomonas aeruginosa* outbreak in a neonatal intensive care unit, **F434**
- mortality**, Diagnostic value of radiography in cases of perinatal death: a population based study, **F521**
Effect of ethnicity on mortality of very low birthweight infants in Israel, **F333**
Excess risk of mortality in very low birthweight triplets: a national, population based study, **F36**
Infant to staff ratios and risk of mortality in very low birthweight infants, **F94**
- multidrug resistance**, Neonatal sepsis in Peshawar., **F160**
- multiorgan failure**, Early dialysis in a neonate with intrauterine lisinopril exposure, **F154**
- multiple pregnancy**, Excess risk of mortality in very low birthweight triplets: a national, population based study, **F36**
- murmur**, Neonatal murmurs: are senior house officers good enough?, **F147**
- Muslim**, Effect of ethnicity on mortality of very low birthweight infants in Israel, **F333**
- naloxone**, Naloxone for narcotic exposed newborn infants: systematic review, **F308**
- narcotic exposure**, Naloxone for narcotic exposed newborn infants: systematic review, **F308**
- near infrared spectroscopy**, Effect of a change in global metabolic rate on peripheral oxygen consumption in neonates, **F143**
The light still shines, but not that brightly? The current status of perinatal near infrared spectroscopy, **F263**
- necrotising enterocolitis**, A ten year, multicentre study of coagulase negative staphylococcal infections in Australasian neonatal units, **F89**
Donor human milk versus formula for preventing necrotising enterocolitis in preterm infants: systematic review, **F11**
Probiotics for preterm infants?, **F354**
- neonatal**, Neonatal shaken baby syndrome—historical inexactitudes, **F161**
Neonatal shaken baby syndrome—lessons to be learned, **F161**
Swaddling and heat loss, **F256**
- neonatal abstinence syndrome**, Treatment of neonatal abstinence syndrome, **F2**
- neonatal haemochromatosis**, Progress in treatment and outcome for children with neonatal haemochromatosis, **F124**
- neonatal intensive care**, Outcomes for high risk New Zealand newborn infants in 1998–1999: a population based, national study, **F15**
- neonatal nurse practitioner**, Developing a neonatal workforce: role evolution and retention of advanced neonatal nurse practitioners, **F426**
- neonatal resuscitation**, Changes in resuscitation practice at birth, **F375**
- neonate**, Nasal CPAP for neonates: what do we know in 2003?, **F168**
- neurodevelopment**, Long term outcome of neonatal meningitis, **F179**
Prophylactic indomethacin for preterm infants: a systematic review and meta-analysis, **F464**
- neurodevelopmental outcome**, A national two year follow up study of extremely low birthweight infants born in 1996–1997, **F29**
- newborn**, Infant to staff ratios and risk of mortality in very low birthweight infants, **F94**
Neonatal meningitis, **F173**
- newborns**, Infection control for SARS in a tertiary neonatal centre, **F405**
- noise intensities**, Noise levels within the ear and post-nasal space in neonates in intensive care, **F315**
- non-contact oxygen delivery**, Wafting does work, **F446**
- nosocomial infection**, Contamination of a milk bank pasteuriser causing a *Pseudomonas aeruginosa* outbreak in a neonatal intensive care unit, **F434**
- nosocomial infections**, A ten year, multicentre study of coagulase negative staphylococcal infections in Australasian neonatal units, **F89**
- obstetrical brachial palsy**, Congenital brachial palsy: incidence, causes, and outcome in the United Kingdom and Republic of Ireland, **F185**
- OMIM 603528**, Dehydrated hereditary stomatocytosis with transient perinatal ascites, **F438**
- ontogenesis**, Faecal elastase 1 levels in premature and full term infants, **F106**
- opiates**, Naloxone for narcotic exposed newborn infants: systematic review, **F308**
Prenatal drug exposure and maternal and infant feeding behaviour, **F391**
- outcome**, Congenital brachial palsy: incidence, causes, and outcome in the United Kingdom and Republic of Ireland, **F185**
Long term outcome of neonatal meningitis, **F179**
- overdose**, Intravenous propacetamol overdose in a term newborn, **F351**
- oxidative stress**, Plasma bilirubin level and oxidative stress in preterm infants, **F119**
- oxygen consumption**, Effect of a change in global metabolic rate on peripheral oxygen consumption in neonates, **F143**
Effect of limb cooling on peripheral and global oxygen consumption in neonates, **F139**
The light still shines, but not that brightly? The current status of perinatal near infrared spectroscopy, **F263**
- oxygen saturation**, Decreasing oxygen saturation in very early preterm newborn infants after transfusion, **F163**
Motion resistant pulse oximetry in neonates, **F505**
- oxygen treatment**, Oxygen administration in infants, **F84**
- pancreas**, Faecal elastase 1 levels in premature and full term infants, **F106**
- paracetamol**, Intravenous propacetamol overdose in a term newborn, **F351**
- parenteral nutrition**, Iodine deficiency associated with parenteral nutrition in extreme preterm infants, **F56**
- parents**, Discharging twins separately from neonatal units, **F445**
- parents' views**, Use of animal surfactant: should we seek consent?, **F351**
- patent ductus arteriosus**, Prophylactic indomethacin for preterm infants: a systematic review and meta-analysis, **F464**
- percutaneous long line**, Pericardial effusion and cardiac tamponade as complications of neonatal long lines: are they really a problem?, **F292**
- pericardial effusion**, Pericardial effusion and cardiac tamponade as complications of neonatal long lines: are they really a problem?, **F292**
- perinatal**, Diagnostic value of radiography in cases of perinatal death: a population based study, **F521**
- perinatal risks**, Parental post-traumatic reactions after premature birth: implications for sleeping and eating problems in the infant, **F400**
- permethrin**, Congenital leukaemia after heavy abuse of permethrin during pregnancy, **F436**
- pH monitoring**, Multipoint measurement of intragastric pH in healthy preterm infants, **F517**
- phototherapy**, Phototherapy with turquoise versus blue light, **F430**
Treatment of jaundice in low birthweight infants, **F459**
- phrenic nerve**, Extubation failure due to phrenic nerve injury, **F72**
- physiology**, Galen (AD 129–200) of Pergamun: anatomist and experimental physiologist, **F441**
- placental respiration**, Dr Erasmus Darwin (1731–1802) of Lichfield and placental respiration, **F346**
- platelets**, Neonatal thrombocytopenia: causes and management, **F359**
- pleurodesis**, Congenital idiopathic chylothorax in neonates: chemical pleurodesis with povidone-iodine (Betadine), **F531**
- polymerase chain reaction (PCR)**, Urine polymerase chain reaction as a screening tool for the detection of congenital cytomegalovirus infection, **F371**
- posthaemorrhagic hydrocephalus**, Hyponatraemia as a consequence of serial liquor punctures in preterm infants with a ventricular access device after posthaemorrhagic hydrocephalus, **F351**
- posthaemorrhagic ventricular dilatation**, Serum S-100 protein does not correlate with cerebral ultrasound scans in preterm infants, **F160**
- povidone-iodine**, Congenital idiopathic chylothorax in neonates: chemical pleurodesis with povidone-iodine (Betadine), **F531**
- pregnancy**, Maternal and umbilical cord erythrocyte omega-3 and omega-6 fatty acids and haemorheology in singleton and twin pregnancies, **F134**
- pre-leukaemic fusion genes**, Absence of leukaemic fusion gene transcripts in preterm infants exposed to diagnostic x rays, **F237**
- premature**, Decreased neutrophil apoptosis in tracheal fluids of preterm infants at risk of chronic lung disease, **F245**
Faecal elastase 1 levels in premature and full term infants, **F106**

- Improved outcomes for very low birthweight infants: evidence from New Zealand national population based data, **F23**
- Outcomes for high risk New Zealand newborn infants in 1998–1999: a population based, national study, **F15**
- premature infant**, Effect of early targeted indomethacin on the ductus arteriosus and blood flow to the upper body and brain in the preterm infant, **F477**
- Multipoint measurement of intragastric pH in healthy preterm infants, **F517**
- prematurity**, A national two year follow up study of extremely low birthweight infants born in 1996–1997, **F29**
- Effects of gestational age and prenatal and perinatal events on the coagulation status in premature infants, **F319**
- Extubation failure due to phrenic nerve injury, **F72**
- Frequent wheeze at follow up of very preterm infants: which factors are predictive?, **F329**
- Jaundice in low birthweight infants: pathobiology and outcome, **F455**
- The EPICure study: growth and associated problems in children born at 25 weeks of gestational age or less, **F492**
- Treatment of jaundice in low birthweight infants, **F459**
- prescribing error**, Intravenous propacetamol overdose in a term newborn, **F351**
- preterm**, An inadequate glycaemic response to glucagon is linked to insulin resistance in preterm infants?, **F62**
- Blood transfusion in preterm neonates, **F78**
- Cognitive development at 5.5 years of children with chronic lung disease of prematurity, **F101**
- Decreasing oxygen saturation in very early preterm newborn infants after transfusion, **F163**
- Donor human milk versus formula for preventing necrotising enterocolitis in preterm infants: systematic review, **F11**
- Growth and body composition in preterm infants with bronchopulmonary dysplasia, **F46**
- Hyponatraemia as a consequence of serial liquor punctures in preterm infants with a ventricular access device after posthaemorrhagic hydrocephalus, **F351**
- Iodine deficiency associated with parenteral nutrition in extreme preterm infants, **F56**
- Limitations of ultrasonography for diagnosing white matter damage in preterm infants, **F275**
- Neonatal thrombocytopenia: causes and management, **F359**
- Neonatologists are using much less dexamethasone, **F432**
- Parental post-traumatic reactions after premature birth: implications for sleeping and eating problems in the infant, **F400**
- Plasma bilirubin level and oxidative stress in preterm infants, **F119**
- Preventing hypothermia at birth in preterm babies: at a cost of overheating some?, **F256**
- Probiotics for preterm infants?, **F354**
- Proinflammatory and anti-inflammatory cytokine responses in preterm infants with systemic infections, **F209**
- Weaning preterm infants: a randomised controlled trial, **F302**
- preterm brain injury**, Magnetic resonance imaging of preterm brain injury, **F269**
- preterm infants**, Absence of leukaemic fusion gene transcripts in preterm infants exposed to diagnostic x rays, **F237**
- Excess risk of mortality in very low birthweight triplets: a national, population based study, **F36**
- Increased serum levels of interleukin 6 are associated with severe intraventricular haemorrhage in extremely premature infants, **F501**
- Recombinant erythropoietin and blood transfusion in selected preterm infants, **F41**
- Use of healthcare resources, family function, and socioeconomic support during the first four years after preterm birth, **F415**
- probiotics**, Probiotics for preterm infants?, **F354**
- progressive ventricular dilatation**, Progressive ventricular dilatation (PVD) over the past 22 years, **F257**
- propacetamol**, Intravenous propacetamol overdose in a term newborn, **F351**
- prophylaxis**, Oral mixed micellar vitamin K for prevention of late vitamin K deficiency bleeding, **F109**
- pseudoaneurysms**, Brachial artery pseudoaneurysm in a premature infant, **F152**
- Pseudomonas aeruginosa**, Contamination of a milk bank pasteuriser causing a *Pseudomonas aeruginosa* outbreak in a neonatal intensive care unit, **F434**
- psychosocial issues**, Implications of carrier identification in newborn screening for cystic fibrosis, **F467**
- radiography**, Diagnostic value of radiography in cases of perinatal death: a population based study, **F521**
- randomised double blind controlled trial**, Recombinant erythropoietin and blood transfusion in selected preterm infants, **F41**
- rebreathing**, Respiratory responses to hypoxia/hypercapnia in small for gestational age infants influenced by maternal smoking, **F217**
- recurrence risk**, Sporadic case of fatal encephalopathy with neonatal onset associated with a T158M missense mutation in MECP2, **F251**
- relapse**, Relapse of neonatal herpes simplex virus infection, **F483**
- research**, Obtaining consent for neonatal research, **F166**
- research ethics**, Consent for clinical research in the neonatal intensive care unit: a retrospective survey and a prospective study, **F280**
- respiratory failure**, Pilot investigation of hypothermia in neonates receiving extracorporeal membrane oxygenation, **F128**
- respiratory mechanics**, Oxygen administration in infants, **F84**
- resuscitation**, Preventing hypothermia at birth in preterm babies: at a cost of overheating some?, **F256**
- retroperitoneal mass**, Unusual complication of a central venous line in a neonate, **F440**
- Retz syndrome**, Sporadic case of fatal encephalopathy with neonatal onset associated with a T158M missense mutation in MECP2, **F251**
- rhesus haemolytic disease**, Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn, **F6**
- S100 protein**, S100 protein content of umbilical cord blood in healthy newborns in relation to mode of delivery, **F67**
- S-100 protein**, Serum S-100 protein does not correlate with cerebral ultrasound scans in preterm infants, **F160**
- Saccharomyces**, Methaemoglobinaemia with concurrent blood isolation of *Saccharomyces* and *Candida*, **F446**
- SARS**, Infection control for SARS in a tertiary neonatal centre, **F405**
- school age**, Neonatal cerebral infarction and visual function at school age, **F487**
- school performance**, Health and school performance of teenagers born before 29 weeks gestation, **F190**
- screening**, Cystic fibrosis newborn screening and detection of carriers, **F448**
- Implications of carrier identification in newborn screening for cystic fibrosis, **F467**
- Urine polymerase chain reaction as a screening tool for the detection of congenital cytomegalovirus infection, **F371**
- senior house officers**, Neonatal murmurs: are senior house officers good enough?, **F147**
- sepsis**, Neonatal sepsis in Peshawar:, **F160**
- sexually transmitted infections**, Management of mothers of neonates with vertically transmitted sexually transmitted infections, **F163**
- shaken baby syndrome**, Neonatal shaken baby syndrome—historical inexactitudes, **F161**
- Neonatal shaken baby syndrome—lessons to be learned, **F161**
- SIDS**, Respiratory responses to hypoxia/hypercapnia in small for gestational age infants influenced by maternal smoking, **F217**
- signal extraction technology**, Motion resistant pulse oximetry in neonates, **F505**
- sleeping problems**, Parental post-traumatic reactions after premature birth: implications for sleeping and eating problems in the infant, **F400**
- small for gestational age**, Effects of gestational age and prenatal and perinatal events on the coagulation status in premature infants, **F319**
- small intestine**, Intestinal permeability in relation to birth weight and gestational and postnatal age, **F52**
- smoking**, Respiratory responses to hypoxia/hypercapnia in small for gestational age infants influenced by maternal smoking, **F217**
- SIDS, smoking, and arousal thresholds: conclusions not supported by data, **F162**
- social class**, Social trends in singleton births and birth weight in Wirral residents, 1990–2001, **F421**
- socioeconomic factors**, Use of healthcare resources, family function, and socioeconomic support during the first four years after preterm birth, **F415**
- sound**, Excessive exposure of sick neonates to sound during transport, **F513**
- staff**, Infant to staff ratios and risk of mortality in very low birthweight infants, **F94**

- Staphylococcus epidermidis**, A ten year, multicentre study of coagulase negative staphylococcal infections in Australasian neonatal units, **F89**
- stomatocytosis**, Dehydrated hereditary stomatocytosis with transient perinatal ascites, **F438**
- stress**, Excessive exposure of sick neonates to sound during transport, **F513**
- subcutaneous fat necrosis of the newborn**, Subcutaneous fat necrosis of the newborn: hypercalcaemia with hepatic and atrial myocardial calcification, **F343**
- sudden infant death syndrome**, SIDS, smoking, and arousal thresholds: conclusions not supported by data, **F162**
- supraventricular tachycardia**, Amiodarone and breast feeding, **F255**
Swallowing induced supraventricular tachycardia in a neonate, **F162**
- surfactant**, Use of animal surfactant: should we seek consent?, **F351**
- swaddling**, Swaddling and heat loss, **F256**
- swallowing**, Swallowing induced supraventricular tachycardia in a neonate, **F162**
- temperature control**, Swaddling and heat loss, **F256**
- thalamus**, Neonatal lenticulostriate vasculopathy: further characterisation, **F410**
- thrombocytopenia**, Neonatal thrombocytopenia: causes and management, **F359**
Subcutaneous fat necrosis of the newborn: hypercalcaemia with hepatic and atrial myocardial calcification, **F343**
- thromboembolism**, Use of low molecular mass heparin (enoxaparin) in newborn infants: a prospective cohort study of 62 patients, **F365**
- thrombosis**, Use of low molecular mass heparin (enoxaparin) in newborn infants: a prospective cohort study of 62 patients, **F365**
- thyroid**, Iodine deficiency associated with parenteral nutrition in extreme preterm infants, **F56**
- thyroid disorders**, Do we need to assess the thyroid function in the infants of mothers with Hashimoto's thyroiditis?, **F258**
- tissue perfusion**, Hypotension in the very low birthweight infant: the old, the new, and the uncertain, **F450**
- tissue plasminogen activator**, Brachial artery pseudoaneurysm in a premature infant, **F152**
- T-piece**, Changes in resuscitation practice at birth, **F375**
- transcutaneous bilirubinometry**, Transcutaneous bilirubin measurement in newborn infants: evaluation of a new spectrophotometric method, **F350**
- transfusion**, Massive acute haemolysis in neonates with glucose-6-phosphate dehydrogenase deficiency, **F534**
- transfusion requirements**, Recombinant erythropoietin and blood transfusion in selected preterm infants, **F41**
- transplantation**, Progress in treatment and outcome for children with neonatal haemochromatosis, **F124**
- transport**, Excessive exposure of sick neonates to sound during transport, **F513**
Neonatal transfers by advanced neonatal nurse practitioners and paediatric registrars, **F509**
- transposition of the great arteries**, Bedside balloon atrial septostomy on neonatal units, **F339**
- treating infected neonatal lines**, Risks of treating infected neonatal lines, **F163**
- triplets**, Excess risk of mortality in very low birthweight triplets: a national, population based study, **F36**
- twins**, Discharging twins separately from neonatal units, **F445**
- ultrasonography**, Limitations of ultrasonography for diagnosing white matter damage in preterm infants, **F275**
- ultrasound**, Low field strength magnetic resonance imaging of the neonatal brain, **F203**
Serum S-100 protein does not correlate with cerebral ultrasound scans in preterm infants, **F160**
- ultrasound guided compression obliteration**, Brachial artery pseudoaneurysm in a premature infant, **F152**
- umbilical cord**, Dr Erasmus Darwin (1731–1802) of Lichfield and placental respiration, **F346**
- umbilical granulomas**, Umbilical granulomas: a randomised controlled trial, **F257**
- vaginal delivery**, Effect of caesarean section on breast milk transfer to the normal term newborn over the first week of life, **F380**
- vasculopathy**, Neonatal lenticulostriate vasculopathy: further characterisation, **F410**
- ventilatory support**, Endotracheal tube fixation in neonates, **F352**
- very low birth weight**, Effect of ethnicity on mortality of very low birthweight infants in Israel, **F333**
Hypotension in the very low birthweight infant: the old, the new, and the uncertain, **F450**
Improved outcomes for very low birthweight infants: evidence from New Zealand national population based data, **F23**
Prophylactic indomethacin for preterm infants: a systematic review and meta-analysis, **F464**
- very low birth weight, Australia**, Infant to staff ratios and risk of mortality in very low birthweight infants, **F94**
- viability**, Non-viable delivery at 20–23 weeks gestation: observations and signs of life after birth, **F199**
- vision**, Neonatal cerebral infarction and visual function at school age, **F487**
- visual evoked potential**, Maternal docosahexaenoic acid supplementation during pregnancy and visual evoked potential development in term infants: a double blind, prospective, randomised trial, **F383**
- vitamin E**, Effects of vitamin E supplementation during erythropoietin treatment of the anaemia of prematurity, **F324**
- vitamin K**, Intestinal absorption of mixed micellar phylloquinone (vitamin K₁) is unreliable in infants with conjugated hyperbilirubinaemia: implications for oral prophylaxis of vitamin K deficiency bleeding, **F113**
Vitamin K—what, why, and when, **F80**
- vitamin K deficiency bleeding**, Oral mixed micellar vitamin K for prevention of late vitamin K deficiency bleeding, **F109**
Vitamin K—what, why, and when, **F80**
- wafting**, Wafting does work, **F446**
- weaning**, Factors associated with weaning in full term and preterm infants, **F296**
Weaning preterm infants: a randomised controlled trial, **F302**
- weight loss**, Neonatal weight loss in breast and formula fed infants, **F472**
- white matter damage**, Limitations of ultrasonography for diagnosing white matter damage in preterm infants, **F275**
- workforce**, Developing a neonatal workforce: role evolution and retention of advanced neonatal nurse practitioners, **F426**
- wrapping**, Preventing hypothermia at birth in preterm babies: at a cost of overheating some?, **F256**
- X chromosome inactivation**, Sporadic case of fatal encephalopathy with neonatal onset associated with a T158M missense mutation in MECP2, **F251**