

ORIGINAL ARTICLE

The Scottish perinatal neuropathology study: clinicopathological correlation in early neonatal deaths

J C Becher, J E Bell, J W Keeling, N McIntosh, B Wyatt

Arch Dis Child Fetal Neonatal Ed 2004;**89**:F399–F407. doi: 10.1136/adc.2003.037606



The supplementary tables can be found at <http://adc.bmjournals.com/supplemental/>.

See end of article for authors' affiliations

Correspondence to: Professor McIntosh, Department of Child Life and Health, Reproductive and Developmental Medicine, 20 Sylvan Place, Edinburgh EH9 1UW, Scotland, UK; neil.mcintosh@ed.ac.uk

Accepted
4 November 2003

Background: A proportion of neonatal deaths from asphyxia have been shown to be associated with pre-existing brain injury.

Objectives: (a) To compare the epidemiology of infants displaying signs of birth asphyxia with those not showing signs; (b) to examine the neuropathology and determine if possible the timing of brain insult comparing asphyxiated with non-asphyxiated infants; (c) to compare the clinical features of those born with birth asphyxia with and without pre-labour damage.

Methods: Over a two year period, all 22 Scottish delivery units collected clinical details on early neonatal deaths. Requests for post mortem included separate requests for detailed neuropathological examination of the brain. Infants were classified into two groups: birth asphyxia and non-birth asphyxia. Clinicopathological correlation was used to attempt to define the time of brain insult.

Results: Detailed clinical data were available on 137 of 174 early neonatal deaths that met the inclusion criteria. Seventy of 88 parents who had agreed to post mortem examination consented to a detailed examination of additional samples from the brain; in 53 of these cases the infant was born in an asphyxiated condition. All asphyxiated and encephalopathic infants, 38% of mature and 52% of preterm infants with features of birth asphyxia but without encephalopathy, and only one of 12 infants without any signs of birth asphyxia showed damage consistent with onset before the start of labour.

Conclusions: In a large proportion of neonatal deaths, brain injury predates the onset of labour. This is more common in infants born in an asphyxiated condition.

The three major causes of neonatal death are lethal malformations, prematurity, and birth asphyxia.¹ Whereas the general public considers major malformations and premature birth as unavoidable mischance, birth asphyxia implies a lack of care in labour. Although birth asphyxia is classically linked to intrapartum hypoxia-ischaemia in full term infants, often proceeding to a neonatal encephalopathy—the so-called hypoxic-ischaemic encephalopathy, a proportion of preterm babies are also born in a neurologically depressed condition almost certainly related to poor oxygenation in labour. Asphyxia is acknowledged to be an imprecise term, but is still used regularly by the profession and parents. It may be implied by one or more of the following features: a low Apgar score²⁻⁷; a baby who is difficult to resuscitate; metabolic acidosis in either the cord^{5, 8, 9} or early neonatal blood samples; the development of neonatal encephalopathy.¹⁰⁻¹² A history of these particular features may be sought retrospectively if an infant goes on to develop neurodevelopmental delay. None of these indicators, when applied prospectively to infants born in poor condition, has good sensitivity, specificity, or predictive value for neurodevelopmental delay or disability, although in full term infants the development of neonatal encephalopathy is more specific.¹³ It is clear that perinatal asphyxia is not likely to be an important factor in the development of every case of neonatal encephalopathy or in most cases of cerebral palsy.¹⁴⁻¹⁹ This view has been endorsed by a statement from the International CP Task Force.^{20, 21}

Obstetric care has seen dramatic changes over the last few decades. Most changes have contributed to the steadily falling stillbirth and neonatal death rates.^{1, 22-25} However, despite better clinical care and widespread use of fetal

monitoring and fetal blood sampling, full term infants continue to be born in a neurologically depressed condition. Such infants cause considerable distress to parents and staff. They contribute both to early neonatal mortality and to the pool of children who display later neurodevelopmental disability with cerebral palsy. Although in some cases obstetric risk factors can be identified, affected children also result from pregnancies and labours that, even when scrutinised critically, appear to be normal.

Litigation for perceived perinatal mismanagement is increasingly common, particularly in relation to infants born in a neurologically depressed condition—usually manifested by a poor Apgar score—and often reflexly labelled birth asphyxia. Some recent anecdotal reports and small series of infants born in poor condition have shown neuropathological abnormality at autopsy that must have preceded the onset of labour. These generally represent the collected experience of specialist referral centres²⁶⁻³⁰ or focus on a particular age group of infants—for example, those born preterm.³¹⁻³⁵ Only rarely do the neuropathological studies include correlation with clinical factors.³⁶⁻³⁸ Further insights into cliniconuropathological correlation in the first weeks of life are now being achieved by neuroimaging.³⁹ This Scottish study was set up to identify neuropathological abnormalities in a population cohort of perinatal deaths and to explore the relation between clinical features and pathological findings. We report here the findings in the neonatal deaths.

The specific aims of this paper are to:

- review the epidemiology (sociodemographic, antenatal, and perinatal factors) of the early neonatal deaths overall

and to compare infants who displayed signs of birth asphyxia with those who did not;

- investigate the neuropathological status in those infants in whom a post mortem was authorised, and to determine whether lesions could be of prenatal origin;
- determine if infants who have pre-existing brain damage are, when born alive, more likely to be born in an asphyxiated condition;
- compare the antepartum and intrapartum course of early neonatal deaths of infants born with birth asphyxia with and without pre-existing damage.

METHODS

Study setting and patients

The Scottish perinatal neuropathology study was a prospective observational and experimental study involving all 22 delivery units within Scotland. Patients were recruited during a two year period for each centre. The study started in January 1996, and recruitment of cases was completed by January 1999. The base study considered all perinatal deaths of infants who were ≥ 24 weeks gestation at birth and ≤ 7 days at time of death delivered in Scotland over the two year period. This paper concerns the epidemiology and neuropathology of the liveborn subset of the study cohort. The stillborn infants presented somewhat different features and will be reported on separately.

Infants with central nervous system or cardiac malformations, major chromosomal abnormalities, or central nervous system infection were excluded because it was felt that the neuropathological changes associated with such conditions might interfere with the interpretation of any changes superimposed by perinatal insult.

Figure 1 lists how the cohort of 692 qualifying perinatal deaths was reduced by various exclusions through the 221 liveborn infants to the 70 infants from whom the brain was available for examination in this study. These 70 infants were classified according to whether they displayed birth asphyxia (BA group) or not (noBA group). Analysis of those who died three days or less after the onset of labour allowed identification of pathological features likely to have predated labour and birth. Placentas were available for histological examination from 41 of the 70 infants.

Ethical and consent procedures

Before the start of the study, each delivery unit obtained approval from their local research ethics committee to approach appropriate parents. As different units received ethical permission at slightly varying times, the spread of data collection was three years, although it was two years for each individual centre. Cases were enrolled at the time of post mortem request by the clinician responsible for the care of the infant during life. A detailed clinical dataset was collected on all infants regardless of enrolment status. The purpose of the study was explained to parents. Signed consent was obtained for autopsy, and on a separate consent form, if authorised also for extended neuropathological research studies on the brain.

Clinical details

For each case a detailed questionnaire was completed by specially trained midwives or other local staff who recorded a battery of clinical information and the results of investigations relating to each pregnancy, labour, delivery, and neonatal course. This was entered into a central database (SPSS) by the study clinical coordinator (JCB). Information on the intrapartum cardiotocograph (CTG) was recorded if available.

Diagnosis of asphyxia

No test is available to accurately diagnose clinically important intrapartum asphyxia. The CTG is notorious for its poor predictive value.^{40–41} As one of the principle aims of the study was to determine if infants with pre-existing brain damage are predisposed to neurological depression at birth which might be labelled as birth asphyxia, we used fairly broad inclusion criteria.

- An Apgar score at five minutes of ≤ 5.0 ; this is the traditional assessment and it is widely recognised that a low five minute Apgar score has an association, although weak, with both neonatal death and morbidity in surviving infants.⁴²
- A cord or initial blood pH of < 7.1 ; obstetric epidemiology has shown that a scalp pH of less than 7.25 is abnormal and delivery is indicated if less than 7.2.^{40–41} The relation between scalp and cord pH is good with a sensitivity of 93%.⁴³ However, the neonate is rarely difficult to resuscitate unless the cord pH is less than 7.0. We arbitrarily chose an intermediate level (pH < 7.1) as indicating some degree of birth asphyxia in this group of early neonatal deaths. Recognising the limitations, we also used (in the absence of a cord pH) a first blood gas with a pH less than 7.1 to indicate asphyxia.
- The presence of grade 2/3 neonatal encephalopathy. This is widely accepted as having a closer association with significant birth asphyxia and long term neurodevelopmental disability.^{13–44} The grading of encephalopathy used was that of Sarnat and Sarnat.⁴⁵

Because of the diverse clinical circumstances, not all criteria were available for assessment in each case. Infants who displayed at least one of these criteria were classified as showing clinical evidence of birth asphyxia (BA group). If none of these criteria were present, the infant was included in the non-asphyxiated group (noBA).

Pathological examination

Autopsies

Autopsies were conducted in six Scottish centres, and the brain was retained in fixative for later examination. In the south east of Scotland, the fixed brains were examined in the Department of Neuropathology at the Western General Hospital, Edinburgh. Elsewhere they were sampled locally according to a previously agreed protocol. Up to 20 representative paraffin embedded blocks were prepared in each case from all areas of the cerebrum (including temporal hippocampus), and from the basal ganglia and thalami, midbrain, pons, medulla, vermis, and cerebellar hemispheres. These blocks were collected centrally for review and further investigation in Edinburgh. Paraffin sections were stained routinely with haematoxylin and eosin and luxol fast blue/cresyl violet (myelin). Selected sections were investigated immunocytochemically for astrocytic status, using an antibody to glial fibrillary acidic protein and for microglia/macrophages (antibodies to CD68 and MHCII) or stained with Perls Prussian blue stain (haemosiderin). The neuropathological appearances in grey and white matter were assessed independently in all cases by two observers (JEB and BW), who were initially blind to the clinical history. Selected cases were also reviewed by JWK. Recorded neuropathological features included neuronal eosinophilia and karyorrhexis, astrocytic hyperplasia, activated microglia and accumulation of macrophages, haemorrhage (recent and older), vascular responses, and foci of mineralisation and of infarction. The neuropathological features were then correlated with the gestational and postnatal age of the infant and




Figure 1 The Scottish perinatal deaths cohort. FT, full term; RDS, respiratory distress syndrome; PT, preterm < 37 weeks; IVH, intraventricular haemorrhage; BA, birth asphyxia; MAS, meconium aspiration syndrome; noBA, no birth asphyxia; HIE, hypoxic-ischaemic encephalopathy; PND, prenatal brain damage; NNU, neonatal unit; noPND, no prenatal brain damage; CVS, cardiovascular system; CNS, central nervous system.

with the criteria of birth asphyxia, in combination and individually.

A judgment of whether the damage dated from before the onset of labour, and was therefore prenatal, was based in part on the presence of patently mature lesions such as established infarcts, previous haemorrhage, or extensive mineralisation. However, these features were present in the minority of brain damaged infants. More diffuse features such as definite macrophage infiltration/accumulation and/or prominent reactive astrocytic hyperplasia in white matter are thought to develop over a period of more than three days (table 1). We estimated that the presence or absence of prenatal brain damage could only be determined reliably in infants who died at ≤ 3 days of age (n = 59).

Placenta

The placenta, cord, and membranes were examined macroscopically, and cord length, placental measurements, and trimmed weight were recorded. Any abnormality was described. Histological samples were taken to include a cross section of the umbilical cord, one strip of membranes (adjacent to the hole through which the baby was delivered, if identifiable), and two blocks of placenta with both fetal

and maternal surfaces. Blocks and slides from the placenta and adnexa were submitted for central review in Edinburgh (JK). Histological evidence of infection, specifically chorioamnionitis in the extraplacental membranes or choriionic plate and funisitis, were recorded, as was villitis if generalised.

Statistical analysis

Data were recorded in SPSS. Descriptive statistics were used to examine the prevalence of clinical variables. The χ^2 test with Yates correction (or Fisher's exact test where sample size was less than 20) was used to compare categorical variables, and the unpaired *t* test or Mann-Whitney U test to compare the difference in continuous variables. Significance was assumed at *p* < 0.05, but we recognise that a large number of tests were performed, and some positive results at this level may have occurred by chance. As the epidemiology was performed on observational data, we leave the reader to consider the implications at this level rather than apply a correction such as that of Bonferroni. The statistical comparison of the pathology of asphyxiated and non-asphyxiated infants was made using χ^2 tests with Yates' correction.

Table 1 Timing of injury to the central nervous system after cerebral insult

Pathological feature	Timing of onset after injury	References
Neuronal eosinophilia	6–24 hours	60–62
Neuronal karyorrhexis	12–48 hours	61, 63–65
Infarcts—necrosis	3–8 hours	60, 65, 66
Infarcts—cavitation	14–42 days	65–68
White matter gliosis	3–11 days	31, 32, 58, 60, 61, 65, 67, 69, 70
Grey matter gliosis	3–5 days	30, 63, 68, 71
Microglial upregulation	3 hours–3 days	60, 61, 66, 67, 71
Macrophage infiltration	3–7 days	63, 65–68
Fresh haemorrhage	Minutes	67
Haemosiderin deposits	2–3 days	27, 67, 72
Mineralisation	3–14 days	58, 60, 65, 67

RESULTS

Population and study cohort

Of the 692 deaths in the two years of the study, 221 were early neonatal deaths corresponding to the estimated early neonatal death rate of 2.5/1000 live births in Scotland.

Of the 137 deaths analysed (fig 1), 90 were classified as BA and 47 as noBA according to our liberal definition. Table 2 shows how they met the criteria for birth asphyxia. Most infants died from the effects of prematurity, congenital anomalies, or “anoxia”. The causes of death included one case each of GM1 gangliosidosis, laryngeal atresia, and diaphragmatic hernia, all of which may have contributed to the clinical picture of asphyxia. Twenty out of 137 (15%) pregnancies studied were twin (19) or triplet (one). Complications of pregnancy were common, in particular, oligohydramnios (20%), intrauterine growth restriction (14%), premature rupture of membranes (23%), and second or third trimester antepartum haemorrhage (29%). Although abnormal serum screening for α fetoprotein and human chorionadotrophin occurred in 14 pregnancies, in only five of these was amniocentesis carried out. The other six amniocenteses were performed for amnioreduction (five) or at maternal request (one). Of 62 cases of fetal anomaly scan, 21 were abnormal (including multiple abnormalities). Two infants were conceived following induction by ovulation stimulating drugs, and two by in vitro fertilisation. Emergency caesarean section took place in 57 (42%) deliveries, of which 21 were performed before the onset of labour and 36 were intrapartum. Two infants were born by elective caesarean section, one because of a previous caesarean section and the other because it was a twin pregnancy. There was no excess over the expected proportion of early neonatal deaths delivered out of hours (2100–0900 and weekends; 61%). An abnormal infection screen was found in 15 of the total group (group B *Streptococcus* (seven), coliforms (four), *Staphylococcus aureus* (two), others (two)), of which eight were thought to have died from overwhelming sepsis (three group B *Streptococcus*; one group A *Streptococcus*; two coliforms; one *Pseudomonas*, and one unidentified).

Seventy neonates were fully enrolled in this study with their parents agreeing to both an autopsy and the extended brain sampling. Of these 70, 53 were thought to be asphyxiated (BA group; 23 mature (≥ 37 weeks) and 30 preterm (24–36 weeks) infants), and 17 did not appear to be asphyxiated (noBA group; three mature (≥ 37 weeks) and 14 preterm (24–36 weeks) infants) (fig 1). The mature infants lived for between 15 minutes and seven days, with only three surviving for more than three days. The preterm infants lived for between five minutes and 6.8 days, with only eight infants surviving for more than three days. The ratio of

asphyxiated (77%) to non-asphyxiated (23%) was slightly skewed in the group of autopsied infants towards asphyxiated cases when compared with the whole cohort of liveborn infants included in the detailed epidemiological survey (n = 137; 66% asphyxiated, 34% non-asphyxiated).

Clinical comparison of BA and noBA cohorts

Detailed supplementary tables can be found at <http://adc.bmjournals.com/supplemental/>. Briefly, the mothers were comparable for age, weight, height, social class, marital status, parity, and all other factors examined (supplementary table 1). Mothers of infants who were born in an asphyxiated state were less likely to have received steroids during pregnancy (20% v 36%, p = 0.036). Hyperemesis (8% v 23%, p = 0.013), placenta praevia (2% v 11%, p = 0.037), intrauterine growth retardation (10% v 23% p = 0.066), and pyrexia or flu-like illness during pregnancy (6% v 17%, p = 0.061) were less common in the BA cohort (supplementary table 2). Markers of fetal distress (supplementary table 3), such as meconium staining and cardiotocograph (CTG) abnormalities,⁴⁰ were significantly more prevalent in the BA cohort (26% v 11%, p = 0.040; 59% v 33%, p = 0.004). Intrapartum infection, indicated by positive vaginal swabs, maternal pyrexia, increased white cell count, or increased C reactive protein, occurred in 12 cases (*Escherichia coli* and other coliforms, group B *Streptococcus*, and *Staphylococcus aureus*) but was not more common in the BA group. Malpresentation was less common in the BA cohort (30% v 45%, p = 0.087).

The noBA cohort were of younger gestation (29 v 32, p = 0.017), lighter, and had a smaller head circumference (supplementary table 4). The BA cohort, who had lower Apgar scores, required more resuscitation as a result. Eighteen (20%) infants in the BA group were asystolic at birth. Infants in the BA cohort were more likely to die early compared with those in the noBA cohort (10.3 h v 43 h, p = 0.002). Of 137 infants, only 106 were admitted to a neonatal unit. Of the remaining 31 infants, five were born in good condition and died suddenly and unexpectedly: three were found dead in their cots on the postnatal ward after transfer from the labour ward, and two suffered a sudden acute deterioration in the labour ward after a normal delivery. Twenty four infants had severe birth asphyxia,

Table 2 Clinical features of birth asphyxia in 137 early neonatal deaths

Features of asphyxia	Full term	Preterm
Total number of infants	38	99
Single feature only		
Apgar ≤ 5 at 5 min	9	35
Cord pH < 7.1	0	1
1st pH < 7.1	1	8
NNE	1	0
Two features		
Low Apgar and low pH	7	9
Low Apgar and NNE	2	1
Low pH and NNE	0	1
Three features		
Low pH, low Apgar, and NNE	11	4
Total with some indication of asphyxia	31 (82%)	59 (60%)

All infants had a five minute Apgar score. Only 12 full term infants and 11 preterm infants had cord pH measured. An additional 22 full term infants and 35 preterm infants had the pH measured on arrival in the local neonatal unit. 16 full term infants at 12 hours of age were not paralysed, and 14 of these had features of an encephalopathy. 19 preterm infants remained alive and non-paralysed at 12 hours; six had an encephalopathy. NNE, Neonatal encephalopathy.

and, although they had signs of life at or shortly after birth, they could not be resuscitated sufficiently to move them to the neonatal unit. One extremely premature infant (a triplet) was given only compassionate care.

Clinical details of infants admitted to a neonatal unit were often limited because of early death (supplementary table 5). Within the first hour of birth, infants in the BA group had a considerably lower initial arterial blood pH ($6.96 \nu 7.25$, $p < 0.001$). In survivors of more than 12 hours, those in the BA group were more likely to have renal dysfunction ($55\% \nu 24\%$, $p = 0.029$) and to require assisted ventilation for poor respiratory drive ($33\% \nu 3\%$, $p < 0.001$). The noBA cohort, in keeping with their shorter gestation, had a greater incidence of respiratory distress syndrome and were more likely to have received exogenous surfactant ($73\% \nu 40\%$, $p = 0.001$) and to have muscular paralysis ($36\% \nu 14\%$, $p = 0.011$). In infants who survived for longer than 12 hours, abnormal neurology was documented in 83% of the BA cohort compared with 20% of the noBA cohort ($p < 0.001$). Seizures were significantly more common (19%) in the BA than in the no BA group (3%) ($p < 0.027$), but many infants ($n = 30$) were treated prophylactically with anticonvulsants or paralysis, and many ($n = 61$) died before seizures might have been expected. The groups were comparable for other features of systemic dysfunction, in particular coagulopathies, necrotising enterocolitis, cardiovascular instability, and glucose homeostasis.

Neuropathological findings and identification of prenatal brain damage

Eighty eight infants underwent autopsy, and 70 parents authorised the additional samples required for this research study (fig 1). Table 3 shows the prevalence of neuropathological abnormalities in these 70 infants, classified into the BA and noBA groups ($53 \nu 17$) and according to their gestation (mature ν preterm). Table 4 shows similar data for the infants aged 3 days and less. A detailed table of clinicopathological correlation for each infant in the group of 27 with putative prenatal brain damage has been provided for the interested reader (supplementary table 6). In table 4, the BA group has been further subdivided according to whether encephalopathy was one of the features of birth asphyxia.

In both mature and preterm infants, the asphyxiated infants were more likely to show brain damage than the non-asphyxiated, although brain damage was not universally present in asphyxiated infants (tables 3 and 4). Some infants showed evidence of continuing brain damage, with recent events such as neuronal eosinophilia and fresh haemorrhage superimposed on older lesions including established infarcts, macrophage accumulation including cells laden with haemosiderin, extensive micromineralisation, and white matter gliosis. Infants with no evidence of asphyxia at birth (mostly preterm infants) were more likely than asphyxiated infants to appear virtually normal on neuropathological examination, and such changes as were present, including haemorrhage and neuronal eosinophilia, appeared to be recent except in two mature infants who displayed prominent gliosis.

In cases in which brain damage was present, a conclusion as to whether this was likely to be of prenatal origin could be achieved only in infants who died at ≤ 3 days of age. This was based on the presence of abnormalities thought to first appear about three days after brain injury. There is no absolute certainty about the time needed for the different responses to become visible (table 1), but the presence of accumulations of macrophages and/or prominent astrocytic hyperplasia in human white or grey matter is generally assumed to require three days or more. Evidence from the literature for this timing is presented in more detail in

supplementary table 7. It is important to note that, of the 27 infants judged to have suffered prenatal brain damage, only four had survived for more than two days, six had survived one to two days, and all the rest (65%) had survived for less than one day from the onset of labour. On this basis, 26 (57%) of the asphyxiated group had evidence suggesting prenatal brain damage compared with one (8%) of the non-asphyxiated group, a highly significant difference ($p < 0.005$) (table 4).

Table 4 also shows that infants in the BA group who were encephalopathic displayed a particularly high prevalence of brain damage. Nine of 10 infants in this group showed macrophages or gliosis, or both, together with other confirmatory signs of continuing damage such as neuronal karyorrhexis and eosinophilia. Table 4 also highlights the fact that many of the brains of non-encephalopathic asphyxiated infants were apparently undamaged prenatally and that even by the time of death in the postnatal period, 31% of mature and 13% of preterm asphyxiated infants in this subgroup had apparently normal brains. Although the non-asphyxiated infants appeared to be more prone to postnatal or intrapartum damage, this difference was not significant ($p < 0.059$). Unsurprisingly, the preterm infants were more susceptible to damage of recent, and therefore probably, postnatal origin than were mature infants.

Clinical factors associated with prenatal brain damage

A careful comparison was made of the pregnancies leading to the births of infants with features of pre-labour damage (PND group, $n = 27$) compared with those without such damage (noPND group, $n = 32$). Fewer mothers in the PND group received antibiotics in pregnancy ($1 \nu 8$, $p = 0.031$), more had caesarean section ($17 \nu 10$, $p = 0.015$) and emergency caesarean section ($17 \nu 9$, $p = 0.007$) for CTG abnormalities ($18 \nu 8$, $p = 0.005$), and more had meconium present in the amniotic fluid ($11 \nu 3$, $p = 0.005$). The Apgar score was 0 at birth in 33% of the PND group, significantly more than in the noPND group ($9 \nu 2$, $p = 0.008$), and the former group were heavier and more mature ($2526 \nu 1824$ g, $p = 0.033$, and $34.6 \nu 31.2$ weeks gestation, $p = 0.051$ respectively). The PND group were more likely to be ventilated after birth for a poor respiratory drive ($8 \nu 3$, $p = 0.037$), and, although both groups were acidotic, had a more acidic first pH ($6.90 \nu 7.08$, $p = 0.022$). The time to spontaneous respiration was longer ($5 \nu 1$ minute, $p = 0.009$), and the five minute Apgar score was correspondingly less good ($2 \nu 5$, $p = 0.021$). Reflecting the larger birth weight and more mature status, they had a higher first blood pressure ($46 \nu 36$ mm Hg, $p = 0.019$) and were less likely to receive surfactant ($4 \nu 13$, $p = 0.024$). The time to death, however, was similar in the two groups ($12 \nu 7$ hours, $p = 0.42$).

No differences in sociodemographic or pregnancy factors were identified between the encephalopathic and non-encephalopathic asphyxiated groups, but CTG abnormalities were present in 80% of the former group and in only 43% of the latter group ($p < 0.04$).

Prenatal damage and the signs of birth asphyxia

Table 5 shows the pathology of prenatal brain damage related to the criteria we used for birth asphyxia. Although the strongest clinical association with the features of pre-labour damage is the development of a neonatal encephalopathy after a low pH and a poor Apgar score at five minutes, it is of note that, of the 22 infants who had only a low Apgar score and then died and had a post mortem examination, 11 showed brain damage. By this evidence, a low Apgar score was the sole clinical indicator of prenatal damage in three of

Table 3 Histological evidence of brain damage in 70 neonates

Pathological feature	BA group (n = 53; asphyxiated infants)		NoBA group (n = 17; non-asphyxiated infants)	
	Mature (n = 23)	Preterm (n = 30)	Mature (n = 3)	Preterm (n = 14)
Neuronal eosinophilia	14 (61)	9 (30)	2	5 (36)
Neuronal karyorrhexis	11 (48)	8 (27)	0	0 (0)
Grey matter infarcts	1 (4)	3 (10)	0	0 (0)
White matter gliosis	11 (48)	14 (47)	2	0 (0)
Grey matter gliosis	7 (30)	5 (17)	1	0 (0)
Microglial upregulation	9 (39)	14 (47)	1	1 (7)
Macrophages	9 (39)	14 (47)	0	0 (0)
Fresh haemorrhage	11 (48)	19 (63)	0	8 (57)
Haemosiderin deposits	0 (0)	1 (3)	0	0 (0)
Mineral deposits	2 (9)	8 (27)	2	1 (7)

Values in parentheses are percentages.
Mature, ≥37 weeks; preterm, 24–36 weeks.

13 mature infants and in eight of 13 preterm infants. Only 16 mature and 19 preterm infants in the PND group survived to 12 hours and remained non-paralysed; of these, 14 mature and six preterm infants had an encephalopathy. Looked at another way, 14 full term infants had clinical neonatal encephalopathy. Eight of these had a post mortem examination, and all had evidence of prenatal damage. Only six preterm infants had neonatal encephalopathy. Two of these had a post mortem, and only one had evidence of prenatal damage.

The placenta

In 41 cases (59% of those who had a post mortem examination), a placenta was available for examination. In seven cases, there was histological evidence of infection, and in 33 cases there was none. All of the seven infected placentas came from infants delivered prematurely. In two (25 and 27 weeks gestation), the inflammation was focal, and in four it was more generalised (at 24, 24, 30, and 35 weeks gestation). The placenta of an additional baby, born at 41 weeks gestation, showed focal acute deciduitis without inflammation of the placenta, membranes, or cord. Only two of these infants had evidence of prenatal brain damage. Thus there was virtually no concordance of placental and brain pathology.

DISCUSSION

Early neonatal deaths

A major aim of this study was to determine the neuropathology in a geographically defined cohort of early neonatal

deaths and to seek associations with events in the mother’s pregnancy, labour, and delivery, and with the infant’s condition at birth and during the period before death. We excluded infants with chromosomal abnormalities and with abnormalities of the cardiovascular and central nervous systems because these might themselves lead to neuropathological changes. We were able to review 137 cases with carefully documented clinical detail, and 70 with extensive neuropathology.

Birth asphyxia criteria

Although intrapartum hypoxia manifesting as birth asphyxia is uncommon and in decline,⁴⁶ it is still viewed as a potentially preventable cause of death or damage often with expensive medicolegal implications. Yet there is much evidence that neurodevelopmental delay and cerebral palsy are associated with birth asphyxia in only a minority of cases and also that most birth asphyxiated infants do not manifest developmental delay or cerebral palsy.^{47 48} We used broad inclusion criteria for the diagnosis of birth asphyxia to ensure that we missed no cases and were able to evaluate the individual clinical features of asphyxia in relation to neuropathological abnormality. This may have led to the inclusion of infants whose poor condition was due to other factors such as sepsis and/or metabolic disease, including case 10 (supplementary table 6) with gangliosidosis GM1. Two thirds of our cohort of 137 infants were born in a poor condition. We used the clinical finding of depression at birth manifested by Apgar scores or fetal/neonatal acidosis as a marker of an acute intrapartum event leading to birth asphyxia. An assessment for neonatal encephalopathy in

Table 4 Histological evidence of brain damage, including putative prenatal damage, in 59 neonates aged 3 days or less

Pathological feature	Encephalopathic		No encephalopathy		No encephalopathy	
	BA group (n = 10; 17%)		BA group (n = 36; 61%)		BA group (n = 13; 22%)	
	Mature (n = 8)	Preterm (n = 2)	Mature (n = 13)	Preterm (n = 23)	Mature (n = 2)	Preterm (n = 11)
Neuronal eosinophilia	8 (100)	1	5 (38)	4 (17)	0	4 (36)
Neuronal karyorrhexis	8 (100)	1	1 (8)	4 (17)	0	0
Grey matter infarcts	0	0	0	0	0	0
White matter gliosis*	7 (88)	1	4 (31)	9 (39)	1	0
Grey matter gliosis*	5 (63)	1	0	1 (4)	0	0
Microglial upregulation	6 (75)	1	2 (15)	9 (39)	0	1 (9)
Macrophages*	7 (88)	1	1 (8)	8 (35)	0	0
Fresh haemorrhage	4 (50)	0	6 (46)	11 (48)	0	5 (45)
Haemosiderin deposits	0	0	0	0	0	0
Mineral deposits	1 (13)	0	1 (8)	6 (26)	1	1 (9)
*Estimated prenatal brain damage	8 (100)	1	5 (38)	12 (52)	1	0

Values in parentheses are percentages.
Mature, ≥37 weeks; preterm, 24–36 weeks.

Table 5 Features of asphyxia and presence of putative prenatal damage (PND)

Features of asphyxia	Full term			Preterm		
	Total		PND	Total		PND
	Clinical	PM	at PM	Clinical	PM	at PM
Single feature only						
Apgar \leq 5 at 5 min	9	7	3	35	15	8
Cord pH $<$ 7.1	0	0	0	1	0	0
1st pH $<$ 7.1	1	1	1	8	5	2
NNE	1	0	0	0	0	0
Two features						
Apgar and low pH	7	5	1	9	3	2
Apgar and NNE	2	0	0	1	0	0
Low pH and NNE	0	0	0	1	0	0
Three features						
Low pH, low Apgar and NNE	11	8	8	4	2	1
Total	31	21	13	59	25	13

One full term infant with prenatal damage had no asphyxia.
PM, Total of 47 infants who died at 3 days or less of age. NNE, Neonatal encephalopathy.

combination with these markers would have been more specific,²¹ but many of our infants died within hours of delivery, and a record of neurological examination was not always obtained. In addition, the administration of muscle relaxants to a fifth of our population precluded such an assessment. Finally, 70% of our group were preterm and thus unlikely to exhibit the classical signs of neonatal encephalopathy.

Clinical

The epidemiological background of this cohort is similar to other recent studies from the developed world.^{42 49-51} Analysis of the maternal sociodemographic information and the detailed data from the pregnancy did not identify any reliable predictors for birth asphyxia or for neuropathological abnormalities. Significant placenta praevia and hyperemesis were protective against asphyxia in general, possibly because these mothers were more intensively monitored. Even taking neonatal encephalopathy in isolation as a marker for prenatal asphyxia, no differences were identified between encephalopathic and non-encephalopathic asphyxiated infants in the pregnancy or sociodemographic factors monitored in this study. A history of pyrexia or flu-like illness in pregnancy has previously been found to be associated with neonatal encephalopathy.⁵¹ Our series did not show this association, and pyrexia was more common in pregnancies that resulted in non-asphyxiated infants. Intrauterine growth restriction has previously been strongly associated with neonatal encephalopathy^{49 50 52} and affected 14% of our population, although not just the asphyxiated infants.

CTG abnormalities are common and are poorly predictive of fetal acidosis.⁵³ Both CTG abnormalities and meconium staining of liquor were more common in infants with prenatal damage in this study, and CTG abnormalities proved to be the only difference between the encephalopathic BA and non-encephalopathic BA groups (80% v 43%, $p < 0.04$). Randomised trials have shown that, although monitoring of fetal heart rate can reduce the numbers of neonatal seizures, there is no change in the incidence of long term neurological damage,⁵⁴ suggesting that some fetal heart rate abnormalities may reflect prior compromise. Although meconium staining alone has a high false positive rate,⁵⁵ it is associated with increased perinatal mortality and morbidity.⁵⁶ It has been hypothesised that intra-amniotic meconium may cause vasoconstriction of the umbilical vessels⁵⁷ inducing fetal hypoxia-ischaemia. This is difficult to substantiate after delivery.

Neuropathology

About half (51%) of the 137 eligible infants had detailed neuropathological investigation. The range of neuropathological abnormalities resembles those reported in previous studies,^{28-35 58 60 61 63-69 73} although the prevalence of neuronal damage and damage to the grey matter is higher than elsewhere. Judgments about neuropathological abnormalities are more difficult in the preterm than in the term brain. Despite these difficulties, comparison of the asphyxiated infants and those not apparently suffering from birth asphyxia shows clear differences in terms of neuropathological changes. Examination confined to infants who died within three days of the start of labour, and separation of the asphyxiated group into those with and without neonatal encephalopathy, identifies a spectrum of damage. Unsurprisingly, the mature infants who died after displaying neonatal encephalopathy are most likely to show neuropathological changes. All of the brains in this study were carefully examined to determine whether any damage could have occurred before the onset of labour. If it is accepted that features such as focal or diffuse astrocytic hyperplasia and parenchymal macrophage accumulation are cellular reactions that require three days to become established, we may conclude that most infants with features of birth asphyxia had sustained brain damage prenatally, including all eight of the full term encephalopathic group. We are unable to establish the age of the damage, but the background of apparently normal brain development suggests that the insult was sustained not long before the start of labour. It is harder to draw conclusions about the preterm infants in this study, but the absence of neuropathological changes in virtually all the mature and most of the preterm infants who did not display asphyxia is reassuring.

We do not underestimate the difficulty of interpreting these neuropathological findings and attributing the time of onset. Every abnormality has been included in supplementary table 6, whether focal or diffuse, recent or old, but we accept that interpretation is subjective. Establishing the timing is difficult because experimental studies are not possible in human infants, although classic studies were able to relate pathology findings to major clinical incidents.^{60 67} Some experts also comment that the results of animal work may not be directly relevant to the human situation.^{60 67} Immunocytochemical investigation is mandatory to separate reactive astrocytosis from myelination gliosis. A number of the classic studies were conducted before cell specific immunocytochemistry became available, although this is not the case for more recent papers. Earlier papers may not

have always included, or added, the duration of labour as a factor in timing, and interpretation may be hampered by longer survival. It is noted that the infants with a history of encephalopathy had survived for more than one day in most instances. We concede that seizure activity may induce and accelerate some of the changes seen in the brains of such infants, but the presence of diffuse astrocytosis in other infants who had survived very few hours and who died with no evidence of seizure activity reinforces the possibility of prenatal origin. A more secure evidence base for timing neuropathological events awaits the evolution of new markers of cell damage and irreversible cell death. The clinical significance of some of the lesions described such as diffuse astrocytosis, and in particular their contribution to the cause of death, remains uncertain.

Correlation of clinical factors and neuropathology

This study has failed to identify any pointers that would predict the birth of a compromised infant. Abnormal CTG, and meconium staining of liquor were the only predictive factors for birth asphyxia or prenatal brain damage. Previous studies have reported an association between oligohydramnios and prenatal brain damage possibly related to impaired blood flow in the umbilical cord. Abnormal CTG was the only clinical factor differentiating the asphyxiated infants who displayed encephalopathy and neuropathological abnormality from those who did not. Recently the presence of prenatal infection has been linked to brain damage. We found no support for this association.

Implications for surviving infants

It is possible that the neuropathological findings reported here represent the most severe end of a spectrum of perinatal brain damage resulting in a fatal outcome, while infants surviving perinatal asphyxia might show lesser degrees of similar pathology. However, the possibility also exists that dead infants and survivors represent two completely different groups in terms of both causation and pathology. Recent neuroimaging studies of surviving neonates with encephalopathy, with or without seizures, have a bearing on these questions. A large study by Cowan *et al*²⁹ concluded, on the basis of magnetic resonance imaging performed in the first two weeks of life, that brain damage in mature infants with neonatal encephalopathy was most often acute and of perinatal onset particularly in an encephalopathic group without seizures. Very few infants in that study displayed evidence of prenatal brain damage on magnetic resonance imaging. Neuropathological corroboration was achieved in very few cases. In the absence of immunocytochemical investigation of gliosis and brain macrophage accumulation in all deaths, their conclusions about the prevalence of prenatal abnormalities may be an underestimate. We have discussed the difficulty of timing the lesion in our own study, in which conclusions on the presence or absence of prenatal brain damage were confined to infants who died less than three days after the onset of labour and based on neuropathological examination rather than imaging. We suggest that the cerebral insult was probably sustained only shortly before the onset of labour (even possibly precipitating the onset of labour). Evidence of continuing neuronal damage was also present in our series, not dissimilar to the findings of Cowan *et al*, but this was often in addition to the damage identified as occurring before labour within the constraints of current knowledge. It might be expected that brain damage in survivors would be less extensive and severe than in those with a fatal outcome. Whether the brain damage observed in our study represents the result of persisting or repeated insult, or the onset of a potentially reversible cascade accruing from a single insult, is uncertain. A multistep

pathological process might present opportunities for intervention to limit further brain damage.

The fact that a significant proportion of clinically asphyxiated infants display no evidence of brain damage, and that infants who are not asphyxiated at birth often display only recent postnatal damage, offers hope for a good clinical outcome if such infants could be identified and "rescued" by medical intervention. This study shows that the current battery of investigations associated with pregnancy and labour remain blunt instruments in accurately predicting the arrival of an asphyxiated and prenatally brain damaged infant. Future work must address the development of methods for detecting antepartum damage so that optimal management of these vulnerable fetuses can be planned. Further evidence is also required on evolution of cellular reactions in the developing brain. The findings in this study support the notion that the birth of a compromised "asphyxiated" encephalopathic infant is not necessarily the result of a mismanaged labour nor the lack of vigilance in pregnancy.

ACKNOWLEDGEMENTS

This study was funded by the Scottish Home and Health Department and by Wellbeing and was dependent on the collaboration of Scottish paediatric pathologists and neuropathologists: Drs N Alsanjari, D Doyle, I Graham, E S Gray, A G Howatson, S Lang, A M Lutfy, J McCullough, J MacKenzie, K J McKenzie, J L McPhie, E R Nairn, and N M Smith. We are also grateful to the midwives and medical staff in the 22 neonatal and obstetric units throughout Scotland whose enthusiasm and help in enrolling cases and collecting data made this study possible. We are also grateful for assistance with manuscript production by Ms Angela Penman. Finally we are indebted to the families who, despite the turmoil surrounding the deaths of their infants and knowing that the results would not personally help them at their time of grief, allowed these detailed investigations to proceed.

Authors' affiliations

J C Becher, N McIntosh, Department of Child Life and Health, University of Edinburgh, Edinburgh, Scotland, UK
 J E Bell, B Wyatt, Department of Pathology, University of Edinburgh
 J W Keeling, Department of Paediatric Pathology, Royal Hospital for Sick Children, Edinburgh

REFERENCES

- 1 Confidential Enquiry into Stillbirths and Deaths in Infancy (CESDI). 8th Annual Report. Maternal and Child Health Consortium. 2001.
- 2 Sykes GS, Molloy PM, Johnson P, *et al*. Do Apgar scores indicate asphyxia? *Lancet* 1982;494-6.
- 3 Fields LM, Entman SS, Boehm FH. Correlation of the one-minute Apgar score and the pH value of umbilical arterial blood. *South Med J* 1983;76:1477-9.
- 4 Paneth N, Fox HE. The relationship of Apgar score to neurologic handicap: a survey of clinicians. *Obstet Gynecol* 1983;61:547-50.
- 5 Josten BE, Johnson TR, Nelson JP. Umbilical cord blood pH and Apgar scores as an index of neonatal health. *Am J Obstet Gynecol* 1987;157:843-8.
- 6 Ruth VJ, Raivio KO. Perinatal brain damage: predictive value of metabolic acidosis and the Apgar score. *BMJ* 1988;297:24-7.
- 7 Gilstrap LC, Leveno KJ, Burris JS, *et al*. Diagnosis of birth asphyxia on the basis of fetal pH, Apgar score, and newborn cerebral dysfunction. *Am J Obstet Gynecol* 1989;161:825-30.
- 8 Amato M, Gambon RC, Von Muralit G. Accuracy of Apgar score and arterial cord-blood pH in diagnosis of perinatal brain-damage assessed by CK-BB isoenzyme measurement. *J Perinat Med* 1986;14:335-8.
- 9 Johnson JWC, Richards DS, Wagaman RA. The case for routine umbilical blood acid-base studies at delivery. *Am J Obstet Gynecol* 1990;162:621-5.
- 10 McShane M, Maguire S, McClure G, *et al*. Birth asphyxia, encephalopathy and outcome. *Ir Med J* 1987;80:421-2.
- 11 Nijima S, Levene MI. Post-asphyxial encephalopathy in a preterm infant. *Dev Med Child Neurol* 1989;31:395-7.
- 12 Nelson KB, Leviton A. How much of neonatal encephalopathy is due to birth asphyxia? *Am J Dis Child* 1991;145:1325-31.
- 13 Levene MI, Grindulus H, Sands C, *et al*. Comparison of two methods of predicting outcome in perinatal asphyxia. *Lancet* 1986:67-9.
- 14 Blair E, Stanley FJ. When can cerebral palsy be prevented? The generation of causal hypotheses by multivariate analysis of a case-control study. *Pediatr Perinat Epidemiol* 1993;7:272-301.
- 15 Nelson KB. Relationship of intrapartum and delivery room events to long-term neurologic outcome [Review]. *Clin Perinatol* 1989;16:995-1007.

- 16 **Yudkin P**, Johnson A, Clover LM, *et al*. Assessing the contribution of birth asphyxia to cerebral palsy in term singletons. *Pediatr Perinat Epidemiol* 1995;**9**:156–70.
- 17 **Nelson KB**, Leviton A. How much of neonatal encephalopathy is due to birth asphyxia? *Am J Dis Child* 1991;**145**:1325–31.
- 18 **Edwards AD**, Nelson KB. Neonatal encephalopathies: time to reconsider the cause of encephalopathies. *BMJ* 1998;**317**:1537–8.
- 19 **Hall DMB**. Birth asphyxia and cerebral palsy. *BMJ* 1989;**299**:279–82.
- 20 **Bakketeig LS**. Only a minor part of cerebral palsy cases begin in labour. *BMJ* 1999;**319**:1016–17.
- 21 **MacLennan A**. A template for defining a causal relation between acute intrapartum events and cerebral palsy: international consensus statement. *BMJ* 1999;**319**:1054–9.
- 22 **Confidential Enquiry into Stillbirths and Deaths in Infancy (CESDI)**. Annual Report for 1 January–December 1993. Part 1: summary of methods and main results. London: Department of Health, 1995.
- 23 **Confidential Enquiry into Stillbirths and Deaths in Infancy (CESDI)**. 5th Annual Report. Maternal and Child Health Consortium, 1998.
- 24 **Confidential Enquiry into Stillbirths and Deaths in Infancy (CESDI)**. 6th Annual Report. Maternal and Child Health Consortium, 1999.
- 25 **Confidential Enquiry into Stillbirths and Deaths in Infancy (CESDI)**. 7th Annual Report. Maternal and Child Health Consortium, 2000.
- 26 **Rorke LB**. Pathology of perinatal brain injury. New York: Raven Press, 1982.
- 27 **Squier M**, Keeling JW. The incidence of prenatal brain injury. *Neuropathol Appl Neurobiol* 1991;**17**:29–38.
- 28 **Marin-Padilla M**. Developmental neuropathology and the impact of perinatal brain damage. I: haemorrhagic lesions in the neocortex. *J Neuropathol Exp Neurol* 1996;**55**:758–73.
- 29 **Marin-Padilla M**. Developmental neuropathology and the impact of perinatal brain damage. II: white matter lesions in the neocortex. *J Neuropathol Exp Neurol* 1997;**56**:219–35.
- 30 **Marin-Padilla M**. Developmental neuropathology and the impact of perinatal brain damage. III: Grey matter lesions in the neocortex. *J Neuropathol Exp Neurol* 1999;**58**:407–29.
- 31 **Golden JA**, Gilles FH, Rudelli RD, *et al*. Frequency of neuropathological abnormalities in very low birth weight infants. *J Neuropathol Exp Neurol* 1997;**56**:472–8.
- 32 **Gilles FH**, Leviton A, Jeffrey SM, *et al*. Groups of histopathologic abnormalities in brains of very low birth weight infants. *J Neuropathol Exp Neurol* 1998;**57**:1026–34.
- 33 **Skullerud K**, Westre B. Frequency and prognostic significance of germinal matrix hemorrhage, periventricular leucomalacia, and pontosubicular necrosis in preterm neonates. *Acta Neuropathol* 1986;**70**:257–61.
- 34 **Leviton A**, Paneth N. White matter damage in preterm newborns: an epidemiologic perspective. *Early Hum Dev* 1990;**24**:1–22.
- 35 **Paneth N**, Rudelli RD, Monte W, *et al*. White matter necrosis in very low birthweight infants: neuropathologic and ultrasonographic findings in infants surviving six days or longer. *J Pediatr* 1990;**116**:975–84.
- 36 **Gaffney G**, Squier M, Johnson A, *et al*. Clinical associations of prenatal ischaemic white matter injury. *Arch Dis Child Fetal Neonatal Ed* 1994;**70**:101–6.
- 37 **Mito T**, Kamei A, Takashima S, *et al*. Clinicopathological study of pontosubicular necrosis. *Neuropediatrics* 1993;**24**:204–7.
- 38 **Ellis WG**, Goetzman BA, Lindenberg JA. Neuropathologic documentation of prenatal brain damage. *Am J Dis Child* 1988;**142**:858–66.
- 39 **Cowan FM**, Rutherford MA, Groenendaal F, *et al*. Origin and timing of brain lesions in term infants with neonatal encephalopathy. *Lancet* 2003;**361**:736–42.
- 40 **NICE**. The use of electronic fetal monitoring; the use and interpretation of cardiotocography in intrapartum fetal surveillance. NICE, 2001.
- 41 **Royal College of Obstetrics and Gynaecology**. The use of electronic fetal monitoring; the use and interpretation of cardiotocography in intrapartum fetal surveillance. RCOG, 2001.
- 42 **Gaffney G**, Sellers S, Flavell V, *et al*. Case-control study of intrapartum care, cerebral palsy, and perinatal death [see comments]. *BMJ* 1994;**308**:743–50.
- 43 **Brandt-Niebelschutz S**, Saling E. Indications for operative termination of labor on cardiotocography and fetal blood analysis: the reliability of these methods. *J Perinat Med* 1994;**22**:19–27.
- 44 **Yudkin P**, Johnson A, Clover LM, *et al*. Clustering of perinatal markers of birth asphyxia and outcome at age five years. *Br J Obstet Gynaecol* 1994;**101**:774–81.
- 45 **Sarnat HB**, Sarnat MS. Neonatal encephalopathy following fetal distress. *Arch Neurol* 1976;**33**:696–705.
- 46 **Kalter H**. Five decade international trends in the relation of perinatal mortality and congenital malformations: stillbirth and neonatal death compared. *Int J Epidemiol* 1991;**20**:173–9.
- 47 **Blair E**, Stanley FJ. Intrapartum asphyxia: a rare cause of cerebral palsy. *J Pediatr* 1988;**112**:515–19.
- 48 **Nelson KB**. What proportion of cerebral palsy is related to birth asphyxia? [editorial]. *J Pediatr* 1988;**112**:572–4.
- 49 **Adamson SJ**, Alessandri LM, Badawi N, *et al*. Predictors of neonatal encephalopathy in full term infants. *BMJ* 1995;**311**:598–602.
- 50 **Badawi N**, Kurinczuk JJ, Keogh JM, *et al*. Intrapartum risk factors for newborn encephalopathy: the Western Australia case-control study. *BMJ* 1998;**317**:1554–8.
- 51 **Badawi N**, Kurinczuk JJ, Keogh JM, *et al*. Antepartum risk factors for newborn encephalopathy: the Western Australia case-control study. *BMJ* 1998;**317**:1549–53.
- 52 **Gunn AJ**, Gunn TR. Changes in risk factors for hypoxic-ischaemic seizures in term infants. *Aust NZ J Obstet Gynaecol* 1997;**37**:36.
- 53 **Steer PJ**, Eigbe F, Lissauer TJ, *et al*. Interrelationships among abnormal cardiotocograms in labour, meconium staining of the amniotic fluid, arterial cord blood pH and apgar scores. *Obstet Gynecol* 1989;**74**:715–21.
- 54 **MacDonald D**, Grant A, Sheridan-Pereira M, *et al*. The Dublin randomised controlled trial of intrapartum fetal heart rate monitoring. *Am J Obstet Gynecol* 1985;**152**:524–39.
- 55 **Nelson KB**, Grether JK. Potentially asphyxiating conditions and spastic cerebral palsy in infants of normal birth weight. *Am J Obstet Gynecol* 1998;**152**:524–39.
- 56 **Berkus M**, Langer O, Samueloff A, *et al*. Meconium-stained amniotic fluid: increased risk for adverse neonatal outcome. *Obstet Gynecol* 1994;**84**:115–20.
- 57 **Alshuler G**. Some placental considerations related to neurodevelopmental and other disorders. *J Child Neurol* 1993;**8**:78–94.
- 58 **Gilles FH**, Murphy SF. Perinatal telencephalic leucoencephalopathy. *J Neurol Neurosurg Psychiatry* 1969;**32**:404–13.
- 59 **Rorke LB**. Anatomical features of the developing brain implicated in pathogenesis of hypoxic-ischaemic injury. *Brain Pathol* 1992;**2**:211–21.
- 60 **Norman MG**. Perinatal brain damage. *Perspect Pediatr Pathol* 1978;**4**:41–92.
- 61 **Low JA**, Robertson DM, Simpson LL. Temporal relationships of neuropathologic conditions caused by perinatal asphyxia. *Am J Obstet Gynecol* 1989;**160**:608–14.
- 62 **Graeber MB**, Blakemore WF, Kreutzberg GW. Cellular pathology of the central nervous system. In: Graham DI, Lantos PL, eds. *Greenfield's neuropathology*. London: Arnold, 2002:123–9.
- 63 **Friede RL**. Ponto-subicular lesions in perinatal anoxia. *Arch Pathol* 1972;**94**:343–54.
- 64 **Wigglesworth JS**, Bridger JE. Neuropathological clues to the timing of early brain lesions. In: Lou HC, Griesen G, Larsen JF, eds. *Brain lesions in the newborn. Alfred Benzon Symposium 37*. Copenhagen: Munksgaard, 1994:165–73.
- 65 **Squier MV**. Acquired diseases of the nervous system. In: Keeling JW, ed. *Fetal and neonatal pathology*. London: Springer Verlag, 2001:633–58.
- 66 **Banker BQ**. The neuropathological effects of anoxia and hypoglycemia in the newborn. *Dev Med Child Neurol* 1967;**9**:544–50.
- 67 **Ellis WG**, Goetzman BW, Lindenberg JA. Neuropathologic documentation of prenatal brain damage. *Am J Dis Child* 1988;**142**:858–66.
- 68 **Kinney HC**, Armstrong DD. Perinatal neuropathology. In: Graham DI, Lantos PL, eds. *Greenfield's neuropathology*. London: Arnold, 2002:519–606.
- 69 **Roessmann U**, Gambetti P. Pathological reaction of astrocytes in perinatal brain injury. Immunohistochemical study. *Acta Neuropathol* 1986;**70**:302–7.
- 70 **Norenberg MD**. Astrocyte responses to CNS injury. *J Neuropathol Exp Neurol* 1994;**53**:213–20.
- 71 **Del Bigio MR**, Becker LE. Microglial aggregation in the dentate gyrus: a marker of mild hypoxic-ischaemic brain insult in human infants. *Neuropathol Appl Neurobiol* 1994;**20**:144–51.
- 72 **Vanezis P**. Bruising: concepts of ageing and interpretation. In: Rutty GN, ed. *Essentials of autopsy practice*. London: Springer-Verlag, 2001:221–40.
- 73 **Norman MG**. Antenatal neuronal loss and gliosis of the reticular formation, thalamus, and hypothalamus. A report of three cases. *Neurology* 1972;**22**:910–16.

PostScript

LETTERS

Pain relief during common neonatal procedures: a survey

We conducted a survey of neonatal pain relief practices for common procedures across the United Kingdom as a baseline for improving our own practice, and we here present the results.

We sent a questionnaire to all Scottish hospitals, and units from the rest of the United Kingdom if they had 40 or more maternity beds ($n = 96$);¹ the response rate was 85%. Analgesia was used in 82% of units for elective intubation, the commonest agent used being morphine (79%), followed by fentanyl. Analgesia was also used in 11% of the units for intravenous cannulation and in 10% for heelpricks. The analgesia most commonly used for cannulation was sucrose or dextrose. Some 5% of units stated that they used morphine for radial arterial lines but these infants were already ventilated and receiving morphine.

These data appear to give a snapshot of current practice, but we cannot know how far unit guidelines translated into the actual experience of the babies. As pain in the neonatal period has immediate and long term consequences,^{1,2} and preterm infants may be exposed to many painful procedures during their hospital stay, there is some way to go before we can claim that neonates are getting optimum pain control.^{1,2} The wider use of sucrose and topical anaesthetics (if safety concerns can be adequately addressed) seem likely to be the quickest routes to improving the situation.

Acknowledgements

We thank all the neonatal units who participated in this survey.

C Rennix, C M Manjunatha, S E Ibhaneebhor

Correspondence to: Neonatal Unit, Wishaw General Hospital, Wishaw ML2 0DP, Lanarkshire, Scotland, UK; Chikka10@hotmail.com

doi: 10.1136/adc.2003.045351

References

- 1 Foster GL. *Good birth guide*, 1st ed. Perth, Australia: Vermillion, 2002.
- 2 Chiswick ML. Assessment of pain in neonates [commentary]. *Lancet* 2000;355:6-8.

US bioethics fall short of world standards

In their study of motion resistant pulse oximetry in neonates, Sahni *et al*¹ obtained approval from their institutional review board and consent from the parents of the infants involved. Nevertheless, the study fails the most basic principles of bioethics, and this calls into question the value of institutional review boards and points to a yawning chasm between American ethical practices and world ethical standards.

The recognised criteria for ethical experimentation are the Nuremberg Code (1947)² and the Declaration of Helsinki (1964) as amended.³ The Nuremberg Code requires the consent of the subject, which obviously could not be obtained in this case. The Declaration of Helsinki provides for the consent of the legal representatives of minor children in certain limited instances:

“For a research subject who is legally incompetent, physically or mentally incapable of giving consent or is a legally incompetent minor, the investigator must obtain informed consent from the legally authorized representative in accordance with applicable law. These groups should not be included in research unless the research is necessary to promote the health of the population represented and this research cannot instead be performed on legally competent persons.”³

This provision is inapplicable in this instance because this research did not promote the health of the population group represented and because this research easily could have been performed on legally competent adults.

Male neonatal non-therapeutic circumcision violates basic human rights to security of the person and to freedom from torture, inhuman, or degrading procedures. A recent study found that neonatal circumcision fails all ethical tests.⁴ Moreover, the Norwegian Council for Medical Ethics advised the Norwegian Medical Association that the circumcision of boys is not consistent with important principles of medical ethics, has no established medical benefit, and causes pain even with the use of local anaesthesia.⁵ Non-therapeutic circumcision of children violates articles 1, 2, and 20 of the *European Convention on Human Rights and Biomedicine*.⁶

The institutional review board must be more than a rubber stamp to approve whatever is proposed. Clearly, world ethical standards were not considered in this instance. It is time for American bioethics boards and committees to adopt world standards.

J V Geisheker, G Hill

Doctors Opposing Circumcision, Suite 42, 2442 NW Market Street, Seattle Washington 98107, USA; iconbuster@earthlink.net

doi: 10.1136/adc.2003.047373

References

- 1 Sahni R, Gupta A, Ohira-Kist K, *et al*. Motion resistant pulse oximetry in neonates. *Archives of Disease in Childhood Fetal and Neonatal Edition* 2003;88:F505. Available at: <http://adc.bmjournals.com/cgi/content/full/fetalneonatal%3b88/6/F505>.
- 2 The Nuremberg Code. *BMJ* 1996;313:1448, Available at: <http://bmj.bmjournals.com/cgi/content/full/313/7070/1448>.

3 Declaration of Helsinki, Adopted by the 18th World Medical Assembly, Helsinki, Finland, June 1964, and later amended. Available at: <http://www.wma.net/e/policy/b3.htm>.

4 Hodges FM, Svoboda JS, Van Howe RS. Prophylactic interventions on children: balancing human rights with public health. *J Med Ethics* 2002;28:10-16, Available at: <http://jme.bmjournals.com/cgi/content/abstract/28/1/10>.

5 Gulbrandsen P. Rituell omskjæring av gutter. *Tidsskr Nor Lægeforen* 2001;121:2994, Available at: http://www.tidsskriftet.no/pls/lts/pa_lt.visSeksjon?vp_SEKS_ID=419642.

6 European Convention on Human Rights and Biomedicine (1997). Adopted at Oviedo, 4 April 1997. Available at: <http://conventions.coe.int/Treaty/en/Treaties/Html/164.htm>.

Time is of the essence

Accurate time keeping is of great import in all resuscitation settings and none more so than in the medicolegal minefield of perinatology. Recent cases have shown accurate documentation is of vital importance, especially the recording of times. Two studies have delineated considerable discrepancies in clock times in patient care settings^{1,2}; these may exacerbate any medicolegal issues. A prospective observational study assessed clock accuracy in paediatric and neonatal resuscitation areas in a hospital with three sites, separated by up to 18 miles. Senior paediatric cover was provided by one middle grade and one consultant, making accuracy of timing extremely important. The accuracy of the consultants' watches was prospectively assessed without warning.

A total of 39 clock times were taken and compared with the true time as per the speaking clock. In total, the mean (SD) difference was -11 (40) seconds. The labour suite clocks had a mean of +6 (20) seconds. The neonatal unit clocks had a mean of -109 (107) seconds. When paediatric resuscitation areas throughout the Trust were compared, the mean was +18 (43) seconds. The consultants' watches had a mean of +8 (54) seconds.

In the maternity hospital, the labour suite clocks have been changed to "satellite controlled" ones costing five pounds ninety nine pence. This has resulted in their close correlation, with obvious medicolegal benefits. The other clinical areas studied did not have this technology in place, but the results are considerably better than in previous studies. The accuracy of the consultants' watches was exceptional, as no warning was given. This is a reminder to all to document times accurately and evidence enough that resuscitation areas in hospitals should use modern technology and have a centrally controlled time system to avoid needless errors in annotation. We leave you to take your own time to decide.

A Crow, M Davidson, J P McClure

Correspondence to: Paediatric Department, Crosshouse Hospital, Kilmarnock, Scotland, UK; markgdavidson@doctors.org.uk

doi: 10.1136/adc.2003.048512

References

- 1 Wong CM, Stenson BJ, Laing AL. As time goes by. *Scott Med J* 2002;**47**:138–9.
- 2 Lerner EB, Billitier AJ 4th, Adolph JE. Ambulance, fire, and police dispatch times compared with the atomic clock. *Prehosp Emerg Care* 2000;**4**:28–30.

Atrial flutter in preterm babies

Atrial flutter is uncommon in neonates without congenital heart disease or cardiac surgery. It forms about 3% of cardiac arrhythmias in the newborn.¹ Although idiopathic atrial flutter can occur in the fetus,^{2,3} accounting for 30% of fetal arrhythmias in one series,⁴ spontaneous conversion often occurs during birth. I share our experience of two preterm babies who had atrial flutter associated with maternal opiate abuse. There are no previous case reports on this association.

The first case was of a 27 week gestation baby born to a mother with mild cerebral palsy who was abusing drugs such as heroin, crack cocaine, and alcohol and was on a methadone programme during pregnancy. The baby was ventilated from birth for hyaline membrane disease. He had withdrawal symptoms from day 2 in spite of a maintenance infusion of diamorphine, which was then gradually increased. On day 3, he suddenly developed one brief narrow complex tachycardia followed by a similar persistent tachycardia. This was initially diagnosed as supraventricular tachycardia, and he received appropriate treatment with no effect. On review by a cardiologist, atrial flutter was confirmed. Echocardiography ruled out any structural heart disease. The atrial flutter lasted for seven hours. The heart finally reverted to a sinus rhythm with a second dose of digoxin. The baby continued to receive a maintenance dose of digoxin. There was no recurrence of the atrial flutter.

The second case is of a 28 week preterm baby born to a mother who was a heroin addict and was on a methadone programme during the last trimester of pregnancy. The baby developed hyaline membrane disease and was initially managed with head box oxygen and then nasal continuous positive airways pressure. From day 2 he needed ventilation (with diamorphine maintenance). He developed withdrawal symptoms and, later, two episodes of atrial flutter (fig 1).



Figure 1 Electrocardiogram showing classical sawtoothed flutter waves.

Diamorphine was increased to control the withdrawal symptoms. He spontaneously reverted to sinus rhythm and had no further episodes of atrial flutter. No structural heart disease was found on echocardiography.

These episodes of atrial flutter clearly happened in conjunction with other symptoms of opiate withdrawal. Sympathetic excitation is known to occur during opiate withdrawal. We do not know if this predisposes preterm babies, in whom atrial excitation occurs more readily, to this type of arrhythmia. Until we have further case reports, we will not be certain about this association, and the occurrence of this arrhythmia in these cases may be coincidental. However, we know that this type of arrhythmia, if persistent, can be serious, and immediate treatment will be life saving. Hence preterm babies should be monitored closely during opiate withdrawal.

S Sivakumar

Neonatal Unit, Birmingham Women's Hospital,
Edgbaston, Birmingham B15 2TL, UK;
s_siva_kumar@yahoo.com

doi: 10.1136/adc.2003.046474

References

- 1 Southall DP, Johnson AM, Shinebourne EA, et al. Frequency and outcome of disorders of cardiac rhythm and conduction in a population of newborn infants. *Pediatrics* 1981;**68**:58–66.
- 2 Lisowski LA, Verheijen PM, Benatar AA, et al. Atrial flutter in the perinatal age group: diagnosis, management and outcome. *J Am Coll Cardiol* 2000;**35**:771–7.
- 3 Naumburg E, Reisenfeld T, Axelsson O. Fetal tachycardia: intrauterine and postnatal course. *Fetal Diagn Ther* 1997;**12**:205–9.
- 4 Jaeggi E, Fouron JC, Drblik SP. Fetal atrial flutter diagnosis: clinical features, treatment, and outcome. *J Pediatr* 1998;**132**:335–9.

Is mesenteric blood flow compromised during phototherapy in preterm neonates?

We have previously reported that abdominal distension, visible "ropy" bowel loops, and bile stained gastric aspirates (manifestations of ileus) without loose watery stools are more often observed in preterm neonates having conventional phototherapy (CPT) than in those not having this treatment.¹ Reported changes in the mesenteric blood flow as well as peripheral blood flow and cardiac output during CPT indicate that mesenteric ischaemia may occur during CPT in preterm neonates.^{2–4} We hypothesised that mesenteric blood flow may be compromised during CPT in preterm neonates who are not being fed. If our hypothesis was true, mesenteric ischaemia may explain ileus during CPT in preterm neonates.

In a prospective observational study, superior mesenteric artery blood flow (maximum, minimum) velocity and resistive index (RI) were measured by ultrasound pulsed Doppler in 14 consecutive preterm neonates before and 8–12 hours after the start of CPT. At the time of the study, they did not have associated common risk factors for ileus such

as patent ductus arteriosus, indomethacin, sepsis, electrolyte imbalance, and enteral feeds. Their ventilatory/oxygen needs were minimal, and cardiovascular support was not required. The birth weight, gestational age, and postnatal age of the enrolled neonates were 885–1410 g, 27–29 weeks, and 2–4 days respectively. The mean (SD) maximum velocity (V_{MAX}) and RI before and after the start of CPT were not significantly different: V_{MAX}, 0.41 (0.13) v 0.50 (0.11) m/s (p = 0.10); RI, 0.73 (0.08) v 0.70 (0.08) (p = 0.10). Minimum velocity after CPT was, however, significantly increased: 0.06 (0.04) v 0.16 (0.05) m/s (p < 0.001). Ileus developed 4.8 (2.1) days after the initiation of CPT in 8/14 neonates despite the absence of the risk factors studied.

Increased superior mesenteric artery end diastolic blood flow velocity may indicate photorelaxation of the mesenteric vascular smooth muscle during CPT.⁵ CPT per se may be a risk factor for ileus in preterm neonates.

R Kadalraja, S K Patole, R Muller, J S Whitehall

Department of Neonatology, Kirwan Hospital for Women, and School of Public Health and Tropical Medicine, James Cook University, Queensland, Australia

Correspondence to: S Patole, Department of Neonatal Paediatrics, KEM Hospital for Women, University of Western Australia, Perth, Western Australia 6008; skpatole@hotmail.com

doi: 10.1136/adc.2004.057646

References

- 1 Kadalraja R, Thomas E, Patole SK, et al. Is phototherapy a risk factor for ileus in high-risk neonates? Proceedings of Pediatric Academic Societies Annual Meeting, Baltimore May 2001. *Pediatr Res* 2001;120A.
- 2 Pezzati M, Biagiotti R, Vangi V, et al. Changes in mesenteric blood flow response to feeding: conventional versus fiber-optic phototherapy. *Pediatrics* 2000;**105**:350–3.
- 3 Wu PYK, Wong WH, Hodgman JE, et al. Changes in blood flow in the skin and muscle with phototherapy. *Pediatr Res* 1974;**8**:257–62.
- 4 Benders MJ, Van Bel F, Van de Bor M. Cardiac output and ductal reopening during phototherapy in preterm infants. *Acta Paediatr* 1999;**88**:1014–19.
- 5 Furchgott RF. Endothelium-dependent relaxation, endothelium derived relaxation factor and photorelaxation of blood vessels. *Semin Perinatol* 1991;**15**:11–15.

CORRECTION

doi: 10.1136/adc.2002.037606corr1

J C Becher, J E Bell, J W Keeling, et al. The Scottish perinatal neuropathology study: clinicopathological correlation in early neonatal deaths (*Arch Dis Child Fetal Neonatal Ed* 2003;**89**:F399–407). In Table 4, the second row, last column heading was published incorrectly and should read: NoBA group (n = 13; 22%). We apologise for this error.

Supplementary Table 1 Maternal sociodemographic details

	All cases (n=137)	BA (n=90)	noBA (n=47)	Sig
Maternal age at booking*	28(6)	28.2(5.9)	26.8(6.5)	0.236
Weight at booking (kg)*	66.5(13.7)	67.7(14.6)	64.6(11.2)	0.271
Height at booking (cm)*	163(7)	163(8)	163(6)	0.566
Social class 1,2 [#]	58(43)	41(46)	17(36)	0.311
Unemployed [#]	16(12)	8(9)	8(18)	0.159
Married [#]	38(28)	21(23)	17(38)	0.111
Smoker at booking [#]	51(38)	36(40)	15(34)	0.373
Primigravida [#]	50(37)	30(34)	20(42)	0.287
No of pregnancies [^]	1(2)	1(2)	1(2)	0.468
History of infertility, previous neonatal death or miscarriage [^]	40(29)	31(34)	9(19)	0.062
Previous termination of pregnancy [#]	24(18)	15(16)	9(19)	0.717
Previous uterine surgery [#]	33(24)	24(27)	9(19)	0.329

* mean, standard deviation and p value (independent samples t-test)

number, percentage and 2-tailed significance (Chi-square test)

[^] median, interquartile range and asymptotic 2-tailed significance (Mann-Whitney U test)

Supplementary Table 2 Pregnancy details

	All cases (n=137)	BA (n=90)	noBA (n=47)	Sig
Unbooked or booked >15 weeks [#]	18(13)	12(13)	6(13)	0.926
Systolic BP at booking [*]	115(15)	117(16)	112(13)	0.098
Diastolic BP at booking [*]	68(12)	68(13)	68(10)	0.970
Multiple pregnancy [#]	20(15)	12(13)	8(18)	0.562
Assisted conception [#]	4(3)	3(3)	1(2)	1.000
Abnormal serum screening [#]	14(16)	8(13)	6(22)	0.371
Amniocentesis performed [#]	11(8)	6(7)	5(11)	0.509
Abnormal fetal anomaly scan [#]	21(34)	16(37)	5(26)	0.403
Antibiotics in pregnancy [#]	22(16)	12(13)	10(22)	0.229
Steroids in pregnancy [#]	35(26)	18(20)	17(38)	0.039
Oligohydramnios [#]	27(20)	14(16)	13(30)	0.091
Hyperemesis [#]	18(13)	7(8)	11(24)	0.010
Placenta praevia >grade1 [#]	7(5)	2(2)	5(11)	0.045
Pyrexia >38°C/flu-like illness during pregnancy [#]	13(10)	5(6)	8(18)	0.061
Anaemia <11g/dl [#]	14(33)	27(30)	17(39)	0.437
Polyhydramnios [#]	11(8)	10(11)	1(2)	0.097
Intrauterine growth restriction <3 rd centile [#]	19(14)	9(10)	10(23)	0.066
Loss of fetal movements reported [#]	14(10)	8(9)	6(14)	0.554
Premature rupture of membranes [#]	31(23)	21(24)	10(23)	0.808
Intrauterine or urinary tract infection [#]	18(13)	10(11)	8(18)	0.331
Pregnancy-induced hypertension [#]	14(10)	7(8)	7(16)	0.235
Antepartum haemorrhage in 2/3 trimesters [#]	40(29)	22(24)	18(40)	0.090
Any of the above complications of pregnancy [#]	112(82)	72(80)	38(81)	0.905

* mean, standard deviation and p value (independent samples t-test)

number, percentage and 2-tailed significance (Chi-square test)

^ median, interquartile range and asymptotic 2-tailed significance (Mann-Whitney U test)

Supplementary Table 3 Intrapartum details

	All cases (n=137)	BA (n=90)	NoBA (n=47)	Sig
Induction of labour [#]	13(10)	8(10)	5(9)	0.764
No labour [#]	29(21)	21(23)	8(18)	0.391
Forceps/Ventouse delivery [#]	10(7)	7(8)	3(6)	1.000
Emergency caesarean section [#]	57(42)	41(45)	16(36)	0.194
Malpresentation of fetus [#]	48(35)	27(29)	21(47)	0.087
Meconium staining [#]	28(20)	23(26)	4(9)	0.040
CTG abnormality reported [#]	51/99(52)	42/71(59)	9/27(31)	0.004
Cord prolapse [#]	9(7)	7(8)	2(4)	0.718
Documented intrapartum infection [#]	12(9)	7(8)	5(11)	0.751
Pyrexia in labour >38°C [#]	11(8)	7(8)	4(9)	1.000
Bleeding in labour [#]	32(23)	21(23)	11(24)	0.993
1 st stage duration (h) [^]	2.1(6.6)	2.5(6.8)	1.6(6)	0.726
2 nd stage duration (m) [^]	5(28)	4(27)	7(30)	0.472
Time of ruptured membranes (h) [^]	2(14)	3(17)	1(6)	0.314
Ruptured membranes >24h [#]	28(21)	20(23)	8(18)	0.451
General anaesthetic in labour [#]	26(19)	19(21)	7(16)	0.378
Epidural in labour [#]	44(32)	33(37)	11(22)	0.115
Opiates in labour [#]	47(34)	28(30)	19(42)	0.276
Delivery out of hours (21:00-08:59 and weekends) [#]	83(61)	57(64)	26(53)	0.362

* mean, standard deviation and p value (independent samples t-test)

number, percentage and 2-tailed significance (Chi-square test)

^ median, interquartile range and asymptotic 2-tailed significance (Mann-Whitney U test)

Supplementary Table 4 Infant newborn details

	All cases (n=137)	BA (n=90)	noBA (n=47)	Sig
Male sex [#]	84 (61)	56 (61)	28 (62)	0.763
Gestation [*]	31 (6.4)	32 (6.8)	29(5)	0.017
< 37 weeks gestation [#]	99 (72)	59 (44)	40 (89)	0.015
Weight (g) [*]	1779 (1193)	1971(1274)	1411 (849)	0.004
Occipitofrontal circumference (cm) [*]	27.8 (6)	28.9(6)	26.1(4)	0.015
Apgar 0 at 1 minute [#]	18 (13)	18 (20)	0	0.001
1 minute Apgar [*]	3 (3)	1(2)	6(3)	<0.001
5 minute Apgar [*]	5(3)	3(3)	8(1)	<0.001
10 minute Apgar [*]	5(4)	2(3)	9(1)	<0.001
Time to establish regular respirations (m) [^]	1 (5)	4 (14)	0 (1)	<0.001
Cardiac massage required [#]	40 (29)	37 (40)	3(7)	<0.001
Intubation required [#]	113 (83)	81 (88)	32(71)	0.001
Age at death (h) [^]	15 (49.2)	10.7 (47.8)	39.1 (50.9)	0.002
Admitted to SCBU [#]	106 (77)	65 (73)	41 (87)	0.046

* mean, standard deviation and p value (independent samples t-test)

number, percentage and 2-tailed significance (Chi-square test)

[^] median, interquartile range and asymptotic 2-tailed significance (Mann-Whitney U test)

Supplementary Table 5 Infant first week details

	All cases (n=106)	BA (n=65)	noBA (n=41)	Sig
< 37 weeks gestation [#]	80 (76)	42(63)	38 (97)	0.001
Initial pH within 1 st hour [*]	7.12 (0.25)	6.96 (0.23)	7.25 (0.12)	<0.001
Time to reach normal pH (h) [^]	4 (5)	4.75 (4.88)	3 (4.32)	0.117
First mean arterial pressure [*]	38(15)	40(17)	36(10)	0.207
Hematuria in first 24h [#]	32 (30)	19 (77)	13(71)	1.000
Creatinine >120 [#]	23 (22)	17 (26)	6 (15)	0.020
Inotrope required [#]	35 (33)	20 (34)	15 (38)	0.644
Colloid required [#]	82 (77)	51 (85)	31(81)	0.362
Surfactant given [#]	56 (53)	26(40)	30(74)	0.001
Respiratory distress syndrome [#]	46 (43)	20(32)	26(70)	0.001
Ventilated for poor respiratory drive [#]	21 (20)	20 (33)	1(3)	<0.001
Abnormal coagulation [#]	46 (43)	28 (80)	18 (78)	1.000
Abnormal infection screen [#]	15 (14)	9 (20)	6(17)	0.839
Abnormal liver function tests	7/97 (7)	7 (44)	0	0.022
Hypoglycemia <2.6 mmol/l [#]	40 (38)	22 (39)	18 (50)	0.480
Hyperglycemia >8 mmol/l [#]	36 (34)	22 (42)	14(33)	0.759
Necrotising enterocolitis [#]	5 (5)	4 (7)	1 (3)	0.646
Seizures [#]	12 (11)	11 (16)	1 (3)	0.027
Muscle relaxant [#]	22 (21)	8 (13)	14 (38)	0.011
Abnormal neurology (alive >12hours) [#]	28/45 (62)	25/30(93)	3/15(7)	<0.001
Abnormal cranial USS [#]	48/68(71)	31/40 (76)	17/28 (62)	0.135
Age at death (h) [^]	35.6 (60.25)	18 (62.7)	43.1 (52.1)	0.175

* mean, standard deviation and p value (independent samples t-test)

number, percentage and 2-tailed significance (Chi-square test)

^ median, interquartile range and asymptotic 2-tailed significance (Mann-Whitney U test)

Supplementary Table 6 Clinical & neuropathological features in neonatal deaths with putative prenatal brain damage

Case	Gestation	Clinical Features	Estimated Total Hours of Labour plus postnatal survival	Neuropathological Features
1	42	1. Normal 2. Abnormal CTG, ruptured uterus 3. Apgar 4 ⁵ , pH 6.91, Seizures, isoelectric EEG, cerebral oedema on USS, died 32h	32 Hours	Cerebral oedema; neuronal eosinophilia & karyorrhexis; microglial activation & focal macrophage infiltration in white matter.
2	40	1. Normal 2. Meconium, abnormal CTG, ruptured uterus 3. Apgar 0 ¹ 0 ⁵ , pH 6.8, HIE 3, died 15hours	28 Hours	Cerebral oedema; fresh microhaemorrhages, neuronal eosinophilia & karyorrhexis; microglial activation; white matter gliosis & damage.
3	40	1. Normal 2. PROM, meconium, abnormal CTG, suspected infection 3. Apgar 1 ⁵ , pH 6.86, HIE 3, isoelectric EEG, seizures, died 42h	61 Hours	Neuronal eosinophilia & karyorrhexis; white matter damage; microglial activation & macrophage accumulation; focally gliotic white matter.
4	40	1. Normal 2. Abnormal CTG, cord prolapse 3. Apgar 0 ¹ 2 ⁵ , pH 6.79, HIE 3, cerebral oedema on USS, isoelectric EEG, died 14h	21 Hours	Cerebral oedema; neuronal eosinophilia; white matter gliosis & amphophilic globules.
5	39	1. Loss FM 32,39wks, essential HT 2. No labour, meconium, abnormal CTG, fetomaternal bleed 3. Apgar 4 ⁵ , pH 7.06, HIE grade 3, died 14h	14 Hours	Neuronal eosinophilia & karyorrhexis; grey matter gliosis; microglial activation and macrophage accumulation.
6	38	1. Unbooked, severe PIH 2. No labour, abnormal CTG, multiple placental infarctions 3. Apgar 0 ¹ 2 ⁵ , pH 6.57, seizures, HIE grade 3, bilateral echogenicity on USS, died 17h	17 Hours	Cerebral oedema, neuronal eosinophilia & karyorrhexis; white matter gliosis & macrophage accumulation.
7	38	1. Loss fetal movements 38/40		Germinal matrix haemorrhage;

		2. No labour, abnormal CTG, fetomaternal bleed 3. Apgar 0 ¹ 0 ⁵ , pH 6.8, severe HIE, died 43h	43 Hours	neuronal eosinophilia & karyorrhexis; white matter damage & gliosis; microglial activation & macrophage accumulation.
8	37	1. Smoking 2. No Labour, abnormal CTG, uterine rupture 3. Apgar 0 ¹ 0 ⁵ , pH 6.69, no cortical activity on EEG, seizures, died 35h	35 Hours	Cerebral oedema; neuronal eosinophilia & karyorrhexis; gliosis grey matter; microglial activation & macrophage accumulation.
9	36	1. Smoking, multiple pregnancy, APH 30-36wks 2. Green vaginal discharge, complex shoulder presentation, abnormal CTG 3. Apgar 0 ¹ 3 ⁵ , pH 6.9, abnormal neurology, abnormal background EEG, died 45h	53 Hours	Cerebral oedema, neuronal eosinophilia & karyorrhexis; microglial activation & macrophage infiltration; white matter gliosis.
10	40	1. Previous NND 2. Unexpected poor condition 3. Apgar 7 ⁵ , pH 6.94, abnormal tone, poor respiratory drive, died 26h	39 Hours	Gangliosidosis – GM1; white matter gliosis.
11	36	1. Smoking, previous anencephalic infant, oligohydramnios, IUGR 2. Breech, meconium 3. Apgar 0 ⁵ , died <1h	4 Hours	Micromineralisation; white matter damage & macrophage accumulation.
12	35	1. Known duodenal atresia, antenatal steroids, ROM 35wks, IUGR 2. Meconium, abnormal CTG 3. Tracheal atresia, oesophageal-pulmonary fistula, Apgar 4 ⁵ , pH 6.8, died 11h	11 Hours	Cerebral oedema; white matter gliosis & microglial activation.
13	35	1. Previous SB, unbooked, massive fetal ascites on USS 2. No labour, meconium, abnormal CTG 3. Apgar 1 ⁵ , pH 6.81, abnormal neurology, died 8h	8 Hours	Mineralised neurons in basal ganglia; fresh microhaemorrhages; neuronal karyorrhexis; white matter gliosis & macrophages.

14	32	1. Known duodenal atresia, amnioreduction, polyhydramnios, antenatal steroids, severe PIH 2. No labour, abnormal CTG 3. Apgar 0 ¹ 0 ⁵ , pH 6.9, died 1h	1 Hour	Cerebral oedema; micromineralisation; focal white matter damage; white matter gliosis.
15	28	1. Multiple pregnancy, antenatal steroids, hydropic 1 st twin 2. Breech, meconium 3. Apgar 1 ⁵ , died 2h	7 Hours	Microhaemorrhages; basal ganglia micromineralisation; white matter damage & gliosis; focal macrophage accumulation.
16	28	1. Smoking, oligohydramnios, IUGR, antenatal steroids 2. No labour, abnormal CTG, breech 3. Apgar 9 ⁵ , pH7.09, Abnormal neurology, USS showed IVH, PVL, died 70h	70 Hours	Germinal matrix haemorrhage with thrombosed vessels; white matter infarction & gliosis; neuronal eosinophilia & karyorrhexis; macrophage accumulation.
17	27	1. Previous SB 25wks, antenatal steroids, oligohydramnios, IUGR 2. No labour, meconium, abnormal CTG 3. Apgar 4 ⁵ , died <1h	0.5 Hours	Fresh microhaemorrhages; neuronal eosinophilia; focal grey matter gliosis; macrophage accumulation in grey & white matter.
18	25	1. High AFP, severe oligohydramnios, IUGR, APH at 16, 20, 25 wks, ROM 17wks 2. PROM, breech 3. Apgar 1 ⁵ , died 2h	6 Hours	Germinal matrix haemorrhage; neuronal eosinophilia; white matter gliosis; microglial activation & macrophage infiltration.
19	24	1. Unbooked, IUGR 2. Breech 3. Apgar 2 ⁵ , died<1h	2 Hours	Focal macrophage accumulation & gliosis in grey matter.
20	24	1.Oligohydramnios, APH 13/40, ROM 21/40, anemia <9g/dl, suspected infection	13 Hours	Germinal matrix haemorrhages; neuronal

		2. PROM 3. Apgar 8 ⁵ , pH 7.08, IVH, died 12h		eosinophilia; focal white matter damage & microglial activation; grey matter gliosis.
21	40	1. APH 6,29wks 2. Cord haemorrhage 3. Apgar 0 ⁵ , died at 1h	2 Hours	Focal grey matter gliosis; microglial activation; focal macrophage accumulation & white matter damage.
22	39	1. Essential HT 2. Meconium, abnormal CTG 3. Apgar 0 ¹ 0 ⁵ , died<1h	3 Hours	White matter diffuse microglial activation & macrophage accumulation; focal gliosis of white matter.
23	26	1. Smoking, low AFP, oligohydramnios, APH 9/40, grade 3 placenta praevia, ROM 26/40, suspected infection, bicornuate uterus 2. PROM, breech, abnormal CTG 3. Apgar 7 ⁵ , pH 7.12, died 11h	29 Hours	Microhaemorrhages; focal microglial activation & macrophage infiltration of white matter.
24	25	1. Severe maternal varicella, heavily sedated and ventilated, antenatal steroids 2. Breech, delivered unexpectedly in adult ITU 3. Apgar 0 ¹ 4 ⁵ , pH 6.9, seizures, severe PVH, died 51h	51 Hours	Neuronal eosinophilia & karyorrhexis; germinal matrix haemorrhage; focal microglial activation & macrophage accumulation.
25	42	1. Amniocentesis for low AFP, loss FM 36,41/40, PIH 2. Abnormal CTG 3. Good condition, Apgar 10 ⁵ , collapse at 1h, died 5h	5 Hours	Diffuse white matter gliosis; amphophilic globules.
26	40	1. Normal 2. Meconium, abnormal CTG 3. Apgar 1 ⁵ , died<1h	10 Hours	Fresh microhaemorrhages; white matter gliosis & focal white matter damage.
27	40	1. Normal 2. Meconium 3. Unexpected poor condition, Apgar 5 ⁵ , pH 7.04, died 13h	18 Hours	Cerebral oedema; focal white matter gliosis & microglial activation.

Key:

1. pregnancy features
2. labour and delivery features
3. resuscitation and neonatal features

ROM rupture of membranes; PROM ruptured membranes >24hours; HT hypertension; IUGR intrauterine growth retardation; PVH periventricular haemorrhage; APH antepartum haemorrhage; FM fetal movements; SB stillbirth; AFP serum alpha fetoprotein

Supplementary Table 7 Timing of CNS Injury after Cerebral Insult

Pathological Features	Timing of Onset after Injury	References
Neuronal Eosinophilia	6 – 24 hours	<p>Norman (1978) described eosinophilia in differentiated neurons 24 -36 hours after hypoxic insult (such as delay in establishing respiration) in a classic study of perinatal brain damage. In quoting the results of animal studies, she points out that these may not be applicable directly to human infants.</p> <p>Low et al (1989) suggests that eosinophilia requires at least 18 hours after a documented insult (results in 16 of 120 perinatal deaths).</p> <p>A major current text of neuropathology (Graeber et al, in Greenfield's Neuropathology 2002) quotes a period of "more than 6" hours after insult with reference to observations in the rat.</p>
Neuronal Karyorrhexis	12 – 48 hours	<p>Friede (1972) described neuronal karyorrhexis in mature infants surviving at least 22 hours after an insult.</p> <p>Low et al (1989) suggested that nuclear pyknosis requires 18 hours to become visible. This study documents the duration of labour as well as postnatal survival but suggests that histological changes do not indicate precisely the timing of the insult.</p> <p>Wigglesworth & Bridger (1994) suggested this change required a time lapse of more than 24 hours from their own studies of perinatal deaths.</p> <p>Squier (2001) suggested a time interval of 12-48 hours from her own experience and a survey of the literature.</p>
Infarcts – Necrosis	3 – 8 hours	<p>Banker (1967) quotes 3 hours as the period needed for coagulation necrosis to become evident but a further 9 hours required for commencing cellular reactions.</p> <p>Norman (1978) describes smudgy eosinophilic coagulation necrosis with axonal balls and pyknotic glial nuclei arising 3-8 hours after a cerebral insult; most likely due to a failure of tissue perfusion.</p> <p>Squier (2001) describes coagulation necrosis and retraction balls occurring within 3 hours of insult, quoting other studies and her own observations.</p>
Infarcts – Cavitation	14 – 42 days	<p>Banker (1967) described cysts appearing in areas of damaged white matter two weeks after the insult.</p> <p>Ellis et al (1988) drew on their own experience to conclude that 14 days was required.</p> <p>Squier (2001) quotes the literature and her own experience in suggesting a period of 14 – 42 days.</p> <p>Kinney & Armstrong (2002) described a delay of "a few weeks".</p>
Reactive Gliosis (white matter)	3 – 11 days	<p>Gilles & Murphy (1969) thought that hypertrophic astrocytes required 3 days to appear and were able to attribute brain damage to the prenatal period on this basis, provided that the astrocytosis was accompanied by other evidence of white matter damage such as glial pyknosis.</p> <p>Roessman & Gambetti (1986) thought that 4 days were required (for the appearance of hypertrophied astrocytes identified categorically by GFAP immunocytochemistry; sometimes present in isolation but is taken as evidence of brain damage).</p> <p>Ellis et al (1988) examined very carefully a series of infants</p>

		<p>in whom astrocytosis was graded as early, established or late. Subtle early astrocytosis could be detected by one day after insult whereas enlarged and hypertrophic astrocytes required 3-5 days.</p> <p>Low et al (1989) indicated that 3 – 5 days were required.</p> <p>Norenberg (1994) concentrated on astrocytic reactions and thought these were maximal by 4 days after injury in humans compared with 24 hours in a rat model – he emphasised that the two situations were not comparable in that the human brain was likely to suffer more widespread damage and required to recover before mounting cellular reactions.</p> <p>The consensus is that reactive astrocytosis requires around 3 days and on this basis, damage has been ascribed in several series to the prenatal period although Norman (1978) and Squier (2001) both draw attention to the potential confusion between reactive astrocytosis and “myelination” gliosis. Stress is generally laid on features accompanying gliosis eg amphophilic globules in and around the walls of vessels or macrophage infiltration in order to attribute significance to the reactive gliosis (Golden et al 1997, Gilles et al 1998).</p>
Reactive Gliosis (grey matter)	3 – 5 days	<p>Friede (1972) described a glial response occurring in pontosubicular necrosis 3-5 days after the insult. If infants with pontosubicular necrosis survived only 1-2 days, the grey matter glial response was very slight.</p> <p>Del Bigio & Becker (1994) described the glial response in damaged dentate gyrus “lagging behind” a microglial response which itself required 1-4 days.</p> <p>Marin-Padilla (1999) emphasised that grey matter damage was repaired by gliosis, unlike white matter which cavitates, but found this phenomenon only in comparatively longer survivors (weeks and months after the insult).</p> <p>Kinney & Armstrong (2002) suggested that 3-5 days were required for grey matter gliosis to follow on neuronal necrosis in both preterm and mature infants.</p>
Microglial Upregulation	3 hours – 3 days	<p>Banker (1967) observed microglial infiltrate about 12 hours after the onset of coagulative necrosis.</p> <p>Norman (1978) found rod cells in necrotic white matter foci only after 2-3 days following an anoxic episode.</p> <p>Ellis et al (1988), quoting animal studies compared with their own human studies, observed microglia 1-3 days after an insult.</p> <p>Low et al (1989) observed rod cells 36-72 hours after a hypoxic episode.</p> <p>Del Bigio & Becker (1994) observed an increased number of rod cells only when survival exceeded 4 days following an insult which had produced grey matter infarcts. This study quotes results in animal experiments where microglial upregulation was observed to commence approximately 1 day after the insult, becoming maximal at 4 days.</p>
Macrophage Infiltration	3 – 7 days	<p>Banker (1967) observed macrophages only 1 week after a documented insult.</p> <p>Friede (1972) observed macrophage responses (foamy cells) in relation to pontosubicular damage 4-5 days after the onset of presumed clinical insult.</p> <p>Ellis et al (1988) documented early, middle and late stages from their own observations of macrophage formation and related this to animal studies.</p>

		<p>Squier (2001) observed macrophages only 4-5 days after injury.</p> <p>Kinney & Armstrong (2002) suggested 3-5 days were required for a macrophage response in relation to pontosubicular necrosis, presumably based on their own experience.</p>
Fresh Haemorrhage	Minutes	<p>Fresh haemorrhage is an acute lesion which is too recent to assist in criteria for prenatal damage (Ellis et al 1988) but which can follow an episode of infarction in white or grey matter or a systemic consumption coagulopathy.</p>
Haemosiderin Deposits	2-3 days	<p>Ellis et al (1988) found that haemosiderin staining macrophages accompanied the appearance of macrophages within and around haemorrhagic lesions and that this change required 3 days to appear.</p> <p>Squier & Keeling (1991) suggest that the presence of pigmented macrophages around haemorrhages requires at least 2 days.</p> <p>Vanesis (2001) states that haemosiderin containing macrophages are found within the brain 3-4 days after injury and this is later than in some other tissues. Vanesis emphasises that no reliance should be placed on animal studies in this regard since there is very considerable interspecies variation.</p>
Mineralisation	3 – 14 days	<p>Norman (1978) believed that ferrugination of neurons could occur 3 days post insult. However other authors have all assumed a longer interval.</p> <p>Ellis et al (1988) observed that a period of 14 days was required for the appearance of perivascular mineralising foci in the kitten model. These foci, and amphophilic globules, may be found in association with white matter gliosis (Gilles & Murphy, 1969) and have been used to date the onset of damage in the prenatal period.</p> <p>Ellis et al (1988) thought that neuronal ferrugination required more than 14 days.</p> <p>Squier (2001) thought that mineralisation required 8-14 days. Amphophilic globules in and around the walls of small vessels may represent leaked plasma proteins.</p>

