

Corrections

doi:10.1136/adc.2011.214239

Sie S D, Wennink J M B, van Driel J J, *et al.* Maternal use of SSRIs, SNRIs and NaSSAs: practical recommendations during pregnancy and lactation. This paper was published in print with an incorrect doi. The print citation is *Arch Dis Child Fetal Neonatal Ed* 2012;97:F472–6.

Appendix A. Finnegan Score

System	Symptoms and signs	S c o r e	Date			
			Day	Evening	Night	comments
Central Nervous System Disturbances	High pitched cry	2				
	Continuous high pitched cry	3				
	Sleeps < 1 hour after feeding	3				
	Sleeps > 2 hours after feeding	2				
	Sleeps > 3 hours after feeding	1				
	Hyperactive Moro reflex	2				
	Markedly hyperactive Moro reflex	3				
	Mild tremors when disturbed	1				
	Moderate-severe tremors when disturbed	2				
	Mild tremors when undisturbed	3				
	Moderate-severe tremors when undisturbed	4				
	Increased muscle tone	2				
	Excoriation	1				
	Myoclonic jerks	3				
	Convulsions	5				
Metabolic/Vasomotor/Respiratory Disturbances	Sweating	1				
	Hyperthermia 37.2- 38.5°C	1				
	Hyperthermia > 38.5°C	2				
	Frequent yawning	1				
	Mottling	1				
	Nasal stuffiness	1				
	Sneezing (>3 times)	1				
	Nasal flaring	2				
	Respiratory rate > 60/min	1				
	Respiratory rate > 60/min with retractions	2				
Gastrointestinal Disturbances	Excessive suckling	1				
	Poor feeding	2				
	Regurgitation	2				
	Projectile vomiting	3				
	Loose stools	2				
	Watery stools	3				
	Total score					