



OPEN ACCESS

Neonatal hyperglycaemia is associated with worse neurodevelopmental outcomes in extremely preterm infants

Itay Zamir ¹, Elisabeth Stoltz Sjöström,² Fredrik Ahlsson,³ Ingrid Hansen-Pupp,⁴ Fredrik Serenius,³ Magnus Domellöf ¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/fetalneonatal-2020-319926>).

¹Department of Clinical Sciences, Pediatrics, Umeå University, Umeå, Sweden

²Department of Food, Nutrition and Culinary Science, Umeå University, Umeå, Sweden

³Department of Women's and Children's Health, Pediatrics, Uppsala University, Uppsala, Sweden

⁴Department of Clinical Sciences Lund, Pediatrics, Lund University, Skåne University Hospital, Lund, Sweden

Correspondence to

Dr Itay Zamir, Department of Clinical Sciences, Pediatrics, Umeå University, Umeå, Sweden; itay.zamir@umu.se

Received 11 June 2020

Revised 13 December 2020

Accepted 15 December 2020



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

To cite: Zamir I, Stoltz Sjöström E, Ahlsson F, et al. *Arch Dis Child Fetal Neonatal Ed* Epub ahead of print: [please include Day Month Year]. doi:10.1136/archdischild-2020-319926

ABSTRACT

Objective To assess the associations between neonatal hyperglycaemia and insulin treatment, versus long-term neurodevelopmental outcomes in children born extremely preterm.

Design and setting Observational national cohort study (Extremely Preterm Infants in Sweden Study) using prospectively and retrospectively collected data. Neurodevelopmental assessment was performed at 6.5 years of age.

Patients 533 infants born <27 gestational weeks during 2004–2007; 436 survivors were assessed at 6.5 years.

Outcome measures Neurodevelopmental disability (NDD), survival without moderate to severe NDD, Wechsler Intelligence Scale for Children IV Full scale intelligence quotient (WISC-IV FSIQ) and Movement Assessment Battery for Children 2 (MABC-2) total score.

Results Duration of neonatal hyperglycaemia >8 mmol/L was associated with WISC-IV scores—for each day with hyperglycaemia there was a decrease of 0.33 points (95% CI 0.03 to 0.62) in FSIQ. Neonatal hyperglycaemia >8 mmol/L occurring on 3 consecutive days was associated with lower MABC-2 scores (adjusted mean difference: –4.90; 95% CI –8.90 to –0.89). For each day with hyperglycaemia >8 mmol/L, there was a decrease of 0.55 points (95% CI 0.17 to 0.93) in MABC-2 total score. Insulin treatment was not associated with any of the outcome measures.

Conclusion Neonatal hyperglycaemia >8 mmol/L was associated with lower intelligence scores and worse motor outcomes at 6.5 years of age. Insulin treatment was not associated with either worsened or improved neurodevelopmental outcomes. Randomised controlled trials are needed to clarify the role of insulin in treating hyperglycaemia in extremely preterm infants.

INTRODUCTION

Children and adolescents born extremely preterm (EPT) are at considerable risk of long-term health problems and often have special educational needs.^{1,2} Approximately 25% of children born EPT have moderate to severe disability at 2.5 years of age,³ and a third have motor impairments at 6.5–15 years of age.^{4,5}

Neonatal hyperglycaemia is common in EPT infants, occurring in up to 80% of infants, and it has been associated with increased mortality and white matter reduction at term age.^{6,7} Furthermore, hyperglycaemia was shown to have a negative effect

What is already known on this topic?

- Extremely preterm infants are prone to neonatal hyperglycaemia.
- Neonatal hyperglycaemia has been associated with multiple morbidities and has been suggested to be associated with neurodevelopmental disability.
- Insulin treatment for neonatal hyperglycaemia has shown conflicting results.

What this study adds?

- Both magnitude and duration of neonatal hyperglycaemia were associated with worse motor outcomes at 6.5 years of age and might be associated with lower intelligence scores.
- These associations were observed at glucose concentrations >8 mmol/L.
- Insulin treatment was not associated with either worsened or improved neurodevelopmental outcomes at 6.5 years of age, but further studies are required.

on brain activity during the first 3 days of life in very preterm (VPT) infants.⁸ However, conflicting results have been published so far regarding the association between hyperglycaemia and later neurodevelopmental outcomes in preterm infants.^{9,10}

Insulin treatment is a treatment strategy for neonatal hyperglycaemia, aiming to improve the glucose utilisation of the preterm infant.¹¹ The benefit of insulin treatment has been debated. Retrospective studies have reported conflicting results, from increased survival at postnatal days 28 and 70 in EPT infants⁶ to no effect and even increased mortality and neurodevelopmental disability (NDD) at 2 years of age in children born VPT.^{10,12} Randomised controlled studies have not shown a beneficial effect of insulin treatment on mortality and neurodevelopmental outcomes in different paediatric patient groups, including preterm infants.^{13–16}

This study aimed at assessing the associations between neonatal hyperglycaemia and insulin treatment, and neurodevelopmental outcomes at 6.5 years of age in children born EPT. We hypothesised that neonatal hyperglycaemia would be associated




Figure 1 Study flow chart. EPT, extremely preterm; MABC-2, Movement Assessment Battery for Children 2; NDD, neurodevelopmental disability; WISC-IV, Wechsler Intelligence Scale for Children IV.

with worse neurodevelopmental outcomes, while insulin treatment would be associated with better neurodevelopmental outcomes.

METHODS

Study population

The Extremely Preterm Infants in Sweden Study (EXPRESS) includes all infants who were born at gestational age <27 weeks between 1 April 2004 and 31 March 2007 in Sweden. Cohort characteristics, perinatal data, glycaemic status data, nutritional intakes, survival, morbidity and neurodevelopmental outcomes at age 6.5 years have been previously published.^{4 6 17–19}

Of 707 live-born infants, 602 survived the first 24 hours of life and 533 (88.5%) of those were included in this study (figure 1). Eleven and 23 infants died before reaching 48 and 72 hours of age, respectively. These infants were excluded from analyses involving neonatal hyperglycaemia occurring on 2 and 3 consecutive days (and insulin treatment in infants exposed to such hyperglycaemia), respectively. Ninety-seven children died before the age of 6.5 years (figure 1), and thus, 436 (72.4%) of the children who survived the first 24 hours of life were

included in analyses regarding neurodevelopmental outcomes. It was previously shown that the perinatal and sociodemographic characteristics and disability rates in the follow-up group did not differ from those who were not followed-up.¹⁷

Perinatal and glycaemic status data

Prospectively collected perinatal data was retrieved from the EXPRESS database. Glycaemic status data collection was performed as previously described.⁶ Briefly, all glucose measurements and insulin treatment data available during the first 28 postnatal days were retrospectively collected from hospital records. For each infant, daily highest and lowest glucose concentrations were registered for the first 28 postnatal days, and the respective means were calculated. As no clear definition of hyperglycaemia in this patient population exists, several prospectively defined definitions were used: a highest glucose measurement >8, >10, >12 or >14 mmol/L, occurring at least once or on 2 or 3 consecutive days during the first 28 postnatal days (12 different definitions in total). Insulin treatment was given according to the clinical judgement of the attending physician, since no uniform national guideline for insulin treatment of preterm infants existed in Sweden during the study period.

Neurodevelopmental outcomes

Prospectively collected data from a neurodevelopmental assessment performed at 6.5 years of age at the seven university hospitals in Sweden was retrieved from the EXPRESS database.¹⁷ Wechsler Intelligence Scale for Children IV (WISC-IV; n=355) was used to measure cognitive ability and was administered by psychologists.²⁰ Movement Assessment Battery for Children 2 (MABC-2; n=345), a standardised test assessing gross and fine motor function (manual dexterity, balance, ball skills), was administered by physiotherapists.²¹ MABC-2 total scores ≤75 and ≤68 are equivalent to ≤15th centile (borderline motor impairment) and ≤5th centile (developmental coordination disorder) in Swedish children, respectively.⁴ The Gross Motor Function Classification System (GMFCS) was used to assess functional level in children with cerebral palsy.²² Hearing assessment was based on the child's need of hearing aids. Visual acuity was assessed by ophthalmologists and classified according to modified WHO criteria. Data regarding maternal education was collected using a parental questionnaire.

Children not tested with WISC-IV but who had been physically examined received an NDD category according to clinical examination and medical record review (n=11). Medical record review was also used to assign an NDD category to children who were not physically examined (n=59).

The NDD categories were previously defined in the EXPRESS cohort as follows: no or mild NDD—having Full Scale Intelligence Quotient (FSIQ) score ≥−2 SD, no/mild cognitive disability on clinical examination or medical record review, no cerebral palsy or cerebral palsy (GMFCS 1), and visual acuity of ≥20/63 in the better eye; moderate to severe NDD—having FSIQ score <−2 SD or moderate/severe cognitive disability on clinical examination or medical record review, or cerebral palsy (GMFCS ≥2), or visual acuity of <20/63 in the better eye, or hearing impairment necessitating the use of hearing aid.¹⁷

Statistical analyses

The analyses were done using SPSS Statistical software (V.26.0; IBM, Armonk, New York, USA). Possible associations were assessed between neonatal hyperglycaemia (yes/no), total duration of neonatal hyperglycaemia (in days), insulin treatment in

hyperglycaemic infants (yes/no) and the following outcomes at 6.5 years of age: NDD (no or mild vs moderate to severe), survival without moderate to severe NDD (vs death or survival with moderate to severe NDD), WISC-IV scores (FSIQ scores) and MABC-2 total scores. The associations were evaluated using generalised linear mixed models with an appropriate linear or logistic link function and with a random intercept to account for clustering effect of twins/triplets. Furthermore, in analyses involving insulin treatment, clustering effect of treating hospital was considered in order to account for clinical practice variation regarding insulin treatment.

Possible confounders were identified using directed acyclic graphs. In multivariable models assessing the exposure of hyperglycaemia, the following possible confounders were considered a priori: gestational age at birth, sex, occurrence of intraventricular haemorrhage grade 3–4 and/or periventricular leukomalacia, and educational status of the mother. Furthermore, the following were considered for the first 14 postnatal days: durations of mechanical ventilation and steroid treatments, surgery due to patent ductus arteriosus or necrotising enterocolitis and number of culture-verified sepsis episodes. Models assessing

the exposure of insulin treatment were adjusted for the duration of hyperglycaemia (in days). These models were in a later stage further adjusted for clinical characteristics that differed between infants treated and not-treated with insulin. Furthermore, all models including MABC-2 scores were adjusted for age at follow-up. Children with missing data in a certain variable were excluded from analyses including that specific variable. The significance level was set to $p < 0.05$.

RESULTS

Cohort characteristics are described in table 1. Infants exposed to neonatal hyperglycaemia >10 mmol/L were more often males, had lower gestational age and weight z-score at birth, higher Clinical Risk Index for Babies (CRIB) score and were treated with steroids and mechanical ventilation for longer periods, and insulin-treated infants had lower gestational age at birth and higher prevalence of perinatal morbidities than non-treated infants with hyperglycaemia (table 1). Short-term morbidity and 1-year survival data, as well as follow-up data at 6.5 years of age, including neurodevelopmental outcomes, were

Table 1 Cohort characteristics

Characteristic	The entire cohort (n=533)	No hyperglycaemia (n=164)	Hyperglycaemia (n=369)	P value*	Hyperglycaemic infants		P value
					No insulin (n=298)	Insulin (n=71)	
Perinatal variables							
Gestational age, weeks, mean (\pm SD)	25.3 (1.1)	25.9 (0.8)	25.1 (1.1)	<0.001	25.2 (1.1)	24.5 (0.9)	<0.001
Birth weight z-score, mean (\pm SD)	-0.80 (1.2)	-0.58 (1.1)	-0.90 (1.3)	0.005	-0.89 (1.3)	-0.95 (1.3)	0.717
Male sex, n (%)	288 (54.0)	77 (47.0)	211 (57.2)	0.031	171 (57.4)	40 (56.3)	0.894
Singleton birth, n (%)	417 (78.2)	131 (79.9)	286 (77.5)	0.571	233 (78.2)	53 (74.6)	0.529
CRIB score, mean (\pm SD)	6.5 (3.7)	5.1 (3.7)	7.1 (3.5)	<0.001	6.7 (3.5)	8.7 (3.2)	<0.001
Antenatal steroid treatment, mean no. of doses (\pm SD)	1.5 (0.7)	1.6 (0.7)	1.5 (0.7)	0.592	1.5 (0.7)	1.7 (0.6)	0.045
Neonatal variables							
Days with mechanical ventilation treatment, median (range)†	5 (0–14)	2 (0–14)	7 (0–14)	<0.001	7 (0–14)	11 (0–14)	<0.001
Days with steroid treatment, mean (\pm SD)†	0.3 (1.5)	0.1 (0.9)	0.4 (1.7)	0.007	0.3 (1.4)	1.1 (2.3)	0.008
Sepsis episodes, median (range)†	0 (0–3)	0 (0–2)	0 (0–3)	0.467	0 (0–3)	0 (0–2)	0.959
IVH grade 3–4 and/or PVL, n (%)	95 (17.8)	29 (17.7)	66 (17.9)	1.000	52 (17.4)	14 (19.7)	0.730
Surgically-treated NEC, n (%)†	11 (2.1)	3 (1.8)	8 (2.2)	1.000	4 (1.3)	4 (5.6)	0.048
Surgically-treated PDA, n (%)†	31 (5.8)	5 (3.0)	26 (7.0)	0.074	13 (4.4)	13 (18.3)	<0.001
Daily highest glucose concentration, mmol/L, mean (\pm SD)‡	7.9 (2.3)	5.8 (0.9)	8.7 (2.2)	<0.001	8.2 (1.8)	10.8 (2.6)	<0.001
Daily lowest glucose concentration, mmol/L, mean (\pm SD)‡	5.6 (1.4)	4.7 (1.0)	6.0 (1.4)	<0.001	5.9 (1.3)	6.5 (1.6)	0.003
Days with hyperglycaemia, mean (\pm SD)‡	3.7 (4.9)	0 (0)	5.3 (5.2)	<0.001	3.8 (3.5)	11.3 (6.6)	<0.001
Socioeconomic variables							
Education status of the mother, n (%)§	(n=405)	(n=127)	(n=278)		(n=224)	(n=54)	
≤9 years	38 (9.4)	8 (6.3)	30 (10.8)	0.041¶	23 (10.3)	7 (13.0)	0.588¶
10–11 years	26 (6.4)	12 (9.4)	14 (5.0)		13 (5.8)	1 (1.9)	
12–13 years	141 (34.8)	35 (27.6)	106 (38.1)		87 (38.8)	19 (35.2)	
14–15 years	100 (24.7)	32 (25.2)	68 (24.5)		51 (22.8)	17 (31.5)	
16 years	45 (11.1)	16 (12.6)	29 (10.4)		25 (11.2)	4 (7.4)	
≥17 years	55 (13.6)	24 (18.9)	31 (11.2)		25 (11.2)	6 (11.1)	
Non-Nordic origin of mother, n (%)	88 (20.2) (n=436)	16 (11.6) (n=138)	72 (24.2) (n=298)	0.002	63 (26.1) (n=241)	9 (15.8) (n=57)	0.101

Hyperglycaemia—glucose concentration measurement >10 mmol/L at least once during the first 28 postnatal days; Student's t-test and χ^2 test were used as appropriate.

*P value for comparison between infants with and without neonatal hyperglycaemia.

†During the first 14 postnatal days.

‡During the first 28 postnatal days.

§Percentage based on known values.

¶P value for overall difference between groups with available data.

CRIB, Clinical Risk Index for Babies; IVH, intraventricular haemorrhage; NEC, necrotising enterocolitis; PDA, patent ductus arteriosus; PVL, periventricular leukomalacia.

Table 2 Univariable and multivariable generalised linear mixed models for the associations between neonatal hyperglycaemia (>8 and >10 mmol/L) and moderate to severe neurodevelopmental disability at 6.5 years of age in children born extremely preterm

Hyperglycaemia definition		Moderate to severe NDD					
		No hyperglycaemia	Hyperglycaemia	Crude OR (95% CI)	P value	aOR (95% CI)	P value
>8 mmol/L	at least once	12/56 (21.4%)	132/380 (34.7%)	1.95 (0.99 to 3.84)	0.055	1.27 (0.60 to 2.67)	0.532
	on ≥2 consecutive days	40/160 (25.0%)	104/276 (37.7%)	1.81 (1.17 to 2.81)	0.008	1.16 (0.68 to 1.97)	0.588
	on ≥3 consecutive days	68/252 (27.0%)	76/184 (41.3%)	1.90 (1.26 to 2.87)	0.002	1.36 (0.83 to 2.24)	0.226
	Days with hyperglycaemia (per 1-day increase)	–	–	1.07 (1.04 to 1.10)	<0.001	1.03 (0.99 to 1.08)	0.176
>10 mmol/L	at least once	37/138 (26.8%)	107/298 (35.9%)	1.53 (0.97 to 2.40)	0.066	1.08 (0.63 to 1.87)	0.775
	on ≥2 consecutive days	76/251 (30.3%)	68/185 (36.8%)	1.34 (0.89 to 2.01)	0.165	0.88 (0.53 to 1.48)	0.637
	on ≥3 consecutive days	91/319 (28.5%)	53/117 (45.3%)	2.07 (1.33 to 3.23)	0.001	1.42 (0.82 to 2.45)	0.214
	Days with hyperglycaemia (per 1-day increase)	–	–	1.08 (1.04 to 1.13)	<0.001	1.04 (0.98 to 1.10)	0.178

A random intercept was used to account for the clustering effect of twins/triplets. Adjusted for gestational age at birth, sex, occurrence of intraventricular haemorrhage grade 3–4 and/or periventricular leukomalacia, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, number of culture-verified sepsis episodes during the first 14 postnatal days and educational status of the mother. NDD, neurodevelopmental disability.

previously published.^{4 17 18} Briefly, 33.6% of the children had moderate to severe NDD at 6.5 years of age. Mean (±SD) FSIQ score was 83.4 (14.8), developmental coordination disorder was found in 22.2% and borderline motor impairment was found in further 13.1% of the children. Children tested with WISC-IV and MABC-2 did not differ in their baseline characteristics from children not tested with either of these tests (results not shown), except for a longer duration of mechanical ventilation treatment in non-tested children (7.3 vs 6.0 days for both tests; $p=0.042$ for WISC-IV and 0.015 for MABC-2).

NDD at 6.5 years of age

Neonatal hyperglycaemia and insulin treatment were not significantly associated with moderate to severe NDD at 6.5 years of age in adjusted models, regardless of hyperglycaemia definition or duration (table 2 and online supplemental tables 1 and 2).

Survival without moderate to severe NDD at 6.5 years of age

Neonatal hyperglycaemia and insulin treatment were not significantly associated with survival without moderate to severe NDD at 6.5 years of age in adjusted models, regardless of hyperglycaemia definition or duration (table 3 and online supplemental tables 3 and 4).

WISC-IV scores

Neonatal hyperglycaemia >8 mmol/L occurring on 2 consecutive days (but not according to other hyperglycaemia definitions) was significantly associated with lower WISC-IV scores at 6.5 years of age in adjusted models (table 4 and online supplemental table 5). A longer total duration of exposure to neonatal hyperglycaemia >8 mmol/L (but not according to other hyperglycaemia definitions) was significantly associated with lower WISC-IV scores. Insulin treatment was not significantly associated with WISC-IV scores, regardless of hyperglycaemia definition or duration (except for hyperglycaemia >14 mmol/L occurring on 3 consecutive days; online supplemental table 6).

MABC-2 scores

Neonatal hyperglycaemia >10, >12 and >14 mmol/L, regardless of frequency of occurrence (except for hyperglycaemia >12 mmol/L occurring on 3 consecutive days and hyperglycaemia >14 mmol/L occurring on 2 consecutive days), as well as hyperglycaemia >8 mmol/L occurring on 3 consecutive days, were significantly associated with lower MABC-2 scores at 6.5 years of age in adjusted models (Cohen's $d<0.2$; table 5 and online supplemental table 7). A longer total duration of exposure to neonatal hyperglycaemia was significantly associated

Table 3 Univariable and multivariable generalised linear mixed models for the associations between neonatal hyperglycaemia (>8 and >10 mmol/L) and survival without moderate to severe neurodevelopmental disability at 6.5 years of age in children born extremely preterm

Hyperglycaemia definition		Survival without moderate to severe NDD					
		No hyperglycaemia	Hyperglycaemia	Crude OR (95% CI)	P value	aOR (95% CI)	P value
>8 mmol/L	at least once	44/73 (60.3%)	249/460 (54.1%)	0.78 (0.46 to 1.31)	0.351	0.87 (0.49 to 1.55)	0.628
	on ≥2 consecutive days	120/184 (65.2%)	173/338 (51.2%)	0.56 (0.38 to 0.82)	0.003	1.16 (0.74 to 1.82)	0.517
	on ≥3 consecutive days	185/281 (65.8%)	108/229 (47.2%)	0.46 (0.32 to 0.67)	<0.001	1.54 (0.99 to 2.38)	0.056
	Days with hyperglycaemia (per 1-day increase)	–	–	0.96 (0.93 to 0.99)	0.003	1.00 (0.96 to 1.04)	0.965
>10 mmol/L	at least once	102/164 (62.2%)	191/369 (51.8%)	0.66 (0.45 to 0.96)	0.034	1.13 (0.71 to 1.78)	0.606
	on ≥2 consecutive days	176/291 (60.5%)	117/231 (50.6%)	0.67 (0.47 to 0.96)	0.031	1.03 (0.66 to 1.59)	0.913
	on ≥3 consecutive days	229/362 (63.3%)	64/148 (43.2%)	0.44 (0.30 to 0.66)	<0.001	1.47 (0.91 to 2.39)	0.115
	Days with hyperglycaemia (per 1-day increase)	–	–	0.94 (0.90 to 0.97)	0.001	1.02 (0.97 to 1.07)	0.400

A random intercept was used to account for the clustering effect of twins/triplets. Adjusted for gestational age at birth, sex, occurrence of intraventricular haemorrhage grade 3–4 and/or periventricular leukomalacia, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, number of culture-verified sepsis episodes during the first 14 postnatal days and educational status of the mother. NDD, neurodevelopmental disability.

Table 4 Univariable and multivariable generalised linear mixed models for the associations between neonatal hyperglycaemia (>8 and >10 mmol/L) and Wechsler Intelligence Scale for Children IV (WISC-IV) score at 6.5 years of age in children born extremely preterm

Hyperglycaemia definition		Mean WISC-IV score		Univariable		Multivariable	
		No hyperglycaemia (n)	Hyperglycaemia (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores (95% CI)	P value
>8 mmol/L	at least once	88.82 (45)	83.28 (310)	-5.53 (-9.90 to -1.16)	0.013	-2.58 (-6.79 to 1.64)	0.230
	on ≥2 consecutive days	88.14 (130)	81.6 (225)	-6.51 (-9.49 to -3.53)	<0.001	-3.17 (-6.26 to -0.08)	0.045
	on ≥3 consecutive days	86.42 (208)	80.63 (147)	-5.79 (-8.77 to -2.80)	<0.001	-2.58 (-5.66 to 0.49)	0.099
	Days with hyperglycaemia (per 1-day increase)	–	–	-0.66 (-0.90 to -0.42)	<0.001	-0.33 (-0.62 to -0.03)	0.029
>10 mmol/L	at least once	86.75 (114)	82.71 (241)	-4.04 (-7.23 to -0.84)	0.013	-1.57 (-4.79 to 1.65)	0.337
	on ≥2 consecutive days	85.78 (208)	81.53 (147)	-4.25 (-7.29 to -1.21)	0.006	-1.06 (-4.21 to 2.09)	0.510
	on ≥3 consecutive days	85.65 (263)	79.36 (92)	-6.29 (-9.69 to -2.89)	<0.001	-2.24 (-5.83 to 1.35)	0.220
	Days with hyperglycaemia (per 1-day increase)	–	–	-0.72 (-1.05 to -0.39)	<0.001	-0.31 (-0.69 to 0.08)	0.117

A random intercept was used to account for the clustering effect of twins/triplets. Adjusted for gestational age at birth, sex, occurrence of intraventricular haemorrhage grade 3–4 and/or periventricular leukomalacia, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, number of culture-verified sepsis episodes during the first 14 postnatal days and educational status of the mother.

with lower MABC-2 scores, regardless of hyperglycaemia definition. Insulin treatment was not significantly associated with MABC-2 scores, regardless of hyperglycaemia definition or duration (online supplemental table 8).

DISCUSSION

In this cohort of EPT infants, we observed that neonatal hyperglycaemia and its duration were associated with lower MABC-2 scores but not with NDD at 6.5 years of age, adjusted for multiple clinical factors. Significant associations with worse

motor outcomes were observed when using hyperglycaemia threshold definition as low as >8 mmol/L. Hyperglycaemia was also associated with lower WISC-IV scores. Insulin treatment was not associated with either worsened or improved neurodevelopmental outcomes at 6.5 years of age.

We report a novel finding of an association between neonatal hyperglycaemia and lower MABC-2 scores. Significant associations were found for exposure to hyperglycaemia >10, >12 and >14 mmol/L of almost all frequency definitions, and only for hyperglycaemia >8 mmol/L occurring on 3 consecutive days.

Table 5 Univariable and multivariable generalised linear mixed models for the associations between neonatal hyperglycaemia (>8 and >10 mmol/L) and Motor Assessment Battery for Children 2 (MABC-2) score at 6.5 years of age in children born extremely preterm

Hyperglycaemia definition		Mean MABC-2 score		Univariable		Multivariable	
		No hyperglycaemia (n)	Hyperglycaemia (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores (95% CI)	P value
>8 mmol/L	at least once	73.49 (44)	64.55 (301)	-8.94 (-14.67 to -3.22)	0.002	-4.73 (-10.20 to 0.74)	0.090
	on ≥2 consecutive days	70.27 (129)	62.99 (216)	-7.27 (-11.19 to -3.36)	<0.001	-3.24 (-7.26 to 0.79)	0.114
	on ≥3 consecutive days	68.88 (204)	61.09 (141)	-7.79 (-11.69 to -3.89)	<0.001	-4.90 (-8.90 to -0.89)	0.017
	Days with hyperglycaemia (per 1-day increase)	–	–	-0.84 (-1.15 to -0.53)	<0.001	-0.55 (-0.93 to -0.17)	0.005
>10 mmol/L	at least once	70.91 (114)	63.16 (231)	-7.75 (-11.83 to -3.68)	<0.001	-4.64 (-8.78 to -0.51)	0.029
	on ≥2 consecutive days	69.48 (203)	60.31 (142)	-9.17 (-13.04 to -5.29)	<0.001	-5.87 (-9.94 to -1.79)	0.005
	on ≥3 consecutive days	68.17 (256)	58.48 (89)	-9.69 (-14.09 to -5.29)	<0.001	-6.05 (-10.71 to -1.39)	0.011
	Days with hyperglycaemia (per 1-day increase)	–	–	-1.18 (-1.59 to -0.77)	<0.001	-0.91 (-1.39 to -0.42)	<0.001

A random intercept was used to account for the clustering effect of twins/triplets. Adjusted for gestational age at birth, sex, occurrence of intraventricular haemorrhage grade 3–4 and/or periventricular leukomalacia, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, number of culture-verified sepsis episodes during the first 14 postnatal days, educational status of the mother, and age at follow-up. MABC-2 total scores ≤75 and ≤68 are equivalent to ≤15th centile (borderline motor impairment) and ≤5th centile (developmental coordination disorder) in Swedish children, respectively.

These findings suggest a possible dose–response relationship between glucose concentrations and the outcome. For example, for each additional day with hyperglycaemia >8 mmol/L, there was observed a decrease of MABC-2 score by 0.55 points, while for each additional day with hyperglycaemia >14 mmol/L there was observed a decrease of MABC-2 score by 1.47 points. These findings might be of clinical importance since developmental coordination disorder is associated with poor health outcomes.²³

It was previously shown that neonatal hyperglycaemia was not associated with neonatal illness or NDD at 2 years of age in children born VPT.⁹ However, the cohort in that study had higher mean gestational age compared with our cohort and neonatal hyperglycaemia was defined differently. The Hyperglycaemia In Neonates Trial (HINT) assessed neurocognitive function at 7 years of age in preterm-born children with neonatal hyperglycaemia, randomised to receive either standard or tight glycaemic control.¹⁶ No associations between glucose concentrations and MABC-2 scores were found, yet all infants included in that trial were hyperglycaemic while we compared hyperglycaemic and non-hyperglycaemic infants.

It has been suggested that preterm neonates exhibit both insulin resistance and impaired insulin production.²⁴ Conflicting results have been published regarding the effect of insulin treatment for neonatal hyperglycaemia on neurodevelopmental outcomes. A retrospective study in preterm infants did not observe an association between insulin treatment and neurodevelopmental outcomes at 1 year of age.¹² Infants included in that study had higher gestational age at birth, CRIB score and incidence of perinatal morbidities than our cohort. Another retrospective case–control study in children born VPT observed that infants treated with insulin for neonatal hyperglycaemia (after not responding to glucose intake reduction) had a higher incidence of abnormal neurological development at 2 years of age compared with infants not exposed to hyperglycaemia.¹⁰ In the HINT study, infants who had glucose concentrations in the range 4–6 mmol/L were more likely to survive without NDD and low WISC-IV scores, yet tight glycaemic control was not associated with a difference in NDD, survival without NDD, WISC-IV or MABC-2 scores at 7 years of age.¹⁶ In our study, a slightly different definition for NDD was used (lower FSIQ score, higher visual acuity, MABC-2 not a criteria) and different confounders were adjusted for. Furthermore, infants treated with insulin included in our study were generally sicker than hyperglycaemic infants not treated with insulin. Despite this, no differences were found in the studied outcomes, which might suggest a potential positive effect of insulin treatment. Further adjusting for variables differing between infants treated and not-treated with insulin did not affect the results. However, no definite conclusion regarding such a positive effect can be drawn from the current observational study design and further intervention studies are needed to delineate the long-term effects of insulin treatment.

The association between neonatal hyperglycaemia and lower WISC-IV scores in comparison with non-hyperglycaemic infants is a novel finding. However, we observed this association only for hyperglycaemia >8 mmol/L and not when higher thresholds were used. This might be due to limited statistical power or a chance finding. Nevertheless, the results call for further investigation regarding the association between neonatal hyperglycaemia and intelligence outcomes.

The mechanisms behind the possible effect of hyperglycaemia on neurocognitive functions are unclear. Lower brain weight and more severe brain damage, especially in the hippocampus region, were observed in hyperglycaemic rats.²⁵ Furthermore, neonatal

hyperglycaemia was found to induce oxidative stress, apoptosis and an increase in inflammatory cytokines, with ensuing microglial activation and astrocytosis.²⁶ These changes resulted in long-term changes in synaptogenesis and behaviour in rats. Higher methylation of leucine-rich alpha-2-glycoprotein 1 (a protein associated with neurodevelopment), as well as upregulation of poly (ADP-ribose) polymerase-1 and NF- κ B expression (associated with microglial activation) were associated with higher glucose concentrations.^{27 28} In EPT infants, hyperglycaemia >8.3 mmol/L on the first day of life was associated with white matter reduction at term-equivalent age.⁷ A meta-analysis concluded that neonatal hypoglycaemia was associated with visual-motor impairments, thus implying a possible effect of hyperglycaemia on motor development.²⁹

To the best of our knowledge, our study is the largest study conducted in EPT infants investigating the associations between neonatal hyperglycaemia and neurodevelopmental outcomes in childhood. Other strengths of this study include its prospective design regarding neurodevelopmental outcomes data and the broad consideration of possible confounders. A limitation of the study is its retrospective design regarding glycaemic status data. Hospitals differed significantly in the frequency of glucose sampling; however, we have previously shown that no site-dependent differences in hyperglycaemia prevalence were found in this cohort.⁶ Insulin treatment might have had an impact on the magnitude and duration of hyperglycaemia and vice versa. Insulin treatment might not have had a sufficient effect on glucose concentrations or was introduced late, as reflected by the longer duration of exposure to hyperglycaemia in insulin-treated infants (table 1). However, the effect of insulin treatment on glucose concentrations could not be assessed, and thus, a definite conclusion regarding the effectiveness of insulin treatment cannot be drawn. Furthermore, as this was an observational study, our results represent associations and not necessarily causal effects. Although potential confounders have been considered, residual confounding cannot be excluded. The effect size of hyperglycaemia on neurodevelopmental outcomes was relatively small and this study was not originally designed and powered to measure the associations reported.

In conclusion, neonatal hyperglycaemia >8 mmol/L and longer duration of hyperglycaemia were associated with lower intelligence scores and worse motor outcomes at 6.5 years of age in children born EPT. Insulin treatment in hyperglycaemic infants was not associated with neurodevelopmental outcomes at 6.5 years of age. This observational study highlights the potential detrimental long-term effects of neonatal hyperglycaemia, while the benefit of insulin treatment remains unclear. Further studies, including adequately powered randomised controlled trials, are needed in order to better define neonatal hyperglycaemia in preterm infants, to guide its treatment and clarify its consequences.

Acknowledgements The authors would like to thank the EXPRESS group and all of the staff who provided help in the study process.

Contributors IZ contributed to the study design, the acquisition and analysis of the data, drafting the initial manuscript and subsequent critical revision. FA and IH-P contributed to the acquisition of the data and critically reviewed the manuscript. ESS, FS and MD contributed to the study design, the acquisition of the data and critically reviewed the manuscript. All authors have approved the final manuscript prior to submission and are accountable for the integrity of the study.

Funding The study was supported by research grants from The Swedish Research Council (Vetenskapsrådet; grant nr 2016-02095) and Västerbotten County Council (ALF; grant nr VLL-640871 and Spjutspets grant VLL 599-2018).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study was approved by the regional ethical committee in Lund, Sweden (Dnr 42/2004, 524/2009), and written informed consent was obtained from all parents.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Itay Zamir <http://orcid.org/0000-0001-9086-7991>

Magnus Domellöf <http://orcid.org/0000-0002-0726-7029>

REFERENCES

- Holsti A, Adamsson M, Hägglöf B, *et al.* Chronic conditions and health care needs of adolescents born at 23 to 25 weeks' gestation. *Pediatrics* 2017;139:e20162215.
- Farooqi A, Hägglöf B, Sedin G, *et al.* Chronic conditions, functional limitations, and special health care needs in 10- to 12-year-old children born at 23 to 25 weeks' gestation in the 1990s: a Swedish national prospective follow-up study. *Pediatrics* 2006;118:e1466–77.
- Serenius F, Källén K, Blennow M, *et al.* Neurodevelopmental outcome in extremely preterm infants at 2.5 years after active perinatal care in Sweden. *JAMA* 2013;309:1810–20.
- Bolk J, Farooqi A, Hafström M, *et al.* Developmental coordination disorder and its association with developmental comorbidities at 6.5 years in apparently healthy children born extremely preterm. *JAMA Pediatr* 2018;172:765–74.
- Holsti A, Adamsson M, Serenius F, *et al.* Two-thirds of adolescents who received active perinatal care after extremely preterm birth had mild or no disabilities. *Acta Paediatr* 2016;105:1288–97.
- Zamir I, Tornevi A, Abrahamsson T, *et al.* Hyperglycemia in extremely preterm infants - insulin treatment, mortality and nutrient intakes. *J Pediatr* 2018;200:104–10.
- Alexandrou G, Skiöld B, Karlén J, *et al.* Early hyperglycemia is a risk factor for death and white matter reduction in preterm infants. *Pediatrics* 2010;125:e584–91.
- Schumacher EM, Larsson PG, Pripp AH, *et al.* The effect of blood glucose and pCO₂ on spectral EEG of premature infants during the first three days of life. *Neonatology* 2014;105:297–305.
- Tottman AC, Alsweller JM, Bloomfield FH, *et al.* Relationship between measures of neonatal glycemia, neonatal illness, and 2-year outcomes in very preterm infants. *J Pediatr* 2017;188:115–21.
- Morgan C. The potential risks and benefits of insulin treatment in hyperglycaemic preterm neonates. *Early Hum Dev* 2015;91:655–9.
- Heald A, Abdel-Latif ME, Kent AL. Insulin infusion for hyperglycaemia in very preterm infants appears safe with no effect on morbidity, mortality and long-term neurodevelopmental outcome. *J Matern Fetal Neonatal Med* 2012;25:2415–8.
- van der Lugt NM, Smits-Wintjens VEJ, van Zwieten PHT, *et al.* Short and long term outcome of neonatal hyperglycemia in very preterm infants: a retrospective follow-up study. *BMC Pediatr* 2010;10:52.
- Srinivasan V, Agus MSD. Tight glucose control in critically ill children--a systematic review and meta-analysis. *Pediatr Diabetes* 2014;15:75–83.
- Beardsall K, Vanhaesebrouck S, Ogilvy-Stuart AL, *et al.* Early insulin therapy in very-low-birth-weight infants. *N Engl J Med* 2008;359:1873–84.
- Sadhvani A, Asaro LA, Goldberg C, *et al.* Impact of Tight Glycemic Control on Neurodevelopmental Outcomes at 1 Year of Age for Children with Congenital Heart Disease: A Randomized Controlled Trial. *J Pediatr* 2016;174:193–8.
- Tottman AC, Alsweller JM, Bloomfield FH, *et al.* Long-term outcomes of hyperglycemic preterm infants randomized to tight glycemic control. *J Pediatr* 2018;193:68–75.
- Serenius F, Ewald U, Farooqi A, *et al.* Neurodevelopmental outcomes among extremely preterm infants 6.5 years after active perinatal care in Sweden. *JAMA Pediatr* 2016;170:954–63.
- EXPRESS Group. Incidence of and risk factors for neonatal morbidity after active perinatal care: extremely preterm infants study in Sweden (express). *Acta Paediatr* 2010;99:978–92.
- Stoltz Sjöström E, Öhlund I, Ahlsson F, *et al.* Nutrient intakes independently affect growth in extremely preterm infants: results from a population-based study. *Acta Paediatr* 2013;102:1067–74.
- Wechsler D. Technical and interpretive manual. In: *Wechsler intelligence scale for children*. 4th edn. San Antonio: Psychological Corporation, 2003.
- Henderson SE, Sugden DA, Barnett AL. *Movement assessment battery for children-2: Movement ABC-2: Examiner's manual*. Pearson, 2007.
- Rosenbaum PL, Palisano RJ, Bartlett DJ, *et al.* Development of the gross motor function classification system for cerebral palsy. *Dev Med Child Neurol* 2008;50:249–53.
- Blank R, Smits-Engelsman B, Polatajko H, *et al.* European Academy for Childhood Disability (EACD): recommendations on the definition, diagnosis and intervention of developmental coordination disorder (long version). *Dev Med Child Neurol* 2012;54:54–93.
- Mitanchez-Mokhtari D, Lahlou N, Kieffer F, *et al.* Both relative insulin resistance and defective islet beta-cell processing of proinsulin are responsible for transient hyperglycemia in extremely preterm infants. *Pediatrics* 2004;113:537–41.
- Tayman C, Yis U, Hirfanoglu I, *et al.* Effects of hyperglycemia on the developing brain in newborns. *Pediatr Neurol* 2014;51:239–45.
- Satrom KM, Ennis K, Sweis BM, *et al.* Neonatal hyperglycemia induces CXCL10/CXCR3 signaling and microglial activation and impairs long-term synaptogenesis in the hippocampus and alters behavior in rats. *J Neuroinflammation* 2018;15:82.
- Cartier J, Piyasena C, Sparrow SA, *et al.* Alterations in glucose concentrations affect DNA methylation at Lrg1 in an ex vivo rat cortical slice model of preterm brain injury. *Eur J Neurosci* 2018;47:380–7.
- Gisslen T, Ennis K, Bhandari V, *et al.* Recurrent hypoinsulinemic hyperglycemia in neonatal rats increases PARP-1 and NF- κ B expression and leads to microglial activation in the cerebral cortex. *Pediatr Res* 2015;78:513–9.
- Shah R, Harding J, Brown J, *et al.* Neonatal glycaemia and neurodevelopmental outcomes: a systematic review and meta-analysis. *Neonatology* 2019;115:116–26.

Online supplementary table 1. Univariable and multivariable generalised linear mixed models for the associations between neonatal hyperglycaemia (>12 and >14 mmol/L) and moderate to severe neurodevelopmental disability at 6.5 years of age in children born extremely preterm.

Hyperglycaemia definition	Moderate to severe NDD		Crude OR (95% CI)	P value	aOR (95% CI)	P value	
	No hyperglycaemia	Hyperglycaemia					
>12 mmol/L	at least once	60/213 (28.2%)	84/223 (37.7%)	1.54 (1.02 to 2.31)	0.039	1.03 (0.63 to 1.70)	0.896
	on ≥ 2 consecutive days	97/314 (30.9%)	47/122 (38.5%)	1.40 (0.90 to 2.18)	0.139	0.93 (0.55 to 1.58)	0.796
	on ≥ 3 consecutive days	114/375 (30.4%)	30/61 (49.2%)	2.21 (1.27 to 3.84)	0.005	1.27 (0.66 to 2.44)	0.479
	Days with hyperglycaemia (per 1 day increase)	-	-	1.09 (1.03 to 1.15)	0.002	1.02 (0.95 to 1.10)	0.547
>14 mmol/L	at least once	79/267 (29.6%)	65/169 (38.5%)	1.48 (0.98 to 2.24)	0.061	1.02 (0.62 to 1.70)	0.927
	on ≥ 2 consecutive days	114/362 (31.5%)	30/74 (40.5%)	1.48 (0.88 to 2.50)	0.141	0.83 (0.44 to 1.54)	0.547
	on ≥ 3 consecutive days	127/405 (31.4%)	17/31 (54.8%)	2.64 (1.25 to 5.58)	0.011	1.51 (0.62 to 3.65)	0.361

Hyperglycaemia definition	Moderate to severe NDD		Crude OR (95% CI)	P value	aOR (95% CI)	P value
	No hyperglycaemia	Hyperglycaemia				
Days with hyperglycaemia (per 1 day increase)	-	-	1.12 (1.04 to 1.20)	0.004	1.03 (0.94 to 1.13)	0.496

A random intercept was used to account for the clustering effect of twins/triplets. Adjusted for gestational age at birth, sex, occurrence of intraventricular haemorrhage grade 3-4 and/or periventricular leukomalacia, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, number of culture-verified sepsis episodes during the first 14 postnatal days, and educational status of the mother.

Online supplementary table 2. Univariable and multivariable generalised linear mixed models for the associations between insulin treatment in hyperglycaemic infants and moderate to severe neurodevelopmental disability at 6.5 years of age in children born extremely preterm.

Hyperglycaemia definition	Moderate to severe NDD in hyperglycaemic infants		Crude OR (95% CI)	P value	aOR* (95% CI)	P value	aOR† (95% CI)	P value	
	No insulin	Insulin							
>8 mmol/L	at least once	102/323 (31.6%)	30/57 (52.6%)	2.33 (1.31 to 4.17)	0.004	1.33 (0.65 to 2.73)	0.441	1.17 (0.56 to 2.47)	0.677
	on ≥2 consecutive days	74/219 (33.8%)	30/57 (52.6%)	1.86 (1.01 to 3.43)	0.048	1.15 (0.55 to 2.41)	0.706	1.03 (0.48 to 2.22)	0.937
	on ≥3 consecutive days	48/131 (36.6%)	28/53 (52.8%)	1.73 (0.88 to 3.40)	0.114	1.28 (0.58 to 2.81)	0.540	1.00 (0.43 to 2.32)	0.999
>10 mmol/L	at least once	77/241 (32.0%)	30/57 (52.6%)	2.22 (1.22 to 4.04)	0.009	1.21 (0.57 to 2.56)	0.616	1.10 (0.51 to 2.38)	0.803
	on ≥2 consecutive days	40/130 (30.8%)	28/55 (50.9%)	2.00 (1.02 to 3.94)	0.045	1.08 (0.48 to 2.41)	0.857	1.05 (0.46 to 2.37)	0.912
	on ≥3 consecutive days	27/70 (38.6%)	26/47 (55.3%)	1.97 (0.91 to 4.23)	0.084	1.37 (0.57 to 3.28)	0.479	1.12 (0.44 to 2.86)	0.812
>12 mmol/L	at least once	54/167 (32.3%)	30/56 (53.6%)	2.15 (1.14 to 4.07)	0.019	1.56 (0.72 to 3.39)	0.264	1.45 (0.66 to 3.20)	0.353
	on ≥2 consecutive days	22/72 (30.6%)	25/50 (50.0%)	1.81 (0.80 to 4.05)	0.151	1.11 (0.43 to 2.83)	0.834	1.01 (0.39 to 2.65)	0.983

Hyperglycaemia definition	Moderate to severe NDD in hyperglycaemic infants		Crude OR (95% CI)	P value	aOR* (95% CI)	P value	aOR† (95% CI)	P value
	No insulin	Insulin						
on ≥ 3 consecutive days	11/26 (42.3%)	19/35 (54.3%)	1.61 (0.55 to 4.74)	0.380	1.24 (0.37 to 4.12)	0.724	0.97 (0.25 to 3.70)	0.958
at least once	37/116 (31.9%)	28/53 (52.8%)	2.30 (1.16 to 4.58)	0.018	1.74 (0.80 to 3.80)	0.163	1.54 (0.68 to 3.48)	0.299
>14 mmol/L								
on ≥ 2 consecutive days	13/39 (33.3%)	17/35 (48.6%)	1.78 (0.67 to 4.72)	0.246	1.15 (0.38 to 3.47)	0.809	1.00 (0.30 to 3.33)	0.999
on ≥ 3 consecutive days	6/10 (60.0%)	11/21 (52.4%)	0.69 (0.13 to 3.65)	0.652	0.55 (0.08 to 3.62)	0.521	0.06 (0.002 to 2.50)	0.134

Random intercepts were used to account for the clustering effects of twins/triplets and treating hospital.

* Adjusted for duration of hyperglycaemia (according to respective definition) during the first 28 postnatal days.

† Adjusted for gestational age at birth, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, and duration of hyperglycaemia (according to respective definition) during the first 28 postnatal days.

Online supplementary table 3. Univariable and multivariable generalised linear mixed models for the associations between neonatal hyperglycaemia (>12 and >14 mmol/L) and survival without moderate to severe neurodevelopmental disability at 6.5 years of age in children born extremely preterm.

Hyperglycaemia definition	Survival without moderate to severe NDD		Crude OR (95% CI)	P value	aOR (95% CI)	P value	
	No hyperglycaemia	Hyperglycaemia					
>12 mmol/L	at least once	154/255 (60.4%)	139/278 (50.0%)	0.66 (0.46 to 0.94)	0.022	1.06 (0.69 to 1.62)	0.798
	on ≥ 2 consecutive days	218/368 (59.2%)	75/154 (48.7%)	0.66 (0.45 to 0.97)	0.035	1.04 (0.65 to 1.64)	0.881
	on ≥ 3 consecutive days	262/430 (60.9%)	31/80 (38.8%)	0.41 (0.25 to 0.68)	0.001	1.48 (0.83 to 2.65)	0.183
	Days with hyperglycaemia (per 1 day increase)	-	-	0.92 (0.88 to 0.97)	0.002	1.02 (0.96 to 1.08)	0.586
>14 mmol/L	at least once	189/315 (60.0%)	104/218 (47.7%)	0.61 (0.43 to 0.88)	0.007	1.22 (0.80 to 1.88)	0.357
	on ≥ 2 consecutive days	249/427 (58.3%)	44/95 (46.3%)	0.62 (0.39 to 0.98)	0.042	0.98 (0.57 to 1.67)	0.936
	on ≥ 3 consecutive days	279/469 (59.5%)	14/41 (34.1%)	0.36 (0.18 to 0.71)	0.004	1.56 (0.71 to 3.42)	0.264

Hyperglycaemia definition	Survival without moderate to severe NDD		Crude OR (95% CI)	P value	aOR (95% CI)	P value
	No hyperglycaemia	Hyperglycaemia				
Days with hyperglycaemia (per 1 day increase)	-	-	0.90 (0.83 to 0.96)	0.003	1.04 (0.95 to 1.13)	0.400

A random intercept was used to account for the clustering effect of twins/triplets. Adjusted for gestational age at birth, sex, occurrence of intraventricular haemorrhage grade 3-4 and/or periventricular leukomalacia, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, number of culture-verified sepsis episodes during the first 14 postnatal days, and educational status of the mother.

Online supplementary table 4. Univariable and multivariable generalised linear mixed models for the associations between insulin treatment in hyperglycaemic infants and survival without moderate to severe neurodevelopmental disability at 6.5 years of age in children born extremely preterm.

Hyperglycaemia definition	Survival without moderate to severe NDD in hyperglycaemic infants		Crude OR (95% CI)	P value	aOR* (95% CI)	P value	aOR† (95% CI)	P value	
	No insulin	Insulin							
>8 mmol/L	at least once	222/389 (57.1%)	27/71 (38.0%)	0.47 (0.28 to 0.81)	0.006	1.56 (0.82 to 2.96)	0.177	1.53 (0.79 to 2.97)	0.203
	on ≥2 consecutive days	146/267 (54.7%)	27/71 (38.0%)	0.57 (0.33 to 1.01)	0.054	1.41 (0.73 to 2.73)	0.301	1.37 (0.70 to 2.69)	0.359
	on ≥3 consecutive days	83/168 (49.4%)	25/64 (39.1%)	0.72 (0.39 to 1.34)	0.297	1.29 (0.62 to 2.66)	0.497	1.21 (0.57 to 2.58)	0.621
>10 mmol/L	at least once	164/298 (55.0%)	27/71 (38.0%)	0.54 (0.31 to 0.93)	0.028	1.27 (0.65 to 2.49)	0.478	1.29 (0.65 to 2.55)	0.472
	on ≥2 consecutive days	90/164 (54.9%)	27/68 (39.7%)	0.63 (0.34 to 1.46)	0.135	1.03 (0.51 to 2.09)	0.933	1.10 (0.53 to 2.27)	0.807
	on ≥3 consecutive days	43/91 (47.3%)	21/57 (36.8%)	0.65 (0.33 to 1.31)	0.229	1.17 (0.51 to 2.66)	0.712	1.07 (0.45 to 2.53)	0.884
>12 mmol/L	at least once	113/209 (54.1%)	26/69 (37.7%)	0.60 (0.33 to 1.07)	0.084	1.23 (0.61 to 2.49)	0.566	1.24 (0.60 to 2.57)	0.554

Hyperglycaemia definition	Survival without moderate to severe NDD in hyperglycaemic infants		Crude OR (95% CI)	P value	aOR* (95% CI)	P value	aOR† (95% CI)	P value
	No insulin	Insulin						
on ≥2 consecutive days	50/93 (53.8%)	25/61 (41.0%)	0.69 (0.34 to 1.40)	0.306	0.93 (0.41 to 2.10)	0.864	0.96 (0.42 to 2.22)	0.924
on ≥3 consecutive days	15/37 (40.5%)	16/43 (37.2%)	0.87 (0.34 to 2.27)	0.775	0.95 (0.33 to 2.80)	0.931	0.92 (0.28 to 2.97)	0.883
at least once	79/154 (51.3%)	25/64 (39.1%)	0.68 (0.36 to 1.29)	0.242	1.10 (0.54 to 2.26)	0.796	1.04 (0.50 to 2.18)	0.919
>14 mmol/L on ≥2 consecutive days	26/51 (51.0%)	18/44 (40.9%)	0.79 (0.33 to 1.90)	0.600	0.93 (0.35 to 2.44)	0.873	0.96 (0.35 to 2.62)	0.929
on ≥3 consecutive days	4/16 (25.0%)	10/25 (40.0%)	2.03 (0.46 to 8.95)	0.340	0.42 (0.08 to 2.20)	0.292	0.26 (0.03 to 2.03)	0.190

Random intercepts were used to account for the clustering effects of twins/triplets and treating hospital.

* Adjusted for duration of hyperglycaemia (according to respective definition) during the first 28 postnatal days.

† Adjusted for gestational age at birth, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, and duration of hyperglycaemia (according to respective definition) during the first 28 postnatal days.

Online supplementary table 5. Univariable and multivariable generalised linear mixed models for the associations between neonatal hyperglycaemia (>12 and >14 mmol/L) and Wechsler Intelligence Scale for Children IV (WISC-IV) score at 6.5 years of age in children born extremely preterm.

Hyperglycaemia definition	Mean WISC-IV score		Univariable		Multivariable		
	No hyperglycaemia (n)	Hyperglycaemia (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores (95% CI)	P value	
>12 mmol/L	at least once	86.03 (177)	81.98 (178)	-4.05 (-7.04 to -1.06)	0.008	-1.07 (-4.10 to 1.96)	0.487
	on ≥ 2 consecutive days	85.24 (262)	80.50 (93)	-4.74 (-8.17 to -1.32)	0.007	-1.43 (-4.87 to 2.02)	0.417
	on ≥ 3 consecutive days	84.78 (314)	78.12 (41)	-6.66 (-11.21 to -2.11)	0.004	-2.07 (-6.57 to 2.44)	0.368
	Days with hyperglycaemia (per 1 day increase)	-	-	-0.78 (-1.24 to -0.32)	0.001	-0.22 (-0.72 to 0.28)	0.383
>14 mmol/L	at least once	86.14 (222)	80.45 (133)	-5.70 (-8.74 to -2.65)	<0.001	-2.88 (-5.97 to 0.20)	0.067
	on ≥ 2 consecutive days	84.91 (299)	79.12 (56)	-5.79 (-9.76 to -1.82)	0.004	-1.41 (-5.37 to 2.56)	0.486
	on ≥ 3 consecutive days	84.45 (334)	76.85 (21)	-7.60 (-13.81 to -1.40)	0.017	-4.13 (-10.21 to 1.96)	0.183

Hyperglycaemia definition	Mean WISC-IV score		Univariable	Multivariable		
	No hyperglycaemia (n)	Hyperglycaemia (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores (95% CI)	P value
Days with hyperglycaemia (per 1 day increase)	-	-	-1.00 (-1.63 to -0.37)	0.002	-0.40 (-1.05 to 0.24)	0.222

A random intercept was used to account for the clustering effect of twins/triplets. Adjusted for gestational age at birth, sex, occurrence of intraventricular haemorrhage grade 3-4 and/or periventricular leukomalacia, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, number of culture-verified sepsis episodes during the first 14 postnatal days, and educational status of the mother.

Online supplementary table 6. Univariable and multivariable generalised linear mixed models for the associations between insulin treatment in hyperglycaemic infants and Wechsler Intelligence Scale for Children IV (WISC-IV) score at 6.5 years of age in children born extremely preterm.

Hyperglycaemia definition		Mean WISC-IV score		Univariable		Multivariable			
		No insulin (n)	Insulin (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores* (95% CI)	P value	Difference in mean scores† (95% CI)	P value
>8 mmol/L	at least once	84.03 (265)	77.61 (45)	-6.43 (-11.01 to -1.85)	0.006	-0.80 (-6.18 to 4.57)	0.769	0.36 (-4.94 to 5.66)	0.894
	on ≥ 2 consecutive days	81.94 (180)	77.86 (45)	-4.08 (-8.85 to 0.69)	0.093	-0.73 (-6.18 to 4.73)	0.793	0.33 (-5.07 to 5.74)	0.904
	on ≥ 3 consecutive days	81.51 (106)	77.29 (41)	-4.22 (-9.13 to 0.69)	0.091	-1.84 (-7.48 to 3.81)	0.521	-0.08 (-5.77 to 5.61)	0.978
>10 mmol/L	at least once	83.34 (196)	77.73 (45)	-5.61 (-10.28 to -0.95)	0.019	-1.48 (-7.09 to 4.12)	0.602	0.30 (-5.31 to 5.91)	0.916
	on ≥ 2 consecutive days	82.26 (103)	78.60 (44)	-3.66 (-8.88 to 1.56)	0.168	-1.13 (-6.90 to 4.63)	0.699	0.43 (-5.52 to 6.38)	0.887
	on ≥ 3 consecutive days	81.67 (55)	75.53 (37)	-6.14 (-12.52 to 0.24)	0.059	-4.33 (-11.42 to 2.77)	0.229	-1.68 (-9.13 to 5.77)	0.654
>12 mmol/L	at least once	82.72 (134)	77.41 (44)	-5.31 (-10.34 to -0.28)	0.039	-3.65 (-9.86 to 2.56)	0.248	-2.46 (-8.63 to 3.70)	0.431
	on ≥ 2 consecutive days	81.89 (54)	80.16 (39)	-1.73 (-7.29 to 3.84)	0.540	0.30 (-6.14 to 6.73)	0.928	1.50 (-5.28 to 8.29)	0.661

Hyperglycaemia definition	Mean WISC-IV score		Univariable		Multivariable			
	No insulin (n)	Insulin (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores* (95% CI)	P value	Difference in mean scores† (95% CI)	P value
on ≥ 3 consecutive days	78.80 (16)	76.76 (25)	-2.04 (-10.20 to 6.12)	0.616	-1.80 (-11.23 to 7.62)	0.701	0.06 (-10.46 to 10.58)	0.991
at least once	81.38 (91)	77.86 (42)	-3.51 (-8.54 to 1.51)	0.169	-2.94 (-8.89 to 3.01)	0.330	-1.97 (-7.64 to 3.69)	0.492
>14 mmol/L on ≥ 2 consecutive days	81.16 (29)	77.90 (27)	-3.26 (-10.35 to 3.83)	0.360	-3.22 (-11.45 to 5.00)	0.435	-3.60 (-11.43 to 4.24)	0.361
on ≥ 3 consecutive days	70.00 (5)	77.94 (16)	7.94 (-4.64 to 20.52)	0.202	8.15 (-6.27 to 22.58)	0.251	16.90 (1.34 to 32.45)	0.035

Random intercepts were used to account for the clustering effects of twins/triplets and treating hospital.

* Adjusted for duration of hyperglycaemia (according to respective definition) during the first 28 postnatal days.

† Adjusted for gestational age at birth, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, and duration of hyperglycaemia (according to respective definition) during the first 28 postnatal days.

Online supplementary table 7. Univariable and multivariable generalised linear mixed models for the associations between neonatal hyperglycaemia (>12 and >14 mmol/L) and Motor Assessment Battery for Children 2 (MABC-2) score at 6.5 years of age in children born extremely preterm.

Hyperglycaemia definition	Mean MABC-2 score		Univariable		Multivariable		
	No hyperglycaemia (n)	Hyperglycaemia (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores (95% CI)	P value	
>12 mmol/L	at least once	70.01 (174)	61.37 (171)	-8.64 (-12.45 to -4.82)	<0.001	-4.88 (-8.81 to -0.95)	0.015
	on ≥ 2 consecutive days	67.95 (256)	59.07 (89)	-8.88 (-13.28 to -4.47)	<0.001	-5.84 (-10.29 to -1.40)	0.010
	on ≥ 3 consecutive days	66.65 (303)	58.62 (42)	-8.03 (-13.86 to -2.19)	0.007	-2.05 (-7.86 to 3.76)	0.489
	Days with hyperglycaemia (per 1 day increase)	-	-	-1.37 (-1.92 to -0.83)	<0.001	-0.93 (-1.52 to -0.34)	0.002
>14 mmol/L	at least once	69.40 (219)	59.28 (126)	-10.12 (-14.08 to -6.16)	<0.001	-6.90 (-10.95 to -2.85)	0.001
	on ≥ 2 consecutive days	66.93 (291)	58.86 (54)	-8.07 (-13.26 to -2.87)	0.002	-3.10 (-8.26 to 2.05)	0.237
	on ≥ 3 consecutive days	66.62 (324)	50.93 (21)	-15.69 (-23.46 to -7.93)	<0.001	-10.73 (-18.38 to -3.09)	0.006

Hyperglycaemia definition	Mean MABC-2 score		Univariable		Multivariable	
	No hyperglycaemia (n)	Hyperglycaemia (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores (95% CI)	P value
Days with hyperglycaemia (per 1 day increase)	-	-	-2.00 (-2.73 to -1.28)	<0.001	-1.47 (-2.21 to -0.72)	<0.001

A random intercept was used to account for the clustering effect of twins/triplets. Adjusted for gestational age at birth, sex, occurrence of intraventricular haemorrhage grade 3-4 and/or periventricular leukomalacia, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, number of culture-verified sepsis episodes during the first 14 postnatal days, educational status of the mother, and age at follow-up. MABC-2 total scores ≤ 75 and ≤ 68 are equivalent to ≤ 15 th centile (borderline motor impairment) and ≤ 5 th centile (developmental coordination disorder) in Swedish children, respectively.

Online supplementary table 8. Univariable and multivariable generalised linear mixed models for the associations between insulin treatment in hyperglycaemic infants and Movement Assessment Battery for Children (MABC-2) score at 6.5 years of age in children born extremely preterm.

Hyperglycaemia definition		Mean MABC-2 score		Univariable		Multivariable			
		No insulin (n)	Insulin (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores* (95% CI)	P value	Difference in mean scores† (95% CI)	P value
>8 mmol/L	at least once	66.21 (257)	54.44 (44)	-11.77 (-17.75 to -5.79)	<0.001	-5.66 (-12.79 to 1.47)	0.119	-3.75 (-10.92 to 3.42)	0.304
	on ≥2 consecutive days	64.76 (172)	55.07 (44)	-9.69 (-16.06 to -3.31)	0.003	-4.98 (-12.33 to 2.37)	0.183	-3.46 (-10.96 to 4.03)	0.363
	on ≥3 consecutive days	64.03 (101)	53.68 (40)	-10.36 (-17.32 to -3.39)	0.004	-7.85 (-16.18 to 0.48)	0.065	-5.49 (-14.66 to 3.67)	0.238
>10 mmol/L	at least once	65.08 (187)	54.54 (44)	-10.54 (-16.88 to -4.20)	0.001	-3.84 (-11.40 to 3.72)	0.318	-0.85 (-8.62 to 6.91)	0.829
	on ≥2 consecutive days	62.80 (99)	54.27 (43)	-8.53 (-15.31 to -1.74)	0.014	-3.96 (-11.70 to 3.79)	0.315	-1.62 (-9.95 to 6.72)	0.702
	on ≥3 consecutive days	62.27 (53)	53.26 (36)	-9.01 (-17.22 to -0.80)	0.032	-6.50 (-16.16 to 3.16)	0.185	-3.55 (-13.97 to 6.86)	0.499
>12 mmol/L	at least once	63.06 (128)	55.27 (43)	-7.79 (-14.61 to -0.96)	0.026	-3.41 (-11.77 to 4.96)	0.422	-0.83 (-9.36 to 7.70)	0.848
	on ≥2 consecutive days	62.59 (51)	55.31 (38)	-7.28 (-15.12 to 0.56)	0.068	-5.22 (-14.27 to 3.84)	0.255	-3.11 (-12.96 to 6.74)	0.532

Hyperglycaemia definition	Mean MABC-2 score		Univariable		Multivariable			
	No insulin (n)	Insulin (n)	Difference in mean scores (95% CI)	P value	Difference in mean scores* (95% CI)	P value	Difference in mean scores† (95% CI)	P value
on ≥ 3 consecutive days	67.80 (17)	53.72 (25)	-14.08 (-26.66 to -1.50)	0.029	-10.54 (-25.03 to 3.95)	0.149	-7.42 (-22.75 to 7.91)	0.332
at least once	60.50 (85)	55.17 (41)	-5.33 (-13.16 to 2.50)	0.180	1.47 (-6.94 to 9.87)	0.730	3.21 (-5.57 to 11.99)	0.470
>14 mmol/L on ≥ 2 consecutive days	65.00 (27)	53.83 (27)	-11.17 (-20.83 to -1.51)	0.024	-4.91 (-15.68 to 5.86)	0.364	-4.21 (-14.51 to 6.09)	0.414
on ≥ 3 consecutive days	63.80 (5)	48.94 (16)	-14.86 (-38.10 to 8.38)	0.197	-14.93 (-42.88 to 13.03)	0.276	-13.33 (-42.79 to 16.13)	0.346

Random intercepts were used to account for the clustering effects of twins/triplets and treating hospital.

* Adjusted for duration of hyperglycaemia (according to respective definition) during the first 28 postnatal days and age at follow-up.

† Adjusted for gestational age at birth, durations of mechanical ventilation and steroid treatments during the first 14 postnatal days, surgery due to patent ductus arteriosus or necrotising enterocolitis during the first 14 postnatal days, and duration of hyperglycaemia (according to respective definition) during the first 28 postnatal days.